§ 522.1055

- (iii) *Limitations*. Administer intramuscularly or subcutaneously. If no improvement is seen after 3 days, treatment should be discontinued and the diagnosis reevaluated. Treatment not to exceed 7 days. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- [43 FR 1942, Jan. 13, 1978, as amended at 48 FR 791, Jan. 7, 1983; 51 FR 15606, Apr. 25, 1986; 52 FR 7832, Mar. 13, 1987; 53 FR 40727, Oct. 18, 1988; 60 FR 29985, June 7, 1995; 61 FR 24441, May 15, 1996; 62 FR 45157, Aug. 26, 1997]

§522.1055 Gleptoferron injection.

- (a) Specifications. Each milliliter contains the equivalent of 200 milligrams of elemental iron as gleptoferron (complex of ferric hydroxide and dextran glucoheptonic acid), and 0.5 percent phenol as a preservative.
- (b) *Sponsor*. See 062408 in §510.600(c) of this chapter.
- (c) *Conditions of use.* It is used in baby pigs as follows:
- (1) For prevention of iron deficiency anemia, administer 200 milligrams of elemental iron intramuscularly on or before 3 days of age.
- (2) For treatment of iron deficiency anemia, administer 200 milligrams of elemental iron intramuscularly.
- $[45\ FR\ 61288,\ Sept.\ 16,\ 1980,\ as\ amended\ at\ 61\ FR\ 18672,\ Apr.\ 29,\ 1996]$

§ 522.1066 Glycopyrrolate injection.

- (a) *Specifications.* Each milliliter of aqueous solution contains 0.2 milligram of glycopyrrolate.
- (b) *Sponsor*. See No. 000031 in §510.600(c) of this chapter.
- (c) Conditions of use. (1) It is indicated as a preanesthetic agent in dogs and cats.
- (2) It is administered intravenously, intramuscularly, or subcutaneously in dogs and intramuscularly in cats at a dosage level of 5 micrograms per pound of body weight (0.25 milliliter per 10 pounds of body weight).
- (3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 21567, May 13, 1983]

§ 522.1077 Gonadorelin injectable.

(a) Specifications. Each milliliter sterile aqueous solution contains 50

micrograms of gonadorelin (as hydrochloride).

- (b) Sponsor. See No. 000856 in \$510.600(c) of this chapter.
- (c) Conditions of use in cattle—(1) Amount. 100 micrograms per cow intramuscularly.
- (2) *Indications for use.* For the treatment of cystic ovaries (ovarian follicular cysts) in cattle to reduce the time to first estrus.
- (3) *Limitations.* For intramuscular use only. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[54 FR 50235, Dec. 5, 1989]

§ 522.1078 Gonadorelin diacetate tetrahydrate injection

- (a) *Specifications.* The drug contains 50 micrograms of gonadorelin diacetate tetrahydrate in each milliliter of sterile solution.
- (b) *Sponsor*. See Nos. 050604 and 057926 in §510.600(c) of this chapter.
- (c) *Conditions of use.* It is used in dairy cows as follows:
 - (1) Amount. 100 micrograms per cow.
- (2) Indications for use. The drug is used for the treatment of ovarian cysts.
- (3) *Limitations.* Administer as a single intramuscular or intravenous injection. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- [43 FR 9804, Mar. 10, 1978, as amended at 45 FR 56798, Aug. 26, 1980; 61 FR 37682, July 19, 1996]

§522.1079 Serum gonadotropin and chorionic gonadotropin.

- (a) Specifications. Each dose consists of 400 international units (I.U.) serum gonadotropin and 200 I.U. chorionic gonadotropin as a freeze-dried powder to be reconstituted with 5 milliliters of sterile aqueous diluent.
- (b) *Sponsor*. See No. 057926 in §510.600(c) of this chapter.
- (c) Conditions of use in swine. (1) Amount. 400 I.U. serum gonadotropin with 200 I.U. chorionic gonadotropin per 5 milliliters dose per animal.
- (2) Indications for use. (i) Gilts. For induction of fertile estrus (heat) in healthy prepuberal (noncycling) gilts.