

800 North Capitol Street, NW., suite 700, Washington, DC.

(14) The stiffness is not less than 50 kiloPascals/second, as determined by a method entitled "Method for Measurement of the Stiffness of Olestra," dated December 19, 1995, which is incorporated by reference in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. Copies are available from the Office of Premarket Approval, Center for Food Safety and Applied Nutrition (HFS-200), Food and Drug Administration, 5100 Paint Branch Pkwy., College Park, MD 20740, or may be examined at the Center for Food Safety and Applied Nutrition's Library, 5100 Paint Branch Pkwy., College Park, MD 20740, or at the Office of the Federal Register, 800 North Capitol St. NW., suite 700, Washington, DC.

(c) Olestra may be used in place of fats and oils in prepackaged ready-to-eat savory (i.e., salty or piquant but not sweet) snacks. In such foods, the additive may be used in place of fats and oils for frying or baking, in dough conditioners, in sprays, in filling ingredients, or in flavors.

(d) To compensate for any interference with absorption of fat soluble vitamins, the following vitamins shall be added to foods containing olestra: 1.9 milligrams alpha-tocopherol equivalents per gram olestra; 51 retinol equivalents per gram olestra (as retinyl acetate or retinyl palmitate); 12 IU vitamin D per gram olestra; and 8 µg vitamin K₁ per gram olestra.

(e)(1) Vitamins A, D, E, and K present in foods as a result of the requirement in paragraph (d) of this section shall be declared in the listing of ingredients. Such vitamins shall not be considered in determining nutrient content for the nutritional label or for any nutrient claims, express or implied.

(i) An asterisk shall follow vitamins A, D, E, and K in the listing of ingredients;

(ii) The asterisk shall appear as a superscript following each vitamin;

(iii) Immediately following the ingredient list an asterisk and statement, "Dietarily insignificant" shall appear prominently and conspicuously as specified in §101.2(c) of this chapter;

(2) Olestra shall not be considered as a source of fat or calories for purposes of §§101.9 and 101.13 of this chapter.

(f) Consistent with its obligation to monitor the safety of all additives in the food supply, including olestra, the Food and Drug Administration will review and evaluate all data and information bearing on the safety of olestra received by the agency after the effective date of this regulation, and will present such data, information, and evaluation to the agency's Food Advisory Committee within 30 months of the effective date of this regulation. The purpose of such presentation will be to receive advice from the Committee on whether there continues to be reasonable certainty that use of olestra in compliance with this regulation is not harmful. The agency will hold such additional Food Advisory Committee meetings on olestra as the agency determines, in its discretion, to be necessary. Based upon the results of this entire process, the FDA will initiate any appropriate regulatory proceedings.

[61 FR 3171, Jan. 30, 1996; 61 FR 11546, Mar. 21, 1996, as amended at 68 FR 46402, Aug. 5, 2003]

§ 172.868 Ethyl cellulose.

The food additive ethyl cellulose may be safely used in food in accordance with the following prescribed conditions:

(a) The food additive is a cellulose ether containing ethoxy (OC₂H₅) groups attached by an ether linkage and containing on an anhydrous basis not more than 2.6 ethoxy groups per anhydroglucose unit.

(b) It is used or intended for use as follows:

(1) As a binder and filler in dry vitamin preparations.

(2) As a component of protective coatings for vitamin and mineral tablets.

(3) As a fixative in flavoring compounds.

§ 172.869 Sucrose oligoesters.

Sucrose oligoesters, as identified in this section, may be safely used in accordance with the following conditions:

(a) Sucrose oligoesters consist of mixtures of sucrose fatty acid esters with an average degree of

§ 172.869

21 CFR Ch. I (4-1-04 Edition)

esterification ranging from four to seven. It is produced by interesterification of sucrose with methyl esters of fatty acids derived from edible fats and oils (including hydrogenated fats and oils). The only solvents which may be used in the preparation of sucrose oligoesters are dimethyl sulfoxide, isobutyl alcohol, and those solvents generally recognized as safe in food.

(b) Sucrose oligoesters meet the specifications in the methods listed in the table in this paragraph. The methods

cited for determining compliance with each specification are incorporated by reference, in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. Copies of the methods may be examined at the Center for Food Safety and Applied Nutrition's Library, room 1C-100, 5100 Paint Branch Pkwy., College Park, MD 20740, or at the Office of the Federal Register, 800 North Capitol St. NW., suite 700, Washington, DC. Copies of the methods are available from the sources listed in the table in this paragraph:

Specification	Limit	Method Cited	Source for Obtaining Method
(1) Sucrose esters	Not less than 90%	"Method for Analyzing the Purity of Sucrose Fatty Acid Esters," issued by Mitsubishi Chemical Corp., June 17, 1998.	Office of Food Additive Safety, Center for Food Safety and Applied Nutrition (HFS-200), Food and Drug Administration, 5100 Paint Branch Pkwy., College Park, MD 20740.
(2) Mono-, di-, and tri-esters ...	Not more than 45%	"Method for Measuring the Ester Distribution of Sucrose Oligoesters," issued by Mitsubishi Chemical Corp., June 17, 1998.	Do.
(3) Tetra-, penta-, hexa-, and hepta-esters.	Not less than 50%	Do.	Do.
(4) Octa-esters	Not more than 40%	Do.	Do.
(5) Free Sucrose	Not more than 0.5%	"Free Sucrose Method," issued by Mitsubishi Chemical Corp., June 17, 1998.	Do.
(6) Acid Value	Not more than 4.0	"Acid Value," Appendix VII, Method I (Commercial Fatty Acids), in the <i>Food Chemicals Codex</i> , 4th ed. (1996), p. 820.	National Academy Press, 2101 Constitution Ave. NW, Washington, DC 20418 (Internet: http://www.nap.edu).
(7) Residue on Ignition	Not more than 0.7%	"Residue on Ignition, Appendix IIC, Method I, in the <i>Food Chemicals Codex</i> , 4th ed. (1996), pp. 751-752, (using a 1-gram sample).	Do.
(8) Residual Methanol	Not more than 10 milligrams/kilogram.	Method listed in the monograph for "Sucrose Fatty Acid Esters" in the First Supplement to the 4th ed. of the <i>Food Chemicals Codex</i> (1997), pp. 44-45.	Do.
(9) Residual Dimethyl Sulfoxide.	Not more than 2.0 milligrams/kilogram.	Do.	Do.
(10) Residual Isobutyl Alcohol	Not more than 10 milligrams/kilogram.	Do.	Do.
(11) Lead	Not more than 1.0 milligram/kilogram.	"Atomic Absorption Spectrophotometric Graphite Furnace Method," Method I, in the <i>Food Chemicals Codex</i> , 4th ed. (1996), pp. 763-765.	Do.

(c) The additive is used as an emulsifier (as defined in §170.3(o)(8) of this chapter) or stabilizer (as defined in §170.3(o)(28) of this chapter) in chocolate and in butter-substitute spreads, at a level not to exceed 2.0 percent; except that the additive may not be used in a standardized food unless permitted by the standard of identity.

[68 FR 50072, Aug. 20, 2003]

§ 172.870 Hydroxypropyl cellulose.

The food additive hydroxypropyl cellulose may be safely used in food, except standardized foods that do not provide for such use, in accordance with the following prescribed conditions:

(a) The additive consists of one of the following:

(1) A cellulose ether containing propylene glycol groups attached by an ether linkage which contains, on an anhydrous basis, not more than 4.6 hydroxypropyl groups per anhydroglucose unit. The additive has a minimum viscosity of 145 centipoises for 10 percent by weight aqueous solution at 25 °C.

(2) A cellulose ether containing propylene glycol groups attached by an ether linkage having a hydroxypropoxy (OC₃H₆OH) content of 5 to 16 percent weight in weight (w/w) on an anhydrous basis, i.e., 0.1 to 0.4 hydroxypropyl groups per anhydroglucose unit. The common name for this form of the additive is low substituted hydroxypropyl cellulose.

(b) The additive is used or intended for use as follows:

(1) The additive identified in paragraph (a)(1) of this section is used or intended for use as an emulsifier, film former, protective colloid, stabilizer, suspending agent, or thickener, in accordance with good manufacturing practice.

(2) The additive identified in paragraph (a)(2) of this section is used or intended for use as a binder and disintegrator in tablets or wafers containing dietary supplements of vitamins and/or minerals. The additive is used in accordance with good manufacturing practice.

[46 FR 50065, Oct. 9, 1981]

§ 172.872 Methyl ethyl cellulose.

The food additive methyl ethyl cellulose may be safely used in food in accordance with the following prescribed conditions.

(a) The additive is a cellulose ether having the general formula $[C_6H_{(10-x-y)}O_5(CH_3)_x(C_2H_5)_y]_n$, where x is the number of methyl groups and y is the number of ethyl groups. The average value of x is 0.3 and the average value of y is 0.7.

(b) The additive meets the following specifications:

(1) The methoxy content shall be not less than 3.5 percent and not more than 6.5 percent, calculated as OCH₃, and the ethoxy content shall be not less than 14.5 percent and not more than 19 percent, calculated as OC₂H₅, both measured on the dry sample.

(2) The viscosity of an aqueous solution, 2.5 grams of the material in 100 milliliters of water, at 20 °C, is 20 to 60 centipoises.

(3) The ash content on a dry basis has a maximum of 0.6 percent.

(c) The food additive is used as an aerating, emulsifying, and foaming agent, in an amount not in excess of that reasonably required to produce its intended effect.

§ 172.874 Hydroxypropyl methylcellulose.

The food additive hydroxypropyl methylcellulose (CAS Reg. No. 9004-65-3) may be safely used in food, except in standardized foods which do not provide for such use if:

(a) The additive complies with the definition and specifications prescribed in the National Formulary, 12th edition.

(b) It is used or intended for use as an emulsifier, film former, protective colloid, stabilizer, suspending agent, or thickener, in accordance with good manufacturing practice.

(c) To insure safe use of the additive, the container of the additive, in addition to being labeled as required by the general provisions of the act, shall be accompanied by labeling which contains adequate directions for use to provide a final product that complies