## § 520.2345f

- (iii) Limitations. Administer orally; continue treatment until symptoms have subsided and the temperature is normal for 48 hours; not for use in animals which are raised for food production; Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- (iv) National Academy of Sciences/National Research Council (NAS/NRC) status. These conditions were NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by §514.111 of this chapter, but may require bioequivalency and safety information.
- (2) Dogs and cats—(i) Amount. 25 milligrams per pound of body weight per day in divided doses every 6 hours.
- (ii) *Indications for use*. Treatment of infections caused by organisms susceptible to tetracycline hydrochloride, such as bacterial gastroenteritis due to *E. coli* and urinary tract infections due to *Staphylococcus* spp. and *E. coli*.
- (iii) Limitations. Administer orally; continue treatment until the temperature has been normal for 48 hours; not for use in food-producing animals; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37329, Aug. 18, 1992]

## § 520.2345f Tetracycline phosphate complex and sodium novobiocin capsules.

- (a) *Specifications*. Each capsule contains the equivalent of 60 milligrams of tetracycline hydrochloride and 60 milligrams of novobiocin.
- (b) *Sponsor*. No. 000009 in §510.600(c) of this chapter.
- (c) Conditions of use. Dogs—(1) Amount. 10 milligrams of each antibiotic per pound of body weight (1 capsule for each 6 pounds) every 12 hours.
- (2) Indications for use. Treatment of acute or chronic canine respiratory infections such as tonsillitis, bronchitis, and tracheobronchitis when caused by pathogens susceptible to tetracycline and/or novobiocin, such as Staphylococcus spp. and Escherichia coli.
- (3) Limitations. Continue treatment for at least 48 hours after the temperature has returned to normal and all evidence of infection has disappeared. As with all antibiotics, appropriate in vitro culturing and susceptibility tests

of samples taken before treatment should be conducted. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37329, Aug. 18, 1992]

## § 520.2345g Tetracycline hydrochloride and sodium novobiocin tablets.

- (a) Specifications. Each tablet contains the equivalent of 60 milligrams of tetracycline hydrochloride and 60 milligrams of novobiocin, or 180 milligrams of tetracycline hydrochloride and 180 milligrams of novobiocin.
- (b) Sponsor. No. 000009 in \$510.600(c) of this chapter.
- (c) Conditions of use. Dogs—(1) Amount. 10 milligrams of each antibiotic per pound of body weight (one single-strength tablet for each 6 pounds or one triple-strength tablet for each 18 pounds).
- (2) Indications for use. Treatment of acute or chronic canine respiratory infections such as tonsillitis, bronchitis, and tracheobronchitis when caused by pathogens susceptible to tetracycline and/or novobiocin, such as Staphylococcus spp. and Escherichia coli.
- (3) Limitations. Continue treatment for at least 48 hours after the temperature has returned to normal and all evidence of infection has disappeared. As with all antibiotics, appropriate in vitro culturing and susceptibility tests of samples taken before treatment should be conducted. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37329, Aug. 18, 1992]

## § 520.2345h Tetracycline hydrochloride, sodium novobiocin, and prednisolone tablets.

- (a) Specifications. Each tablet contains the equivalent of 60 milligrams of tetracycline hydrochloride, 60 milligrams of novobiocin, and 1.5 milligrams of prednisolone or 180 milligrams of tetracycline hydrochloride, 180 milligrams of novobiocin, and 4.5 milligrams of prednisolone.
- (b) *Sponsor*. See No. 000009 in §510.600(c) of this chapter.
- (c) Conditions of use. Dogs—(1) Amount. 10 milligrams of each antibiotic and 0.25 milligram of prednisolone per pound of body weight (one