

§ 522.778

21 CFR Ch. I (4–1–10 Edition)

§ 522.778 Doxycycline hyclate.

(a) *Specifications.* Doxycycline hyclate solution contains 8.5 percent doxycycline activity. A syringe of *N*-methyl-2-pyrrolidone and poly (DL-lactide) mixed with a syringe of doxycycline produces 0.5 milliliter of solution.

(b) *Sponsor.* See 000009 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(1) *Dogs*—(i) *Amount.* Apply subgingivally to periodontal pocket(s) of affected teeth.

(ii) *Indications for use.* For treatment and control of periodontal disease.

(iii) *Limitations.* Do not use in dogs less than 1-year old. Use of tetracyclines during tooth development has been associated with permanent discoloration of teeth. Do not use in pregnant bitches. Use in breeding dogs has not been evaluated. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[63 FR 8349, Feb. 19, 1998, as amended at 65 FR 45878, July 26, 2000]

§ 522.784 Doxylamine succinate injection.

(a) *Specifications.* Each milliliter of the drug contains 11.36 mg of doxylamine succinate.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in conditions in which antihistaminic therapy may be expected to alleviate some signs of disease in horses, dogs, and cats.¹

(2) It is administered to horses at a dosage level of 25 mg per hundred pounds of body weight. It is administered to dogs and cats at a dosage level of 0.5 to 1 mg per pound of body weight. Doses may be repeated at 8 to 12 hours, if necessary, to produce desired effect. Intravenous route is not recommended for dogs and cats and should be injected slowly in horses. Intramuscular and subcutaneous administration should be by divided injection sites.¹

(3) Not for use in horses intended for food.¹

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13858, Mar. 27, 1975, as amended at 42 FR 60140, Nov. 25, 1977; 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 522.800 Droperidol and fentanyl citrate injection.

(a) *Specifications.* Droperidol and fentanyl citrate injection is a sterile solution containing 20 milligrams of droperidol and 0.4 milligram of fentanyl citrate per cubic centimeter.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in dogs as an analgesic and tranquilizer and for general anesthesia.

(2) It is administered as follows:

(i) For analgesia and tranquilization administer according to response desired, as follows:

(a) Intramuscularly at the rate of 1 cubic centimeter per 15 to 20 pounds of body weight in conjunction with atropine sulfate administered at the rate of 0.02 milligram per pound of body weight, or

(b) Intravenously at the rate of 1 cubic centimeter per 25 to 60 pounds of body weight in conjunction with atropine sulfate administered at the rate of 0.02 milligram per pound of body weight.

(ii) For general anesthesia administer according to response desired, as follows:

(a) Intramuscularly at the rate of 1 cubic centimeter per 40 pounds of body weight in conjunction with atropine sulfate administered at the rate of 0.02 milligram per pound of body weight and followed in 10 minutes by an intravenous administration of sodium pentobarbital at the rate of 3 milligrams per pound of body weight, or

(b) Intravenously at the rate of 1 cubic centimeter per 25 to 60 pounds of body weight in conjunction with atropine sulfate administered at the rate of 0.02 milligram per pound of body weight and followed within 15 seconds by an intravenous administration of sodium pentobarbital at the rate of 3 milligrams per pound of body weight.