

(c) *Conditions of use in dogs and cats*—
(1) *Amount*. Apply once daily by rubbing into or spraying a light covering on the infected site and the immediate surrounding vicinity. Continue treatment for 2 to 4 weeks until infection is completely eradicated as determined by appropriate laboratory examination.

(2) *Indications for use*. For topical treatment of infections caused by *Microsporium canis*, *Microsporium gypseum*, and *Trichophyton mentagrophytes*.

(3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[71 FR 13542, Mar. 16, 2006]

§ 524.1445 Miconazole, polymixin B, and prednisolone suspension.

(a) *Specifications*. Each milliliter of suspension contains 23 milligrams (mg) miconazole nitrate, 0.5293 mg polymixin B sulfate, and 5 mg prednisolone acetate.

(b) *Sponsor*. See No. 012578 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs*—(1) *Amount*. Instill five drops in the ear canal twice daily for 7 consecutive days.

(2) *Indications for use*. For the treatment of canine otitis externa associated with susceptible strains of yeast (*Malassezia pachydermatis*) and bacteria (*Staphylococcus pseudintermedius*).

(3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[75 FR 4693, Jan. 29, 2010]

§ 524.1446 Milbemycin oxime solution.

(a) *Specifications*. Each tube contains 0.25 milliliter of a 0.1 percent solution of milbemycin oxime.

(b) *Sponsor*. See No. 058198 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. One tube administered topically into each external ear canal.

(2) *Indications for use*. For the treatment of ear mite (*Otodectes cynotis*) infestations in cats and kittens 4 weeks of age and older. Effectiveness is maintained throughout the life cycle of the ear mite.

(3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[65 FR 13905, Mar. 15, 2000, as amended at 66 FR 13849, Mar. 8, 2001]

§ 524.1451 Moxidectin.

(a) *Specifications*. Each milliliter contains 5 milligrams of moxidectin (0.5 percent solution).

(b) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.426 of this chapter.

(d) *Special considerations*. See § 500.25 of this chapter.

(e) *Conditions of use*—(1) *Amount*. 0.5 milligrams moxidectin per kilogram (2.2 pounds) of body weight.

(2) *Indications for use*. Beef and dairy cattle: For treatment and control of internal and external parasites: gastrointestinal roundworms (*Ostertagia ostertagi* (adult and L4, including inhibited larvae), *Haemonchus placei* (adult and L4), *Trichostrongylus axei* (adult and L4), *T. colubriformis* (adult and L4), *Cooperia oncophora* (adult and L4), *C. pectinata* (adult), *C. punctata* (adult and L4), *C. spatulata* (adult), *C. surnabada* (adult and L4), *Bunostomum phlebotomum* (adult), *Oesophagostomum radiatum* (adult and L4), *Nematodirus helvetianus* (adult and L4)); lungworms (*Dictyocaulus viviparus*, adult and L4); cattle grubs (*Hypoderma bovis*, *H. lineatum*); mites (*Chorioptes bovis*, *Psoroptes ovis* (*P. communis* var. *bovis*)); lice (*Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*, *Bovicola* (*Damalinea*) *bovis*); and horn flies (*Haematobia irritans*). To control infections and to protect from reinfection with *H. placei* for 14 days after treatment, *O. radiatum* and *O. ostertagi* for 28 days after treatment, and *D. viviparus* for 42 days after treatment.

(3) *Limitations*. A withdrawal period has not been established for this product on preruminating calves. Do not use on calves to be processed for veal.

[63 FR 14036, Mar. 24, 1998, as amended at 65 FR 36617, June 9, 2000; 66 FR 46370, Sept. 5, 2001]