

**Food and Drug Administration, HHS**

**§556.515**

in milk from dairy animals and 1 part per million in the uncooked edible tissues of cattle, chickens, turkeys, and ducks.

[47 FR 18590, Apr. 30, 1982]

**§556.470 Nystatin.**

A tolerance of zero is established for residues of nystatin in or on eggs and the uncooked edible tissues of swine and poultry.

**§556.480 Oleandomycin.**

Tolerances are established for negligible residues of oleandomycin in uncooked edible tissues of chickens, turkeys, and swine at 0.15 part per million.

**§556.490 Ormetoprim.**

(a) [Reserved]

(b) *Tolerances.* A tolerance of 0.1 part per million (ppm) is established for negligible residues of ormetoprim in uncooked edible tissues of chickens, turkeys, ducks, salmonids, catfish, and chukar partridges.

[64 FR 26672, May 17, 1999]

**§556.495 Oxfendazole.**

*Cattle:* A tolerance is established for total oxfendazole residues in edible cattle tissues based on a marker residue concentration of 0.8 part per million (ppm) fenbendazole in the target liver tissue. A fenbendazole concentration of 0.8 ppm in liver corresponds to a total safe concentration of oxfendazole residues of 1.7 ppm in liver. The safe concentrations of total oxfendazole residues in other uncooked edible cattle tissues are: muscle, 0.84 ppm; kidney, 2.5 ppm; and fat, 3.3 ppm. A tolerance refers to the concentration of marker residue in the target tissue selected to monitor for total drug residue in the target animal. A safe concentration is the total residue considered safe in edible tissue.

[55 FR 46943, Nov. 8, 1990]

**§556.500 Oxytetracycline.**

(a) *Acceptable daily intake (ADI).* The ADI for total tetracycline residues (chlortetracycline, oxytetracycline, and tetracycline) is 25 micrograms per kilogram of body weight per day.

(b) *Beef cattle, dairy cattle, calves, swine, sheep, chickens, turkeys, finfish, and lobster.* Tolerances are established for the sum of residues of the tetracyclines including chlortetracycline, oxytetracycline, and tetracycline, in tissues and milk as follows:

(1) 2 parts per million (ppm) in muscle.

(2) 6 ppm in liver.

(3) 12 ppm in fat and kidney.

(4) 0.3 ppm in milk.

[63 FR 57246, Oct. 27, 1998, as amended at 66 FR 46370, Sept. 5, 2001; 69 FR 6557, Feb. 11, 2004]

**§556.510 Penicillin.**

Tolerances are established for residues of penicillin and the salts of penicillin in food as follows:

(a) 0.05 part per million (negligible residue) in the uncooked edible tissues of cattle.

(b) Zero in the uncooked edible tissues of chickens, pheasants, quail, swine, and sheep; in eggs; and in milk or in any processed food in which such milk has been used.

(c) 0.01 part per million in the uncooked edible tissues of turkeys.

[40 FR 13942, Mar. 27, 1975, as amended at 43 FR 32749, July 28, 1978]

**§556.513 Piperazine.**

A tolerance of 0.1 part per million piperazine base is established for edible tissues of poultry and swine.

[64 FR 23019, Apr. 29, 1999]

**§556.515 Pirlimycin.**

(a) *Acceptable daily intake (ADI).* The ADI for total residues of pirlimycin is 0.01 milligrams per kilogram of body weight per day.

(b) *Tolerances*—(1) *Cattle*—(i) *Liver (the target tissue).* The tolerance for parent pirlimycin (the marker residue) is 0.5 part per million (ppm).

(ii) *Muscle.* The tolerance for parent pirlimycin (the marker residue) is 0.3 ppm.

(iii) *Milk.* The tolerance for parent pirlimycin (the marker residue in cattle milk) is 0.4 ppm.

(2) [Reserved]

[65 FR 61091, Oct. 16, 2000]