

Food and Drug Administration, HHS

§ 640.21

in § 610.40 of this chapter and § 640.5 (a), (b), and (c).

[53 FR 117, Jan. 5, 1988, as amended at 66 FR 31165, June 11, 2001]

§ 640.15 Segments for testing.

Segments collected in integral tubing shall meet the following standards:

(a) One or more segments shall be provided with each unit of Whole Blood or Red Blood Cells when issued or re-issued.

(b) Before they are filled, all segments shall be marked or identified so as to relate them to the donor of that unit of red cells.

(c) All segments accompanying a unit of Red Blood Cells shall be filled at the time the blood is collected or at the time the final product is prepared.

[66 FR 40890, Aug. 6, 2001]

§ 640.16 Processing.

(a) *Separation.* Within the timeframe specified in the directions for use for the blood collecting, processing, and storage system used, Red Blood Cells may be prepared either by centrifugation, done in a manner that will not tend to increase the temperature of the blood, or by normal undisturbed sedimentation. A portion of the plasma sufficient to insure optimal cell preservation shall be left with the red cells except when a cryoprotective substance or additive solution is added for prolonged storage.

(b) *Sterile system.* All surfaces that come in contact with the red cells shall be sterile and pyrogen-free.

(c) *Final containers.* Final containers used for Red Blood Cells shall be the original blood containers unless the method of processing requires a different container. The final container shall meet the requirements for blood containers prescribed in § 640.2(c). At the time of filing, if a different container is used, it shall be marked or identified by number or other symbol so as to relate it to the donor of that unit of red cells.

[38 FR 32089, Nov. 20, 1973, as amended at 43 FR 34460, Aug. 4, 1978; 50 FR 4139, Jan. 29, 1985; 64 FR 45372, Aug. 19, 1999; 66 FR 1836, Jan. 10, 2001; 66 FR 40890, Aug. 6, 2001]

§ 640.17 Modifications for specific products.

Red Blood Cells Frozen: A cryophylactic substance may be added to the Red Blood Cells for extended manufacturers' storage at -65° C or colder, provided the manufacturer submits data considered by the Director, Center for Biologics Evaluation and Research, as adequately demonstrating through in vivo cell survival and other appropriate tests that the addition of the substance, the materials used and the processing methods results in a final product that meets the required standards of safety, purity, and potency for Red Blood Cells, and that the frozen product will maintain those properties for the prescribed dating period. Section 640.11 (a) and (b) do not apply while a cryophylactic substance is present.

[38 FR 32089, Nov. 20, 1973, as amended at 41 FR 18292, May 3, 1976; 49 FR 23834, June 8, 1984; 50 FR 4139, Jan. 29, 1985; 55 FR 11013, Mar. 26, 1990; 63 FR 16685, Apr. 6, 1998]

Subpart C—Platelets

§ 640.20 Platelets.

(a) *Proper name and definition.* The proper name of this product shall be Platelets. The product is defined as platelets collected from one unit of blood and resuspended in an appropriate volume of original plasma, as prescribed in § 640.24(d).

(b) *Source.* The source material for Platelets is plasma which may be obtained by whole blood collection or by plateletpheresis.

[40 FR 4304, Jan. 29, 1975, as amended at 47 FR 49021, Oct. 29, 1982; 50 FR 4139, Jan. 29, 1985; 72 FR 45887, Aug. 16, 2007]

§ 640.21 Suitability of donors.

(a) Whole blood donors shall meet the criteria for suitability prescribed in § 640.3.

(b) [Reserved]

(c) Plateletpheresis donors must meet the criteria for suitability as prescribed in §§ 640.3 and 640.63(c)(6) or as described in an approved biologics license application (BLA) or an approved

§ 640.22

supplement to a BLA. Informed consent must be obtained as prescribed in § 640.61.

[40 FR 4304, Jan. 29, 1975, as amended at 49 FR 23834, June 8, 1984; 64 FR 56453, Oct. 20, 1999; 72 FR 45887, Aug. 16, 2007]

§ 640.22 Collection of source material.

(a) Whole blood used as the source of Platelets shall be collected as prescribed in § 640.4.

(b) [Reserved]

(c) If plateletpheresis is used, the procedure for collection must be as prescribed in §§ 640.62, 640.64 (except paragraph (c)), and 640.65, or as described in an approved biologics license application (BLA) or an approved supplement to a BLA.

(d) The phlebotomy shall be performed by a single uninterrupted venipuncture with minimal damage to, and minimal manipulation of, the donor's tissue.

[40 FR 4304, Jan. 29, 1975, as amended at 45 FR 27927, Apr. 25, 1980; 49 FR 23834, June 8, 1984; 50 FR 4139, Jan. 29, 1985; 55 FR 11013, Mar. 26, 1990; 59 FR 49351, Sept. 28, 1994; 64 FR 45372, Aug. 19, 1999; 64 FR 56453, Oct. 20, 1999; 72 FR 45887, Aug. 16, 2007]

§ 640.23 Testing the blood.

(a) Blood from which plasma is separated for the preparation of Platelets shall be tested as prescribed in § 610.40 of this chapter and § 640.5 (a), (b), and (c).

(b) The tests shall be performed on a sample of blood collected at the time of collecting the source blood, and such sample container shall be labeled with the donor's number before the container is filled.

[40 FR 4304, Jan. 29, 1975, as amended at 50 FR 4139, Jan. 29, 1985; 53 FR 117, Jan. 5, 1988; 64 FR 45372, Aug. 19, 1999; 66 FR 1836, Jan. 10, 2001; 66 FR 31165, June 11, 2001]

§ 640.24 Processing.

(a) Separation of plasma and platelets and resuspension of the platelets must be in a closed system. Platelets must not be pooled during processing unless the platelets are pooled as specified in the directions for use for the blood collecting, processing, and storage system approved for such use by the Director, Center for Biologics Evaluation and Research.

21 CFR Ch. I (4–1–10 Edition)

(b) Immediately after collection, the whole blood or plasma shall be held in storage between 20 and 24 °C unless it must be transported from the collection center to the processing laboratory. During such transport, all reasonable methods shall be used to maintain the temperature as close as possible to a range between 20 and 24 °C until it arrives at the processing laboratory where it shall be held between 20 and 24 °C until the platelets are separated. The platelet concentrate shall be separated within 4 hours or within the timeframe specified in the directions for use for the blood collecting, processing, and storage system.

(c) The time and speed of centrifugation must have been demonstrated to produce an unclumped product, without visible hemolysis, that yields a count of not less than 5.5×10^{10} platelets per unit in at least 75 percent of the units tested.

(d) The volume of original plasma used for resuspension of the platelets shall be determined by the maintenance of a pH of not less than 6.2 during the storage period. The pH shall be measured on a sample of platelets which has been stored for the maximum dating period at the selected storage temperature. One of the following storage temperatures shall be used continuously:

(1) 20 to 24 °C.

(2) 1 to 6 °C.

(e) Final containers used for Platelets shall be colorless and transparent to permit visual inspection of the contents; any closure shall maintain a hermetic seal and prevent contamination of the contents. The container material shall not interact with the contents, under the customary conditions of storage and use, in such a manner as to have an adverse effect upon the safety, purity, potency, or efficacy of the product. At the time of filling, the final container shall be marked or identified by number so as to relate it to the donor.

[40 FR 4304, Jan. 29, 1975, as amended at 42 FR 10983, Feb. 25, 1977; 47 FR 49021, Oct. 29, 1982; 50 FR 4139, Jan. 29, 1985; 63 FR 16685, Apr. 6, 1998; 64 FR 45372, Aug. 19, 1999; 66 FR 1836, Jan. 10, 2001; 66 FR 40890, Aug. 6, 2001; 72 FR 45887, Aug. 16, 2007; 73 FR 7464, Feb. 8, 2008]