

§ 520.2150b

§ 520.2150b Stanozolol chewable tablets.

(a) *Specifications.* Each chewable tablet contains 2 milligrams of stanozolol.

(b) *Sponsor.* No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) Used as an anabolic steroid treatment in dogs.

(2) Administered orally to small breeds of dogs, ½ to 1 tablet twice daily for several weeks; to large breeds of dogs, 1 to 2 tablets twice daily for several weeks.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 38114, Sept. 20, 1985, as amended at 55 FR 23076, June 6, 1990]

§ 520.2158 Streptomycin/dihydrostreptomycin oral dosage forms.

§ 520.2158a Streptomycin sulfate oral solution.

(a) *Specifications.* Solution containing 25 percent streptomycin sulfate.

(b) *Sponsor.* See Nos. 033008 and 055462 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.610 of this chapter.

(d) *Conditions of use.* Use in drinking water as follows:

(1) *Calves and swine*—(i) *Amount.* 10 to 15 milligrams per pound (mg/pound) of body weight (1.0 to 1.5 grams per gallon).

(ii) *Indications for use.* Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella* spp. susceptible to streptomycin.

(iii) *Limitations.* Calves: Do not administer for more than 5 days. Swine: Do not administer for more than 4 days. Prepare fresh solution daily. Calves: Withdraw 2 days before slaughter. As sole source of streptomycin. Warning: Certain strains of bacteria may develop a tolerance for streptomycin. Consult a veterinarian or animal pathologist for diagnosis.

(2) *Chickens*—(i) *Amount.* 10 to 15 mg/pound of body weight (0.6 to 0.9 grams per gallon).

(ii) *Indications for use.* Treatment of nonspecific infectious enteritis caused by organisms susceptible to streptomycin.

(iii) *Limitations.* Chickens: Do not administer for more than 5 days. With-

21 CFR Ch. I (4–1–11 Edition)

draw 4 days before slaughter. Do not administer to chickens producing eggs for human consumption. Prepare fresh solution daily. As sole source of streptomycin. Warning: Certain strains of bacteria may develop a tolerance for streptomycin. Consult a veterinarian or animal pathologist for diagnosis.

[57 FR 37327, Aug. 18, 1992, as amended at 58 FR 47211, Sept. 8, 1993; 63 FR 51821, Sept. 29, 1998]

§ 520.2158b Dihydrostreptomycin tablets.

(a) *Specifications.* Each tablet contains 37.5 milligrams dihydrostreptomycin (as the sulfate) with 375 milligrams chlorhexidine dihydrochloride.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See §§ 556.120 and 556.200 of this chapter.

(d) *Conditions of use.* *Calves*—(1) *Amount.* 150 milligrams of dihydrostreptomycin and 1.5 grams of chlorhexidine dihydrochloride per 100 pounds of body weight per day.

(2) *Indications for use.* Treatment of bacterial scours in calves.

(3) *Limitations.* Administer orally once a day for 5 days; withdraw 3 days before slaughter.

[57 FR 37327, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992]

§ 520.2158c Dihydrostreptomycin oral suspension.

(a) *Specifications.* Each milliliter contains 1.25 milligrams dihydrostreptomycin (as the sulfate) with 12.5 milligrams chlorhexidine dihydrochloride.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See §§ 556.120 and 556.200 of this chapter.

(d) *Conditions of use.* *Calves*—(1) *Amount.* 150 milligrams of dihydrostreptomycin and 1.5 grams of chlorhexidine dihydrochloride per 100 pounds of body weight per day.

(2) *Indications for use.* Treatment of bacterial scours in calves.

(3) *Limitations.* Administer orally once a day for 5 days; withdraw 3 days before slaughter.

[57 FR 37327, Aug. 18, 1992]