

§ 522.2690

21 CFR Ch. I (4-1-11 Edition)

month of age or after weaning in heifers intended for reproduction. Safety and effectiveness have not been established in veal calves. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

(2) *Feedlot lambs*—(i) *Amount.* 12 mg zeranol (one implant consisting of 1 pellet containing 12 mg zeranol) per implant dose.

(ii) *Indications for use.* For increased rate of weight gain and improved feed conversion.

(iii) *Limitations.* Implant subcutaneously in ear only. Do not use in breeding animals. Do not implant animals within 40 days of slaughter. Safety and effectiveness have not been established in veal calves. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

(3) *Steers fed in confinement for slaughter*—(i) *Amount.* 72 mg zeranol (one implant consisting of 6 pellets, each pellet containing 12 mg zeranol) per implant dose.

(ii) *Indications for use.* For increased rate of weight gain and improved feed efficiency.

(iii) *Limitations.* Implant subcutaneously in ear only. Safety and effectiveness have not been established in veal calves. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

(4) *Pasture cattle (slaughter, stocker, feeder steers, and heifers)*—(i) *Amount.* 138 mg zeranol (one implant consisting of 7 pellets, each of 6 pellets containing 20 mg zeranol and a seventh pellet containing 18 mg zeranol) per implant dose.

(ii) *Indications for use.* For increased rate of weight gain.

(iii) *Limitations.* Implant subcutaneously in ear only. Safety and effectiveness have not been established in veal calves. A withdrawal period has not been established for this product in

preruminating calves. Do not use in calves to be processed for veal.

[59 FR 19639, Apr. 25, 1994; 60 FR 26360, May 17, 1995, as amended at 62 FR 61625, Nov. 19, 1997; 64 FR 46840, Aug. 27, 1999; 67 FR 6867, Feb. 14, 2002; 70 FR 6764, Feb. 9, 2005]

§ 522.2690 Zinc gluconate.

(a) *Specifications.* Each milliliter of solution contains 13.1 milligrams zinc as zinc gluconate neutralized to pH 7.0 with L-arginine.

(b) *Sponsor.* See No. 067647 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs*—(1) *Amount.* The volume injected into each testicle is based on testicular width as determined by measuring each testicle at its widest point using a metric scale (millimeter) caliper.

(2) *Indications for use.* Intratesticular injection for chemical sterilization of 3- to 10-month-old male dogs.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[68 FR 26995, May 19, 2003]

PART 524—OPHTHALMIC AND TOPICAL DOSAGE FORM NEW ANIMAL DRUGS

Sec.

- 524.86 Amitraz liquid.
- 524.154 Bacitracin or bacitracin zinc-neomycin sulfate-polymyxin B sulfate ophthalmic ointment.
- 524.155 Bacitracin zinc-polymyxin B sulfate-neomycin sulfate-hydrocortisone or hydrocortisone acetate ophthalmic ointment.
- 524.390 Chloramphenicol ophthalmic and topical dosage forms.
- 524.390a Chloramphenicol ophthalmic ointment.
- 524.390b Chloramphenicol ophthalmic solution.
- 524.390d Chloramphenicol-prednisolone ophthalmic ointment.
- 524.402 Chlorhexidine.
- 524.450 Clotrimazole cream.
- 524.463 Copper naphthenate.
- 524.520 Cuprimyxin cream.
- 524.575 Cyclosporine ophthalmic ointment.
- 524.590 Diclofenac.
- 524.660 Dimethyl sulfoxide ophthalmic and topical dosage forms.
- 524.660a Dimethyl sulfoxide solution.
- 524.660b Dimethyl sulfoxide gel.
- 524.770 Doramectin.
- 524.775 Emodepside and praziquantel.