

§ 524.450

(c) *Conditions of use in dogs, cats, and horses*—(1) *Indications for use.* For use as a topical antiseptic ointment for surface wounds.

(2) *Limitations.* Do not use in horses intended for human consumption.

[72 FR 265, Jan. 4, 2007]

§ 524.450 Clotrimazole cream.

(a) *Specifications.* Each gram of cream contains 10 milligrams of clotrimazole.

(b) *Sponsor.* See 000859 in § 510.600(c).

(c) *Conditions of use*—(1) *Amount.* Apply ¼-inch ribbon of cream per square inch of lesion once daily for 2 to 4 weeks.

(2) *Indications of use.* For the treatment of fungal infections of dogs and cats caused by *Microsporum canis* and *Trichophyton mentagrophytes*.

(3) *Limitations.* Wash hands thoroughly after use to avoid spread of infection. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 48128, July 18, 1980]

§ 524.463 Copper naphthenate.

(a) *Amount.* The drug is a 37.5 percent solution of copper naphthenate.

(b) *Sponsors.* See Nos. 000856, 017135, and 058829 in § 510.600(c) of this chapter.

(c) *Conditions of use in horses*—(1) *Amount.* Apply daily to affected hooves until fully healed.

(2) *Indications for use.* As an aid in treating horses and ponies for thrush caused by organisms susceptible to copper naphthenate.

(3) *Limitations.* Use on horses and ponies only. Avoid contact around eyes. Do not contaminate feed. Do not use in horses intended for human consumption.

[47 FR 4250, Jan. 29, 1982, as amended at 68 FR 55825, Sept. 29, 2003; 71 FR 38074, July 5, 2006]

§ 524.520 Cuprimyxin cream.

(a) *Specifications.* The drug contains 0.5 percent cuprimyxin (6-methoxy-1-phenazinol 5, 10-dioxide, cupric complex) in an aqueous cream base.

(b) *Sponsor.* See No. 063238 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) Cuprimyxin is a broad spectrum antibacterial and antifungal cream for the topical treat-

21 CFR Ch. I (4–1–11 Edition)

ment of superficial infections in horses, dogs, and cats caused by bacteria, dermatophytes (*Trichophyton* spp., *Microsporum* spp.) and yeast (*Candida albicans*) affecting skin, hair, and external mucosae.

(2) The cream is applied twice daily to affected areas by rubbing into lesions. Treatment should be continued for a few days after clinical recovery to avoid possible relapses.

(3) After application to cutaneous areas, a change in color from dark green to pink is due to the liberation of free myxin from its copper complex.

(4) If no response is seen within seven days, diagnosis and therapy should be reevaluated. If any adverse local reaction is observed after topical application, discontinue treatment.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 45 FR 56799, Aug. 26, 1980; 66 FR 46706, Sept. 7, 2001]

EFFECTIVE DATE NOTE: At 76 FR 17778, Mar. 31, 2011, § 524.520 was removed, effective April 11, 2011.

§ 524.575 Cyclosporine ophthalmic ointment.

(a) *Specifications.* Each gram of ointment contains 2 milligrams of cyclosporine.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* Apply a ¼-inch strip of ointment to the affected eye(s) every 12 hours.

(2) *Indications for use.* For management of chronic keratoconjunctivitis sicca (KCS) and chronic superficial keratitis (CSK) in dogs.

(3) *Limitations.* Place ointment directly on cornea or into the conjunctival sac. Safety of use in puppies, pregnant or breeding animals has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[60 FR 48651, Sept. 20, 1995, as amended at 62 FR 48940, Sept. 18, 1997]

§ 524.590 Diclofenac.

(a) *Specifications.* Each gram of cream contains 10 milligrams diclofenac sodium.