

**§ 606.122**

**21 CFR Ch. I (4–1–10 Edition)**

transfusion shall be labeled: “Contains (name of antibody).”

(5) Recovered plasma labels shall include:

(i) In lieu of an expiration date, the date of collection of the oldest material in the container.

(ii) The statement as applicable: “Caution: For Manufacturing Use Only”; or “Caution: For Use in Manufacturing Noninjectable Products Only.” If the recovered plasma has a reactive screening test for evidence of infection due to a communicable disease agent(s) under § 610.40 of this chapter, or is collected from a donor with a previous record of a reactive screening test for evidence of infection due to a communicable disease agent(s) under § 610.40 of this chapter, the recovered plasma must be labeled as required under § 610.40(h)(2)(ii)(E) of this chapter.

(iii) For recovered plasma not meeting the requirements for manufacture into licensable products, the statement: “Not for Use in Products Subject to License Under Section 351 of the Public Health Service Act.”

(f) Blood and blood components determined to be unsuitable for transfusion shall be prominently labeled: “NOT FOR TRANSFUSION”, and the label shall state the reason the unit is considered unsuitable. The provision does not apply to recovered plasma labeled according to paragraph (e)(5) of this section.

(g) [Reserved]

(h) The following additional information shall appear on the label for blood or blood components shipped in an emergency, prior to completion of required tests, in accordance with § 640.2(f) of this chapter:

(1) The statement: “FOR EMERGENCY USE ONLY BY \_\_\_\_\_.”

(2) Results of any tests prescribed under §§ 610.40, and 640.5 (a), (b), or (c) of this chapter completed before shipment.

(3) Indication of any tests prescribed under §§ 610.40, and 640.5 (a), (b), or (c) of this chapter and not completed before shipment.

(i) The following additional information shall appear on the label for Whole Blood or Red Blood Cells intended for autologous infusion:

(1) Information adequately identifying the patient, e.g., name, blood group, hospital, and identification number.

(2) Date of donation.

(3) The statement: “FOR AUTOLOGOUS USE ONLY.”

(4) In place of the blood group label, each container of blood intended for autologous use and obtained from a donor who fails to meet any of the donor suitability requirements under § 640.3 of this chapter or who is reactive in the hepatitis tests prescribed under § 610.40 of this chapter shall be prominently and permanently labeled: “FOR AUTOLOGOUS USE ONLY.”

(5) Units of blood originally intended for autologous use, except those labeled as prescribed under paragraph (i)(4) of this section, may be issued for homologous transfusion provided the container label complies with all applicable provisions of paragraphs (b) through (e) of this section. In such case, the special label required under paragraph (i) (1), (2), and (3) of this section shall be removed or otherwise obscured.

(j) A tie-tag attached to the container may be used for providing the information required by paragraph (e) (1)(iii), (2)(ii), and (4), (h), or (i)(1), (2), and (3) of this section.

[50 FR 35469, Aug. 30, 1985, as amended at 53 FR 116, Jan. 5, 1988; 55 FR 11014, Mar. 26, 1990; 57 FR 10814, Mar. 31, 1992; 59 FR 23636, May 6, 1994; 63 FR 16685, Apr. 6, 1998; 64 FR 45371, Aug. 19, 1999; 66 FR 31162, June 11, 2001; 67 FR 4907, Feb. 1, 2002; 69 FR 9171, Feb. 26, 2004; 70 FR 14984, Mar. 24, 2005]

**§ 606.122 Instruction circular.**

An instruction circular shall be available for distribution if the product is intended for transfusion. The instruction circular shall provide adequate directions for use, including the following information:

(a) Instructions to mix the product before use.

(b) Instructions to use a filter in the administration equipment.

(c) The statement “Do Not Add Medications” or an explanation concerning allowable additives.

(d) A description of the product, its source, and preparation, including the

name and proportion of the anticoagulant used in collecting the Whole Blood from each product is prepared.

(e) Statements that the product was prepared from blood that was negative when tested for antibody to Human Immunodeficiency Virus (HIV) and non-reactive for hepatitis B surface antigen by FDA required tests and nonreactive when tested for syphilis by a serologic test for syphilis (STS).

(f) The statements: "Warning. The risk of transmitting infectious agents is present. Careful donor selection and available laboratory tests do not eliminate the hazard."

(g) The names of cryoprotective agents and other additives that may still be present in the product.

(h) The names and results of all tests performed when necessary for safe and effective use.

(i) The use of the product, indications, contraindications, side effects and hazards, dosage and administration recommendations.

(j) [Reserved]

(k) For Red Blood Cells, the instruction circular shall contain:

(1) Instructions to administer a suitable plasma volume expander if Red Blood Cells are substituted when Whole Blood is the indicated product.

(2) A warning not to add Lactated Ringer's Injection U.S.P. solution to Red Blood Cell products.

(1) For Platelets, the instruction circular shall contain:

(1) The approximate volume of plasma from which a sample unit of Platelets is prepared.

(2) Instructions to begin administration as soon as possible, but not more than 4 hours after entering the container.

(m) For Plasma, the instruction circular shall contain:

(1) A warning against further processing of the frozen product if there is evidence of breakage or thawing.

(2) Instructions to thaw the frozen product at a temperature between 30 and 37 °C.

(3) When applicable, instructions to begin administration of the product within 6 hours after thawing.

(4) Instructions to administer to ABO-group-compatible recipients.

(5) A statement that this product has the same hepatitis risk as Whole Blood; other plasma volume expanders without this risk are available for treating hypovolemia.

(n) For Cryoprecipitated AHF, the instruction circular shall contain:

(1) A statement that the average potency is 80 or more International Units of antihemophilic factor.

(2) The statement: "Usually contains at least 150 milligrams of fibrinogen"; or, alternatively, the average fibrinogen level determined by assay of representative units.

(3) A warning against further processing of the product if there is evidence of breakage or thawing.

(4) Instructions to thaw the product for no more than 15 minutes at a temperature of between 30 and 37 °C.

(5) Instructions to store at room temperature after thawing and to begin administration as soon as possible but no more than 4 hours after entering the container or after pooling and within 6 hours after thawing.

(6) A statement that 0.9 percent Sodium Chloride Injection U.S.P. is the preferred diluent.

(7) Adequate instructions for pooling to ensure complete removal of all concentrated material from each container.

(8) The statement: "Good patient management requires monitoring treatment responses to Cryoprecipitated AHF transfusions with periodic plasma factor VIII or fibrinogen assays in hemophilia A and hypofibrinogenemic recipients, respectively."

[50 FR 35470, Aug. 30, 1985, as amended at 53 FR 116, Jan. 5, 1988; 64 FR 45371, Aug. 19, 1999]

### Subpart H—Laboratory Controls

#### § 606.140 Laboratory controls.

Laboratory control procedures shall include:

(a) The establishment of scientifically sound and appropriate specifications, standards and test procedures to assure that blood and blood components are safe, pure, potent and effective.