

## § 640.66

continuance in the plasmapheresis program will not interfere with or jeopardize the treatment of the syphilitic donor.

(3) A donor identification system shall be established that positively identifies each donor and relates such donor directly to his blood and its components as well as to his accumulated records and laboratory data. Such system shall include either a photograph of each donor which shall be used on each visit to confirm the donor's identity, or some other method that provides equal or greater assurance of positively identifying the donor.

(4) The amount of whole blood, not including anticoagulant, removed from a donor during a manual plasmapheresis procedure or in any 2-day period shall not exceed 1,000 milliliters unless the donor's weight is 175 pounds or greater, in which case the amount of whole blood, not including anticoagulant, removed from the donor during a manual plasmapheresis procedure or in any 2-day period shall not exceed 1,200 milliliters.

(5) The amount of whole blood, not including anticoagulant, removed from a donor during a manual plasmapheresis procedure within a 7-day period shall not exceed 2,000 milliliters unless the donor's weight is 175 pounds or greater, in which case the amount of whole blood, not including anticoagulant, removed from a donor during a manual plasmapheresis procedure within a 7-day period shall not exceed 2,400 milliliters.

(6) No more than 500 milliliters of whole blood shall be removed from a donor at one time, unless the donor's weight is 175 pounds or greater, in which case no more than 600 milliliters of whole blood shall be removed from the donor at one time.

(7) The plasma shall be separated from the red blood cells immediately after blood collection. The maximum feasible volume of red blood cells shall be returned to the donor before another unit is collected.

(8) The volume of plasma collected during an automated plasmapheresis collection procedure shall be consistent with the volumes specifically approved by the Director, Center for Biologics Evaluation and Research, and

## 21 CFR Ch. I (4-1-10 Edition)

collection shall not occur less than 2 days apart or more frequently than twice in a 7-day period.

[38 FR 32089, Nov. 20, 1973, as amended at 41 FR 10769, Mar. 12, 1976; 64 FR 45373, Aug. 19, 1999; 64 FR 56453, Oct. 20, 1999]

### § 640.66 Immunization of donors.

If specific immunization of a donor is to be performed, the selection and scheduling of the injection of the antigen, and the evaluation of each donor's clinical response, shall be by a qualified licensed physician or physicians. The administration of the antigen may be performed by a licensed physician or a trained person under his supervision. Any material used for immunization shall be either a product licensed under section 351 of the Public Health Service Act for such purpose or one specifically approved by the Director, Center for Biologics Evaluation and Research, Food and Drug Administration. Immunization procedures shall be on file at each plasmapheresis center where immunizations are performed.

[38 FR 32089, Nov. 20, 1973, as amended at 49 FR 23834, June 8, 1984; 55 FR 11013, Mar. 26, 1990]

### § 640.67 Laboratory tests.

Each unit of Source Plasma shall be tested for evidence of infection due to communicable disease agents as required under § 610.40 of this chapter.

[66 FR 31165, June 11, 2001]

### § 640.68 Processing.

(a) *Sterile system.* All administration and transfer sets inserted into blood containers used for processing Source Plasma intended for manufacturing into injectable or noninjectable products and all interior surfaces of plasma containers used for processing Source Plasma intended for manufacturing into injectable products shall be sterile, pyrogen-free, nontoxic, and compatible with the contents under normal conditions of use. Only Sodium Chloride Injection USP shall be used as a red blood cell diluent. If the method of separation of the plasma intended for injectable products involves a system in which an airway must be inserted into the plasma container, the airway shall be sterile and constructed so as to