

**Environmental Protection Agency**

**§ 180.603**

(d) *Indirect or inadvertent residues.*  
[Reserved]

[69 FR 40781, July 7, 2004, as amended at 71 FR 52487, Sept. 6, 2006; 74 FR 9377, Mar. 4, 2009]

**§ 180.601 Cyazofamid; tolerances for residues.**

(a) *General.* Tolerances are established for residues of the fungicide cyazofamid, including its metabolites and degradates, in or on the commodities in the following table. Compliance with the tolerance levels specified in the following table is to be determined by measuring only the sum of 4-chloro-2-cyano-*N,N*-dimethyl-5-(4-methylphenyl)-1*H*-imidazole-1-sulfonamide and its metabolite, 4-chloro-5-(4-methylphenyl)-1*H*-imidazole-2-carbonitrile, calculated as the stoichiometric equivalent of cyazofamid, in or on the following commodities:

| Commodity                                  | Parts per million |
|--|-------------------|
| Brassica, head and stem, subgroup 5A ..... | 1.2               |
| Brassica, leafy greens, subgroup 5B .....  | 12.0              |
| Carrot, roots .....                        | 0.09              |
| Hop dried cones .....                      | 10.0              |
| Okra .....                                 | 0.40              |
| Potato .....                               | 0.02              |
| Spinach .....                              | 9.0               |
| Turnip, greens .....                       | 12.0              |
| Vegetable, cucurbit, group 9 .....         | 0.10              |
| Vegetable, fruiting, group 8 .....         | 0.40              |

(b) *Section 18 emergency exemptions.*  
[Reserved]

(c) *Tolerances with regional registrations.* Tolerances with regional registrations are established for residues of the fungicide cyazofamid, including its metabolites and degradates, in or on the commodities in the following table. Compliance with the tolerance levels specified in the following table is to be determined by measuring only the sum of 4-chloro-2-cyano-*N,N*-dimethyl-5-(4-methylphenyl)-1*H*-imidazole-1-sulfonamide and its metabolite, 4-chloro-5-(4-methylphenyl)-1*H*-imidazole-2-carbonitrile, calculated as the stoichiometric equivalent of cyazofamid, in or on the following commodities:

| Commodity   | Parts per million |
|-------------|-------------------|
| Grape ..... | 1.5               |

(d) *Indirect or inadvertent residues.*  
[Reserved]

[69 FR 58299, Sept. 30, 2004, as amended at 73 FR 21839, Apr. 23, 2008; 74 FR 32453, July 8, 2009; 75 FR 40751, July 14, 2010]

**§ 180.602 Spiroxamine; tolerances for residues.**

(a) *General.* Tolerances are established for residues of the fungicide spiroxamine, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified in the following table is to be determined by measuring only spiroxamine, [(8-(1,1-dimethylethyl)-*N*-ethyl-*N*-propyl-1,4-dioxaspiro[4,5]decane-2-methanamine) in or on the commodities.

| Commodity  | Parts per million |
|--|-------------------|
| Artichoke, globe, import <sup>1</sup> .....          | 0.7               |
| Asparagus <sup>1</sup> .....                         | 0.05              |
| Banana (import) .....                                | 3.0               |
| Grape (import) .....                                 | 1.0               |
| Hop, dried cones .....                               | 50                |
| Vegetable, fruiting, crop group 8 <sup>1</sup> ..... | 1.2               |

<sup>1</sup> No U.S. registration as of December 1, 2010.

(b) *Section 18 emergency exemptions.*  
[Reserved]

(c) *Tolerances with regional registrations.* [Reserved]

(d) *Indirect or inadvertent residues.*  
[Reserved]

[69 FR 42570, July 16, 2004, as amended at 75 FR 74640, Dec. 1, 2010]

**§ 180.603 Dinotefuran; tolerances for residues.**

(a) *General.* (1) Tolerances are established for residues of dinotefuran, (*RS*)-1-methyl-2-nitro-3-((tetrahydro-3-furanyl)methyl)guanidine, including its metabolites and degradates, in or on the commodities listed in the following table. Compliance with the tolerance levels specified below is to be determined by measuring only the sum of dinotefuran and its metabolites DN, 1-methyl-3-(tetrahydro-3-furylmethyl)guanidine, and UF, 1-methyl-3-(tetrahydro-3-furylmethyl)urea, calculated as the stoichiometric equivalent of dinotefuran, in or on the commodities listed in the table below: