

§ 29.1071

Middle Belt Flue-cured, normally characterized by a thinner body and lighter color shade and produced principally in a section lying between the Piedmont and coastal plains regions of Virginia and North Carolina, may be classified as Type 11b.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

§ 29.1071 Type 12.

That type of flue-cured tobacco commonly known as Eastern Flue-cured or Eastern Carolina Flue-cured, produced principally in the coastal plains section of North Carolina, north of the South River.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

§ 29.1072 Type 13.

That type of flue-cured tobacco commonly known as Southeastern Flue-cured or South Carolina Flue-cured, produced principally in the coastal plains section of South Carolina and the southeastern counties of North Carolina, south of the South River.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

§ 29.1073 Type 14.

That type of flue-cured tobacco commonly known as Southern Flue-cured, produced principally in the southern section of Georgia, in northern Florida, and to some extent in Alabama.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

§ 29.1074 Type 92.

That type of flue-cured tobacco commonly known as Foreign-grown Flue-cured, produced in countries other than the United States.

[49 FR 16756, Apr. 20, 1984. Redesignated at 51 FR 25027, July 10, 1986]

7 CFR Ch. I (1-1-11 Edition)

§ 29.1075 Undried.

The condition of unfermented tobacco which has not been air-dried or steam-dried.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

§ 29.1076 Uniformity.

An element of quality which describes the consistency of a lot of tobacco as it is prepared for market. Uniformity is expressed as a percentage in grade specifications. (See Rule 13.)

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

§ 29.1077 Unsound (U).

Damaged. (See Rule 21.)

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

§ 29.1078 Unstemmed.

A form of tobacco, including whole leaf and leaf scrap, from which the stems or midribs have not been removed.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

§ 29.1079 Variegated (K).

Any tobacco that does not blend with the normal colors of the types; any leaf of which 20 percent or more of its surface is grayish, mottled, bleached, doty-faced, scalded, or sunbaked. (See Rule 15.)

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

§ 29.1080 Variegated dark red (KD).

A dark brownish-red discoloration which usually results from excessive sunbaking during the growing process or from storing cured tobacco over extended periods of time. Any leaf of which 20 percent or more of its surface

Agricultural Marketing Service, USDA

§ 29.1085

is dark brownish-red may be described as variegated dark red.

[48 FR 29671, June 28, 1983. Redesignated at 49 FR 16756, Apr. 20, 1984 and 51 FR 25027, July 10, 1986]

§ 29.1081 Variegated red or scorched (KR).

A red discoloration which usually results from excessive heat in the curing process. Any leaf of which 20 percent or more of its surface has been reddened in the curing process may be described as variegated red or scorched. (See Rule 16.)

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

§ 29.1082 Waste.

The portion or portions of the web of tobacco leaves which have been lost or rendered less serviceable for use in tobacco products, including:

- (a) Portions which have decomposed or largely decomposed by field diseases and field-firing, pole-burning, bulk-burning;
- (b) portions which are dead, lifeless, and do not have sufficient strength or stability to hold together

§ 29.1085 Width.

The relative breadth of a tobacco leaf expressed in relation to its length. Width, as an element of quality, does not apply to tobacco in strip form. (See Elements of Quality Chart.)

Elements		Degrees			
Maturity	Immature	Unripe	Mature	Ripe	Mellow.
Leaf structure	Tight	Close	Firm	Open.	
Body	Heavy	Fleshy	Medium	Thin.	
Oil	Lean	Oily	Rich.		
Color intensity	Pale	Weak	Moderate	Strong	Deep.
Width	Stringy	Narrow	Normal	Spready.	
Length		(¹)	(¹)	(¹).	
Uniformity		(²)	(²)	(²).	
Injury tolerance		(²)	(²)	(²).	
Waste tolerance		(²)	(²)	(²).	

¹ Expressed in inches.
² Expressed in percentage.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, and at 48 FR 29671, June 28, 1983. Redesignated and amended at 49 FR 16756, Apr. 20, 1984, and further redesignated at 51 FR 25027, July 10, 1986]

in the normal manufacturing process due to excessive injury of any kind.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

§ 29.1083 Wet (W).

Any sound tobacco containing excessive moisture to the extent that it is in unsafe or doubtful-keeping order. Wet applies to any tobacco which is not damaged but which is likely to damage if treated in the customary manner. (See Rule 22.)

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

§ 29.1084 Whitish-lemon (LL).

A whitish-yellow color which usually results during wet growing seasons when rain leaches or washes out the yellow color from the leaf. Any leaf of which 20 percent or more of its leaf surface has whitish-yellow color may be described as whitish-lemon.

[48 FR 29671, June 28, 1983. Redesignated at 49 FR 16756, Apr. 20, 1984 and 51 FR 25027, July 10, 1986]