

## § 520.1120b

of breeding age; animals should be re-treated in 3 to 4 weeks.

[40 FR 13838, Mar. 27, 1975, as amended at 45 FR 10333, Feb. 15, 1980; 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61624, Nov. 19, 1997]

### § 520.1120b Haloxon boluses.

(a) *Chemical name.* 3-Chloro-7-hydroxy-4-methylcoumarin bis (2-chloroethyl) phosphate.

(b) *Specifications.* Each bolus contains 10.1 grams of haloxon.

(c) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(d) *Related tolerances.* See § 556.310 of this chapter.

(e) *Conditions of use.* (1) Haloxon bolus is an anthelmintic used in cattle for the control of gastrointestinal roundworms of the genera *Haemonchus*, *Ostertagia*, *Trichostrongylus* and *Cooperia*.

(2) It is administered by giving one bolus per approximately 500 pounds body weight (35 to 50 milligrams per kilogram of body weight).

(3) For most effective results, re-treat animals in 3 to 4 weeks. If reinfection is likely to occur, additional re-treatments may be necessary.

(4) Do not use any drug, pesticide or other chemical having cholinesterase inhibiting activity either simultaneously or within a few days before or after treatment with haloxon.

(5) Do not treat animals within one week of slaughter.

(6) Do not treat dairy animals of breeding age or older.

[40 FR 13838, Mar. 27, 1975, as amended at 44 FR 61591, Oct. 29, 1979; 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

### § 520.1130 Hetacillin.

(a) *Specifications.*—(1) Each capsule or tablet contains hetacillin potassium equivalent to 50, 100, or 200 milligrams (mg) of ampicillin.

(2) Each milliliter of suspension contains hetacillin potassium equivalent to 50 mg of ampicillin.

(b) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs and cats.*—(1) *Amount.*—(i) *Dogs.* Administer 5 mg per pound (lb) of body weight orally, twice daily. In severe infections, ad-

## 21 CFR Ch. I (4–1–12 Edition)

minister 5 mg/lb three times daily, or up to 10 mg/lb twice daily. For stubborn urinary tract infections, administer up to 20 mg/lb twice daily.

(ii) *Cats.* Administer 50 mg twice daily.

(2) *Indications for use.* For the treatment of respiratory tract infections, urinary tract infections, gastrointestinal infections, skin infections, soft tissue infections, and postsurgical infections associated with strains of organisms susceptible to hetacillin potassium.

(3) *Limitations.* Federal law restricts this drug to use only by or on the order of a licensed veterinarian.

[75 FR 10166, Mar. 5, 2010]

### § 520.1157 Iodinated casein tablets.

(a) *Specifications.* Each 1-gram tablet contains 25 milligrams of iodinated casein.

(b) *Sponsor.* See No. 017762 in § 510.600(c) of this chapter.

(c) *Conditions of use.*—(1) *Amount.* ½ to 1 tablet per 10 pounds of body weight (equivalent to 0.5 to 2.5 milligrams of iodinated casein per pound of body weight).

(2) *Indications for use.* For dogs for apparent decreased thyroid activity where the signs are alopecia, scaliness of the skin surface, loss of hair, seborrhea, thickening of the skin, hyperpigmentation, and lethargy.

(3) *Limitations.* If no response is observed in 30 to 45 days, the drug should be withdrawn and the diagnosis reconsidered. Do not use in the presence of cardiac disease, ischemia, adrenal insufficiency, or nephrosis. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[49 FR 22469, May 30, 1984]

### § 520.1158 Iodochlorhydroxyquin boluses.

(a) *Specifications.* Each bolus contains 10 grams of iodochlorhydroxyquin.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *Conditions of use.*—(1) *Amount.* 1 bolus (10 grams) daily for a 1,000-pound horse.

(2) *Indications for use.* For treatment of equine diarrhea.