

dermatitis and for the relief of superficial inflammation caused by chemical and physical abrasions and burns.

(2) A small amount is applied to the affected area two or three times daily.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 62 FR 40932, July 31, 1997]

§ 524.981b Fluocinolone acetonide solution.

(a) *Specifications.* The drug contains 0.01 percent fluocinolone acetonide in propylene glycol with citric acid.

(b) *Sponsor.* See No. 099207 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is indicated for the relief of pruritus and inflammation associated with otitis externa and certain superficial acute and chronic dermatoses in the dog. It is also indicated for the relief of pruritus and inflammation associated with acute otitis externa and certain superficial acute and chronic dermatoses in the cat.

(2) A small amount of solution is applied to the affected area 2 or 3 times daily.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 62 FR 40932, July 31, 1997]

§ 524.981c Fluocinolone acetonide, neomycin sulfate cream.

(a) *Specifications.* The drug contains 0.025 percent fluocinolone acetonide and 0.5 percent neomycin sulfate (0.35 percent neomycin base).

(b) *Sponsor.* See No. 099207 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in the relief of pruritus and inflammation associated with superficial acute and chronic dermatoses in dogs. It is used in the treatment of such conditions as allergic and acute moist dermatoses and nonspecific dermatoses in dogs. It is used in the treatment of wound infections in dogs and cats.

(2) A small amount is applied to the infected area two or three times daily.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 62 FR 40932, July 31, 1997]

§ 524.981d Fluocinolone acetonide, dimethyl sulfoxide solution.

(a) *Specifications.* Each milliliter of solution contains 0.01 percent fluocinolone acetonide and 20 percent dimethyl sulfoxide with propylene glycol and citric acid.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs for the relief of impaction commonly present in apparently normal anal sacs, for the reversal of inflammatory changes associated with abnormal anal sacs, and to counteract the offensive odor of anal sac secretions.

(2) It is administered by instillation of 1 to 2 milliliters into each anal sac following expression of anal sac contents. It may be necessary to repeat treatment at 60-day intervals to maintain an odor-free state. The total dosage used should not exceed 2 milliliters per anal sac per treatment.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 61 FR 5507, Feb. 13, 1996]

§ 524.981e Fluocinolone acetonide, dimethyl sulfoxide otic solution.

(a) *Specifications.* Each milliliter of solution contains 0.01 percent of fluocinolone acetonide in 60 percent dimethyl sulfoxide with propylene glycol and citric acid.

(b) *Sponsor.* See No. 000033 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs for the relief of pruritis and inflammation associated with acute and chronic otitis.

(2) It is administered at 4 to 6 drops (0.2 milliliter) twice daily into the ear canal for a maximum period of 14 days. The total dosage used should not exceed 17 milliliters. The ear canal should be cleansed by some appropriate method prior to instillation of the solution and the ear should be massaged gently following instillation.