Food and Drug Administration, HHS

	Limitations
Chlorine, as sodium hypochlorite, not to exceed 0.055 pound of chlorine per pound of dry starch; 0.45 percent of active oxygen obtained from hydrogen peroxide; and propylene oxide, not to exceed 25 percent. Sodium hydroxide, not to exceed 1 percent.	Residual propylene chlorohydrin not more than 5 parts per millior in food starch-modi- fied.

- (h) Food starch may be modified by a combination of the treatments prescribed by paragraphs (a), (b), and/or (i) of this section and any one of the treatments prescribed by paragraph (c), (d), (e), (f), or (g) of this section, subject to any limitations prescribed by the paragraphs named.
- (i) Food starch may be modified by treatment with the following enzymes:

Enzyme	Limitations
Alpha-amylase (E.C. 3.2.1.1)	The enzyme must be generally recognized as safe or approved as a food additive for this purpose. The resulting nonsweet nutritive saccharide polymer has a dextrose equivalent of less than 20.
Beta-amylase (E.C. 3.2.1.2). Glucoamylase (E.C. 3.2.1.3). Isoamylase (E.C. 3.2.1.68). Pullulanase (E.C. 3.2.1.41).	

 $[42\ FR\ 14491,\ Mar.\ 15,\ 1977,\ as\ amended\ at\ 43\ FR\ 11697,\ Mar.\ 21,\ 1978;\ 46\ FR\ 32015,\ June\ 19,\ 1981;\ 57\ FR\ 54700,\ Nov.\ 20,\ 1992;\ 58\ FR\ 21100,\ Apr.\ 19,\ 1993;\ 66\ FR\ 17509,\ Apr.\ 2,\ 2001]$

§ 172.894 Modified cottonseed products intended for human consumption.

The food additive modified cottonseed products may be used for human consumption in accordance with the following prescribed conditions:

- (a) The additive is derived from:
- (1) Decorticated, partially defatted, cooked, ground cottonseed kernels; or
- (2) Decorticated, ground cottonseed kernels, in a process that utilizes *n*-hexane as an extracting solvent in such a way that no more than 60 parts per million of *n*-hexane residues and less than 1 percent fat by weight remain in the finished product; or
- (3) Glandless cottonseed kernels roasted to attain a temperature of not less than 250 °F in the kernel for not less than 5 minutes for use as a snack

food, or in baked goods, or in soft candy; or

- (4) Raw glandless cottonseed kernels may be used in hard candy where the kernel temperature during cooking will exceed 250 °F for not less than 5 minutes.
- (b) The additive is prepared to meet the following specifications:
- (1) Free gossypol content not to exceed 450 parts per million.
- (2) It contains no added arsenic compound and therefore may not exceed a maximum natural background level of 0.2 part per million total arsenic, calculated as As.
- (c) To assure safe use of the additive, the label of the food additive container shall bear, in addition to other information required by the act, the name of the additive as follows:
- (1) The additive identified in paragraph (a)(1) of this section as "partially defatted, cooked cottonseed flour".
- (2) The additive identified in paragraph (a)(2) of this section as "defatted cottonseed flour".
- (3) The additive identified in paragraph (a)(3) of this section as "roasted glandless cottonseed kernels".
- (4) The additive identified in paragraph (a)(4) of this section as "raw glandless cottonseed kernels for use in cooked hard candy".
- (d) The Food and Drug Administration and the Environmental Protection Agency have determined that glandless cottonseed kernels permitted for use by this section are a distinct commodity from glanded cottonseed.

§172.896 Dried yeasts.

Dried yeast (Saccharomyces cerevisiae and Saccharomyces fragilis) and dried torula yeast (Candida utilis) may be safely used in food provided the total folic acid content of the yeast does not exceed 0.04 milligram per gram of yeast (approximately 0.008 milligram of pteroyglutamic acid per gram of yeast).

§172.898 Bakers yeast glycan.

Bakers yeast glycan may be safely used in food in accordance with the following conditions:

(a) Bakers yeast glycan is the comminuted, washed, pasteurized, and