

Feed component	Limitations (percent)
BHT (butylated hydroxytoluene) .....	2
Methionine hydroxy analog and its calcium salts .....	1
Piperazine, piperazine salts .....	0.8
Sodium propionate .....	1
Urea .....	1
Vitamins .....	3

(c) It is used in feed as an anticaking agent in an amount not to exceed that reasonably required to accomplish its intended effect and in no case in an amount to exceed 2 percent by weight of the finished feed.

**§573.960 Sorbitan monostearate.**

The food additive sorbitan monostearate may be safely used alone or in combination with polysorbate 60 as an emulsifier in mineral premixes and dietary supplements for animal feeds.

**§573.980 Taurine.**

The food additive taurine (2-aminoethanesulfonic acid) may be safely used in feed in accordance with the following prescribed conditions:

- (a) It is used as a nutritional supplement in the feed of growing chickens.
- (b) It is added to complete feeds so that the total taurine content does not exceed 0.054 percent of the feed.
- (c) To assure safe use of the additive, the label and labeling shall bear in addition to the other information required by the Act:
  - (1) The name of the additive.
  - (2) The quantity of the additive contained therein.
  - (3) Adequate directions for use.

**§573.1000 Verxite.**

The food additive verxite may be safely used in animal feed in accordance with the following prescribed conditions:

- (a) The additive is a magnesium-aluminum-iron silicate conforming to one of the following:
  - (1)(i) Verxite granules: The additive contains a minimum of 98 percent of hydrobiotite; it is thermally expanded and has a bulk density of from 5 to 9 pounds per cubic foot.
  - (ii) It is used or intended for use:
    - (a) In poultry feed at a level not to exceed 5 percent of the weight of the finished feed as a nonnutritive bulking

agent for restricting calorie intake in pullet replacement feeds.

(b) As an anticaking or blending agent, pelleting aid, or nonnutritive carrier for the incorporation of nutrients in poultry, swine, dog, or ruminant feeds, in an amount not to exceed that necessary to accomplish its intended effect and in no case to exceed 1.5 percent of the dog feed or 5 percent of the final feed for other animals.

(2)(i) Verxite flakes: The additive contains a minimum of 98 percent of hydrobiotite; it has a bulk density of from 20 to 30 pounds per cubic foot.

(ii) It is used or intended for use as an anticaking or blending agent in ruminant feeds in an amount not to exceed that necessary to accomplish its intended effect and in no case to exceed 1 percent by weight of the final feed for ruminants.

(3)(i) Verxite grits: The additive contains a minimum of 80 percent of hydrobiotite; it has a bulk density of from 40 to 50 pounds per cubic foot.

(ii) It is used or intended for use as a partial roughage replacement in ruminant feeds in an amount not to exceed that necessary to accomplish its intended effect and in no case to exceed 1 percent by weight of the final feed.

(b) To assure safe use of the additive, the label of any feed additive supplement, feed additive concentrate, feed additive premix, or complete feed prepared therefrom shall bear, in addition to the other information required by the Act, the name of the additive (verxite granules, verxite flakes, or verxite grits), adequate directions for use, and, when the additive is present in excess of 1 percent, a statement of the quantity of the additive contained therein and the term "nonnutritive" in juxtaposition therewith.

**§573.1010 Xanthan gum.**

The food additive xanthan gum may be safely used in animal feed as follows:

- (a) The food additive is xanthan gum as defined in §172.695 of this chapter and meets all of the specifications thereof.
  - (b) It is used or intended for use as a stabilizer, emulsifier, thickener, suspending agent, or bodying agent in animal feed as follows: