

## § 29.3520

### § 29.3520 Grade.

A subdivision of a type according to group, quality, and color.

### § 29.3521 Grademark.

A grademark normally consists of three symbols which indicate group, quality, and color. A letter is used to indicate group, a number to indicate quality, and a letter or letters to indicate color. For example, B3D means Heavy Leaf, third quality, and dark-brown color.

### § 29.3522 Green (G).

A term applied to green-colored, immature, or crude tobacco. Any leaf which has a green color affecting 20 percent or more of its leaf surface may be described as green. (See Rule 19, § 29.3620.)

### § 29.3523 Group.

A division of a type covering closely related grades based on certain characteristics which are related to stalk position, body, or the general quality of the tobacco. Groups in Dark Air-cured types are: Wrappers (A), Heavy Leaf (B), Thin Leaf (C), Lugs (X), Non-descript (N), and Scrap (S).

[51 FR 40408, Nov. 7, 1986]

### § 29.3524 Injury.

Hurt or impairment from any cause except the fungus or bacterial diseases which attack tobacco in its cured state. (See definition of Damage, § 29.3514.) Injury to tobacco may be caused by field diseases, insects, or weather conditions; insecticides, fungicides, or cell growth inhibitors; nutritional deficiencies or cesses; or improper fertilizing, harvesting, curing, or handling. Injured tobacco includes dead, burned, hail-cut torn, broken, frostbitten, sunburned, sunscalded, scorched, fire-killed, bulk-burnt, steam-burnt, house-burnt bleached, bruised, discolored, or deformed leaves; or tobacco affected by wildfire rust, frog-eye, mosaic, root rot, wilt, black shank, or other diseases. (See Rule 15, § 29.3616.)

### § 29.3525 Leaf.

Whole, unstemmed leaf. Leaf, when applied to tobacco in strip form, shall

## 7 CFR Ch. I (1-1-14 Edition)

describe the divided unit of a whole leaf.

[49 FR 16759, Apr. 20, 1984]

### § 29.3526 Leaf scrap.

A byproduct of unstemmed tobacco. Leaf scrap results from handling unstemmed tobacco and consists of loose and tangled whole or broken leaves.

[30 FR 9207, July 23, 1965. Redesignated at 49 FR 16759, Apr. 20, 1984]

### § 29.3527 Leaf structure.

The cell development of a leaf as indicated by its porosity. (See Elements of Quality, § 29.3586.)

[30 FR 9207, July 23, 1965. Redesignated at 49 FR 16759, Apr. 20, 1984]

### § 29.3528 Leaf surface.

The roughness or smoothness of the web or lamina of a tobacco leaf. Leaf surface is affected to some extent by the size and shrinkage of the veins or fibers (See Elements of Quality, § 29.3586.)

[30 FR 9207, July 23, 1965. Redesignated at 49 FR 16759, Apr. 20, 1984]

### § 29.3529 Length.

The linear measurement of cured tobacco leaves from the butt of the midrib to the extreme tip. (See Standard Tobacco Sizes, § 29.3591.)

[51 FR 40408, Nov. 7, 1986]

### § 29.3530 Lot.

A pile, basket, bulk, or more than one bale, case, hogshead, tierce, package, or other definite package unit.

[30 FR 9207, July 23, 1965. Redesignated at 49 FR 16759, Apr. 20, 1984]

### § 29.3531 Maturity.

The degree of ripeness. (See Elements of Quality, § 29.3586, and Rule 16, § 29.3617.)

[30 FR 9207, July 23, 1965. Redesignated at 49 FR 16759, Apr. 20, 1984]