not present at the time of the prior decision or apart of the evidentiary record in support of the higher-level review during the informal conference in accordance with paragraph (d) of this section. Any expenses incurred by the claimant in connection with the informal conference are the responsibility of the claimant.

- (i) De novo review. The higher-level adjudicator will consider only those decisions and claims for which the claimant has requested higher-level review, and will conduct a de novo review giving no deference to the prior decision, except as provided in §3.104(c).
- (j) Difference of opinion. The higherlevel adjudicator may grant a benefit sought in the claim under review based on a difference of opinion (see §3.105(b)). However, any finding favorable to the claimant is binding except as provided in §3.104(c) of this part. In addition, the higher-level adjudicator will not revise the outcome in a manner that is less advantageous to the claimant based solely on a difference of opinion. The higher-level adjudicator may reverse or revise (even if disadvantageous to the claimant) prior decisions by VA (including the decision being reviewed or any prior decision) on the grounds of clear and unmistakable error under $\S 3.105(a)(1)$ or (a)(2), as applicable, depending on whether the prior decision is finally adjudicated.
- (k) Notice requirements. Notice of a decision made under this section will include all of the elements described in §3.103(f), a general statement indicating whether evidence submitted while the record was closed was not considered, and notice of the options available to have such evidence considered.

(Authority: 38 U.S.C. 5109A and 7105(d))

[84 FR 173, Jan. 18, 2019]

PART 4—SCHEDULE FOR RATING DISABILITIES

Subpart A—General Policy in Rating

Sec.

- Essentials of evaluative rating 4.1
- 4.2 Interpretation of examination reports.
- 4.3 Resolution of reasonable doubt.
- Evaluation of evidence.
- 4.7 Higher of two evaluations.

- 4.9 Congenital or developmental defects.
- 4.10 Functional impairment.
- Effect of change of diagnosis. 4.13
- Avoidance of pyramiding. 4.14
- Total disability ratings.
- Total disability ratings for compensation based on unemployability of the individual.
- 4.17 Total disability ratings for pension based on unemployability and age of the individual.
- 4.17a Misconduct etiology.
- 4.18 Unemployability.
- Age in service-connected claims. 4.19
- 4.20 Analogous ratings.
- Application of rating schedule. 4.21
- Rating of disabilities aggravated by active service.
- 4.23 Attitude of rating officers.
- 4.24 Correspondence.
- Combined ratings table. 4.25
- 4.26 Bilateral factor.
- Use of diagnostic code numbers.
- Prestabilization rating from date of discharge from service.
- 4.29 Ratings for service-connected disabilities requiring hospital treatment or observation.
- 4.30 Convalescent ratings.
- 4.31 Zero percent evaluations.

Subpart B—Disability Ratings

THE MUSCULOSKELETAL SYSTEM

- Functional loss 4 40
- 4.41 History of injury.
- 4.42 Complete medical examination of injury cases.
- 4.43 Osteomyelitis.
- The bones. 4.44
- 4.45 The joints.
- Accurate measurement.
- 4.47-4.54 [Reserved]
- 4.55 Principles of combined ratings for muscle injuries.
- 4.56 Evaluation of muscle disabilities.
- Static foot deformities.
- Arthritis due to strain.
- Painful motion. 4.60 [Reserved]
- 4.61 Examination.
- Circulatory disturbances.
- Loss of use of hand or foot.
- Loss of use of both buttocks. 4.64
- [Reserved] 4.65
- Sacroiliac joint. 4.66
- 4.67 Pelvic bones.
- 4.68 Amputation rule.
- Dominant hand.
- Inadequate examinations. 4.70
- Measurement of ankylosis and joint motion.
- 4.71a Schedule of ratings—musculoskeletal system.
- 4.72 [Reserved]
- 4.73 Schedule of ratings—muscle injuries.

THE ORGANS OF SPECIAL SENSE

- 4.75 General considerations for evaluating visual impairment.
- 4.76 Visual acuity.
- 4.76a Computation of average concentric contraction of visual fields.
- 4.77 Visual fields.
- 4.78 Muscle function.
- 4.79 Schedule of ratings—eye.
- 4.80-4.84 [Reserved]

IMPAIRMENT OF AUDITORY ACUITY

- 4.85 Evaluation of hearing impairment.
- 4.86 Exceptional patterns of hearing impairment.
- $4.87 \quad \text{Schedule of ratings---ear.}$
- 4.87a Schedule of ratings—other sense organs.

INFECTIOUS DISEASES, IMMUNE DISORDERS AND NUTRITIONAL DEFICIENCIES

- 4.88 [Reserved]
- 4.88a Chronic fatigue syndrome.
- 4.88b Schedule of ratings—infectious diseases, immune disorders and nutritional deficiencies.
- 4.88c Ratings for inactive nonpulmonary tuberculosis initially entitled after August 19, 1968.
- 4.89 Ratings for inactive nonpulmonary tuberculosis in effect on August 19, 1968.

THE RESPIRATORY SYSTEM

- 4.96 Special provisions regarding evaluation of respiratory conditions.
- 4.97 Schedule of ratings—respiratory system.

THE CARDIOVASCULAR SYSTEM

- 4.100 Application of the evaluation criteria for diagnostic codes 7000-7007, 7011, and 7015-7020.
- 4.101-4.103 [Reserved]
- 4.104 Schedule of ratings—cardiovascular system.

THE DIGESTIVE SYSTEM

- 4.110 Ulcers.
- 4.111 Postgastrectomy syndromes.
- 4.112 Weight loss.
- 4.113 Coexisting abdominal conditions.
- 4.114 Schedule of ratings—digestive system.

THE GENITOURINARY SYSTEM

- 4.115 Nephritis.
- 4.115a Ratings of the genitourinary system—dysfunctions.
- 4.115b Ratings of the genitourinary system—diagnoses.

GYNECOLOGICAL CONDITIONS AND DISORDERS OF THE BREAST

4.116 Schedule of ratings—gynecological conditions and disorders of the breast.

THE HEMATOLOGIC AND LYMPHATIC SYSTEMS

4.117 Schedule of ratings—hemic and lymphatic systems.

THE SKIN

4.118 Schedule of ratings—skin.

THE ENDOCRINE SYSTEM

4.119 Schedule of ratings—endocrine system.

NEUROLOGICAL CONDITIONS AND CONVULSIVE DISORDERS

- 4.120 Evaluations by comparison.
- 4.121 Identification of epilepsy.
- 4.122 Psychomotor epilepsy.
- 4.123 Neuritis, cranial or peripheral.
- 4.124 Neuralgia, cranial or peripheral.
- 4.124a Schedule of ratings—neurological conditions and convulsive disorders.

MENTAL DISORDERS

- 4.125 Diagnosis of mental disorders.
- 4.126 Evaluation of disability from mental disorders.
- 4.127 Intellectual disability (intellectual developmental disorder) and personality disorders.
- 4.128 Convalescence ratings following extended hospitalization.
- 4.129 Mental disorders due to traumatic stress.
- 4.130 Schedule of ratings—Mental disorders.

DENTAL AND ORAL CONDITIONS

- 4.149 [Reserved]
- 4.150 Schedule of ratings—dental and oral conditions.
- APPENDIX A TO PART 4—TABLE OF AMEND-MENTS AND EFFECTIVE DATES SINCE 1946
- MENTS AND EFFECTIVE DATES SINCE 1946
 APPENDIX B TO PART 4—NUMERICAL INDEX OF
 DISARILITIES
- APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES

AUTHORITY: 38 U.S.C. 1155, unless otherwise noted.

SOURCE: 29 FR 6718, May 22, 1964, unless otherwise noted.

Subpart A—General Policy in Rating

§ 4.1 Essentials of evaluative rating.

This rating schedule is primarily a guide in the evaluation of disability resulting from all types of diseases and injuries encountered as a result of or incident to military service. The percentage ratings represent as far as can practicably be determined the average impairment in earning capacity resulting from such diseases and injuries and

Department of Veterans Affairs

their residual conditions in civil occupations. Generally, the degrees of disability specified are considered adequate to compensate for considerable loss of working time from exacerbations or illnesses proportionate to the severity of the several grades of disability. For the application of this schedule, accurate and fully descriptive medical examinations are required, with emphasis upon the limitation of activity imposed by the disabling condition. Over a period of many years, a veteran's disability claim may require reratings in accordance with changes in laws, medical knowledge and his or her physical or mental condition. It is thus essential, both in the examination and in the evaluation of disability, that each disability be viewed in relation to its history.

[41 FR 11292, Mar. 18, 1976]

§ 4.2 Interpretation of examination reports.

Different examiners, at different times, will not describe the same disability in the same language. Features of the disability which must have persisted unchanged may be overlooked or a change for the better or worse may not be accurately appreciated or described. It is the responsibility of the rating specialist to interpret reports of examination in the light of the whole recorded history, reconciling the various reports into a consistent picture so that the current rating may accurately reflect the elements of disability present. Each disability must be considered from the point of view of the veteran working or seeking work. If a diagnosis is not supported by the findings on the examination report or if the report does not contain sufficient detail, it is incumbent upon the rating board to return the report as inadequate for evaluation purposes.

[41 FR 11292, Mar. 18, 1976]

§4.3 Resolution of reasonable doubt.

It is the defined and consistently applied policy of the Department of Veterans Affairs to administer the law under a broad interpretation, consistent, however, with the facts shown in every case. When after careful consideration of all procurable and assem-

bled data, a reasonable doubt arises regarding the degree of disability such doubt will be resolved in favor of the claimant. See §3.102 of this chapter.

[40 FR 42535, Sept. 15, 1975]

§ 4.6 Evaluation of evidence.

The element of the weight to be accorded the character of the veteran's service is but one factor entering into the considerations of the rating boards in arriving at determinations of the evaluation of disability. Every element in any way affecting the probative value to be assigned to the evidence in each individual claim must be thoroughly and conscientiously studied by each member of the rating board in the light of the established policies of the Department of Veterans Affairs to the end that decisions will be equitable and just as contemplated by the requirements of the law.

§4.7 Higher of two evaluations.

Where there is a question as to which of two evaluations shall be applied, the higher evaluation will be assigned if the disability picture more nearly approximates the criteria required for that rating. Otherwise, the lower rating will be assigned.

§4.9 Congenital or developmental defects.

Mere congenital or developmental defects, absent, displaced or supernumerary parts, refractive error of the eye, personality disorder and mental deficiency are not diseases or injuries in the meaning of applicable legislation for disability compensation purposes.

 $[41~{
m FR}~11292,~{
m Mar.}~18,~1976]$

§4.10 Functional impairment.

The basis of disability evaluations is the ability of the body as a whole, or of the psyche, or of a system or organ of the body to function under the ordinary conditions of daily life including employment. Whether the upper or lower extremities, the back or abdominal wall, the eyes or ears, or the cardiovascular, digestive, or other system, or psyche are affected, evaluations are based upon lack of usefulness, of these

parts or systems, especially in self-support. This imposes upon the medical examiner the responsibility of furnishing, in addition to the etiological, anatomical, pathological, laboratory and prognostic data required for ordinary medical classification, full description of the effects of disability upon the person's ordinary activity. In this connection, it will be remembered that a person may be too disabled to engage in employment although he or she is up and about and fairly comfortable at home or upon limited activity.

[41 FR 11292, Mar. 18, 1976]

§4.13 Effect of change of diagnosis.

The repercussion upon a current rating of service connection when change is made of a previously assigned diagnosis or etiology must be kept in mind. The aim should be the reconciliation and continuance of the diagnosis or etiology upon which service connection for the disability had been granted. The relevant principle enunciated in §4.125, entitled "Diagnosis of mental disorders," should have careful attention in this connection. When any change in evaluation is to be made, the rating agency should assure itself that there has been an actual change in the conditions, for better or worse, and not merely a difference in thoroughness of the examination or in use of descriptive terms. This will not, of course, preclude the correction of erroneous ratings, nor will it preclude assignment of a rating in conformity with §4.7.

 $[29 \ FR \ 6718, \ May \ 22, \ 1964, \ as \ amended \ at \ 61 \ FR \ 52700, \ Oct. \ 8, \ 1996]$

§4.14 Avoidance of pyramiding.

The evaluation of the same disability under various diagnoses is to be avoided. Disability from injuries to the muscles, nerves, and joints of an extremity may overlap to a great extent, so that special rules are included in the appropriate bodily system for their evaluation. Dyspnea, tachycardia, nervouses, fatigability, etc., may result from many causes; some may be service connected, others, not. Both the use of manifestations not resulting from service-connected disease or injury in establishing the service-connected eval-

uation, and the evaluation of the same manifestation under different diagnoses are to be avoided.

§ 4.15 Total disability ratings.

The ability to overcome the handicap of disability varies widely among individuals. The rating, however, is based primarily upon the average impairment in earning capacity, that is, upon the economic or industrial handicap which must be overcome and not from individual success in overcoming it. However, full consideration must be given to unusual physical or mental effects in individual cases, to peculiar effects of occupational activities, to defects in physical or mental endowment preventing the usual amount of success in overcoming the handicap of disability and to the effect of combinations of disability. Total disability will be considered to exist when there is present any impairment of mind or body which is sufficient to render it impossible for the average person to follow a substantially gainful occupation; Provided, That permanent total disability shall be taken to exist when the impairment is reasonably certain to continue throughout the life of the disabled person. The following will be considered to be permanent total disability: the permanent loss of the use of both hands, or of both feet, or of one hand and one foot, or of the sight of both eyes, or becoming permanently helpless or permanently bedridden. Other total disability ratings are scheduled in the various bodily systems of this schedule.

§4.16 Total disability ratings for compensation based on unemployability of the individual.

(a) Total disability ratings for compensation may be assigned, where the schedular rating is less than total, when the disabled person is, in the judgment of the rating agency, unable to secure or follow a substantially gainful occupation as a result of service-connected disabilities: *Provided* That, if there is only one such disability, this disability shall be ratable at 60 percent or more, and that, if there are two or more disabilities, there shall be at least one disability ratable at 40

percent or more, and sufficient additional disability to bring the combined rating to 70 percent or more. For the above purpose of one 60 percent disability, or one 40 percent disability in combination, the following will be considered as one disability: (1) Disabilities of one or both upper extremities, or of one or both lower extremities, including the bilateral factor, if applicable, (2) disabilities resulting from common etiology or a single accident, (3) disabilities affecting a single body system, e.g. orthopedic, digestive, respiratory, cardiovascular-renal, neuropsychiatric, (4) multiple injuries incurred in action, or (5) multiple disabilities incurred as a prisoner of war. It is provided further that the existence or degree of nonservice-connected disabilities previous orunemployability status will be disregarded where the percentages referred to in this paragraph for the service-connected disability or disabilities are met and in the judgment of the rating agency such service-connected disabilities render the veteran unemployable. Marginal employment shall not be considered substantially gainful employment. For purposes of this section, marginal employment generally shall be deemed to exist when a veteran's earned annual income does not exceed the amount established by the U.S. Department of Commerce, Bureau of the Census, as the poverty threshold for one person. Marginal employment may also be held to exist, on a facts found basis (includes but is not limited to employment in a protected environment such as a family business or sheltered workshop), when earned annual income exceeds the poverty threshold. Consideration shall be given in all claims to the nature of the employment and the reason for termination.

(Authority: 38 U.S.C. 501)

(b) It is the established policy of the Department of Veterans Affairs that all veterans who are unable to secure and follow a substantially gainful occupation by reason of service-connected disabilities shall be rated totally disabled. Therefore, rating boards should submit to the Director, Compensation Service, for extra-schedular consideration all cases of veterans who are un-

employable by reason of service-connected disabilities, but who fail to meet the percentage standards set forth in paragraph (a) of this section. The rating board will include a full statement as to the veteran's service-connected disabilities, employment history, educational and vocational attainment and all other factors having a bearing on the issue.

[40 FR 42535, Sept. 15, 1975, as amended at 54 FR 4281, Jan. 30, 1989; 55 FR 31580, Aug. 3, 1990; 58 FR 39664, July 26, 1993; 61 FR 52700, Oct. 8, 1996; 79 FR 2100, Jan. 13, 2014]

§4.17 Total disability ratings for pension based on unemployability and age of the individual.

All veterans who are basically eligible and who are unable to secure and follow a substantially gainful occupation by reason of disabilities which are likely to be permanent shall be rated as permanently and totally disabled. For the purpose of pension, the permanence of the percentage requirements of §4.16 is a requisite. When the percentage requirements are met, and the disabilities involved are of a permanent nature, a rating of permanent and total disability will be assigned if the veteran is found to be unable to secure and follow substantially gainful employment by reason of such disability. Prior employment or unemployment status is immaterial if in the judgment of the rating board the veteran's disabilities render him or her unemployable. In making such determinations, the following guidelines will be used:

- (a) Marginal employment, for example, as a self-employed farmer or other person, while employed in his or her own business, or at odd jobs or while employed at less than half the usual remuneration will not be considered incompatible with a determination of unemployability, if the restriction, as to securing or retaining better employment, is due to disability.
- (b) Claims of all veterans who fail to meet the percentage standards but who meet the basic entitlement criteria and are unemployable, will be referred by the rating board to the Veterans Service Center Manager or the Pension

§4.17a

Management Center Manager under §3.321(b)(2) of this chapter.

(Authority: 38 U.S.C. 1155; 38 U.S.C. 3102)

[43 FR 45348, Oct. 2, 1978, as amended at 56 FR 57985, Nov. 15, 1991; 71 FR 28586, May 17, 2006; 74 FR 26959, June 5, 2009]

§ 4.17a Misconduct etiology.

A permanent and total disability rating under the provisions of §§4.15, 4.16 and 4.17 will not be precluded by reason of the coexistence of misconduct disability when:

- (a) A veteran, regardless of employment status, also has innocently acquired 100 percent disability, or
- (b) Where unemployable, the veteran has other disabilities innocently acquired which meet the percentage requirements of §§4.16 and 4.17 and would render, in the judgment of the rating agency, the average person unable to secure or follow a substantially gainful occupation.

 $[40~{\rm FR}~42536,~{\rm Sept.}~15,~1975,~{\rm as~amended~at}~43~{\rm FR}~45349,~{\rm Oct.}~2,~1978]$

§4.18 Unemployability.

A veteran may be considered as unemployable upon termination of employment which was provided on account of disability, or in which special consideration was given on account of the same, when it is satisfactorily shown that he or she is unable to secure further employment. With amputations, sequelae of fractures and other residuals of traumatism shown to be of static character, a showing of continuous unemployability from date of incurrence, or the date the condition reached the stabilized level, is a general requirement in order to establish the fact that present unemployability is the result of the disability. However, consideration is to be given to the circumstances of employment in individual claims, and, if the employment was only occasional, intermittent, tryout or unsuccessful, or eventually terminated on account of the disability, present unemployability may be attributed to the static disability. Where unemployability for pension previously has been established on the basis of combined service-connected and nonservice-connected disabilities and the service-connected disability or disabilities have increased in severity, §4.16 is for consideration.

[40 FR 42536, Sept. 15, 1975, as amended at 43 FR 45349, Oct. 2, 1978]

§4.19 Age in service-connected claims.

Age may not be considered as a factor in evaluating service-connected disability; and unemployability, in service-connected claims, associated with advancing age or intercurrent disability, may not be used as a basis for a total disability rating. Age, as such, is a factor only in evaluations of disability not resulting from service, *i.e.*, for the purposes of pension.

[29 FR 6718, May 22, 1964, as amended at 43 FR 45349, Oct. 2, 1978]

§ 4.20 Analogous ratings.

When an unlisted condition is encountered it will be permissible to rate under a closely related disease or injury in which not only the functions affected, but the anatomical localization and symptomatology are closely analogous. Conjectural analogies will be avoided, as will the use of analogous ratings for conditions of doubtful diagnosis, or for those not fully supported by clinical and laboratory findings. Nor will ratings assigned to organic diseases and injuries be assigned by analogy to conditions of functional origin.

§4.21 Application of rating schedule.

In view of the number of atypical instances it is not expected, especially with the more fully described grades of disabilities, that all cases will show all the findings specified. Findings sufficiently characteristic to identify the disease and the disability therefrom, and above all, coordination of rating with impairment of function will, however, be expected in all instances.

[41 FR 11293, Mar. 18, 1976]

§ 4.22 Rating of disabilities aggravated by active service.

In cases involving aggravation by active service, the rating will reflect only the degree of disability over and above the degree existing at the time of entrance into the active service, whether the particular condition was noted at the time of entrance into the active service, or it is determined upon the

evidence of record to have existed at that time. It is necessary therefore, in all cases of this character to deduct from the present degree of disability the degree, if ascertainable, of the disability existing at the time of entrance into active service, in terms of the rating schedule, except that if the disability is total (100 percent) no deduction will be made. The resulting difference will be recorded on the rating sheet. If the degree of disability at the time of entrance into the service is not ascertainable in terms of the schedule, no deduction will be made.

§ 4.23 Attitude of rating officers.

It is to be remembered that the majority of applicants are disabled persons who are seeking benefits of law to which they believe themselves entitled. In the exercise of his or her functions, rating officers must not allow their personal feelings to intrude; an antagonistic, critical, or even abusive attitude on the part of a claimant should not in any instance influence the officers in the handling of the case. Fairness and courtesy must at all times be shown to applicants by all employees whose duties bring them in contact, directly or indirectly, with the Department's claimants.

[41 FR 11292, Mar. 18, 1976]

§ 4.24 Correspondence.

All correspondence relative to the interpretation of the schedule for rating disabilities, requests for advisory opinions, questions regarding lack of clarity or application to individual cases involving unusual difficulties, will be addressed to the Director, Compensation Service. A clear statement will be made of the point or points upon which information is desired, and the complete case file will be simultaneously forwarded to Central Office. Rating agencies will assure themselves that the recent report of physical examination presents an adequate picture of the claimant's condition. Claims in regard to which the schedule evaluations are considered inadequate or excessive, and errors in the schedule will be similarly brought to attention.

[41 FR 11292, Mar. 18, 1976, as amended at 79 FR 2100, Jan. 13, 2014]

§ 4.25 Combined ratings table.

Table I, Combined Ratings Table, results from the consideration of the efficiency of the individual as affected first by the most disabling condition, then by the less disabling condition, then by other less disabling conditions, if any, in the order of severity. Thus, a person having a 60 percent disability is considered 40 percent efficient. Proceeding from this 40 percent efficiency, the effect of a further 30 percent disability is to leave only 70 percent of the efficiency remaining after consideration of the first disability, or 28 percent efficiency altogether. The individual is thus 72 percent disabled, as shown in table I opposite 60 percent and under 30 percent.

(a) To use table I, the disabilities will first be arranged in the exact order of their severity, beginning with the greatest disability and then combined with use of table I as hereinafter indicated. For example, if there are two disabilities, the degree of one disability will be read in the left column and the degree of the other in the top row, whichever is appropriate. The figures appearing in the space where the column and row intersect will represent the combined value of the two. This combined value will then be converted to the nearest number divisible by 10. and combined values ending in 5 will be adjusted upward. Thus, with a 50 percent disability and a 30 percent disability, the combined value will be found to be 65 percent, but the 65 percent must be converted to 70 percent to represent the final degree of disability. Similarly, with a disability of 40 percent, and another disability of 20 percent, the combined value is found to be 52 percent, but the 52 percent must be converted to the nearest degree divisible by 10, which is 50 percent. If there are more than two disabilities, the disabilities will also be arranged in the exact order of their severity and the combined value for the first two will be found as previously described for two disabilities. The combined value, exactly as found in table I, will be combined with the degree of the third disability (in order of severity). The combined value for the three disabilities will be found in the space where the column and row intersect, and if there

are only three disabilities will be converted to the nearest degree divisible by 10, adjusting final 5's upward. Thus, if there are three disabilities ratable at 60 percent, 40 percent, and 20 percent, respectively, the combined value for the first two will be found opposite 60 and under 40 and is 76 percent. This 76 will be combined with 20 and the combined value for the three is 81 percent. This combined value will be converted to the nearest degree divisible by 10 which is 80 percent. The same procedure will be employed when there are four or more disabilities. (See table I).

(b) Except as otherwise provided in this schedule, the disabilities arising from a single disease entity, e.g., arthritis, multiple sclerosis, cerebrovascular accident, etc., are to be rated separately as are all other disabiling conditions, if any. All disabilities are then to be combined as described in paragraph (a) of this section. The conversion to the nearest degree divisible by 10 will be done only once per rating decision, will follow the combining of all disabilities, and will be the last procedure in determining the combined degree of disability.

TABLE I—COMBINED RATINGS TABLE
[10 combined with 10 is 19]

	10	20	30	40	50	60	70	80	90
_	.=		40						
	27 28	35	43	51	60	68	76 76	84 84	92 92
	 28 29	36 37	44 45	52 53	60 61	68 68	76	84	92
	30	38	45	53	61	69	77	84	92
	31	38	46	54	62	69	77	85	92
	32	39	47	54	62	70	77	85	92
	33	40	48	55	63	70	78	85	93
	 33	41	48	56	63	70	78	85	93
	 34	42	49	56	64	71	78	85	93
	 35	42	50	57	64	71	78	86	93
	 36	43	50	57	65	72	79	86	93
	37	44	51	58	65	72	79	86	93
	38	45	52	59	66	72	79	86	93
	39	46	52	59	66	73	80	86	93
	40	46	53	60	67	73	80	87	93
	41	47	54	60	67	74	80	87	93
	42	48	55	61	68	74	81	87	94
	 42	49	55	62	68	74	81	87	94
	 43	50	56	62	69	75	81	87	94
	 44	50	57	63	69	75	81	88	94
39	 45	51	57	63	70	76	82	88	94
40	 46	52	58	64	70	76	82	88	94
41	 47	53	59	65	71	76	82	88	94
42	 48	54	59	65	71	77	83	88	94
43	 49	54	60	66	72	77	83	89	94
44	 50	55	61	66	72	78	83	89	94
45	 51	56	62	67	73	78	84	89	95
46	 51	57	62	68	73	78	84	89	95
47	 52	58	63	68	74	79	84	89	95
48	 53	58	64	69	74	79	84	90	95
49	 54	59	64	69	75	80	85	90	95
50	 55	60	65	70	75	80	85	90	95
	 56	61	66	71	76	80	85	90	95
	 57	62	66	71	76	81	86	90	95
	 58	62	67	72	77	81	86	91	95
	 59	63	68	72	77	82	86	91	95
	 60	64	69	73	78	82	87	91	96
	 60	65	69	74	78	82	87	91	96
	 61	66	70	74	79	83	87	91	96
	 62	66	71	75	79	83	87	92	96
	 63	67	71	75	80	84	88	92	96
	 64	68	72	76 77	80	84	88	92	96
	65	69	73	77	81	84	88	92	96
	66	70 70	73 74	77	81	85	89	92 93	96
	 67		1	78 70	82	85	89		96
	68 69	71 72	75 76	78 79	82	86 86	89 90	93 93	96
					83				97
	 69	73	76	80	83	86	90	93	97
	70	74 74	77	80	84	87	90	93	97
	71 72	74 75	78	81	84	87 88	90	94	97
69	 ı /2	/5	78	81	85	88	91	94	97

Department of Veterans Affairs

TABLE I—COMBINED RATINGS TABLE—Continued
[10 combined with 10 is 19]

	10	20	30	40	50	60	70	80	90
70	73	76	79	82	85	88	91	94	97
71	74	77	80	83	86	88	91	94	97
72	75	78	80	83	86	89	92	94	97
73	76	78	81	84	87	89	92	95	97
74	77	79	82	84	87	90	92	95	97
75	78	80	83	85	88	90	93	95	98
76	78	81	83	86	88	90	93	95	98
77	79	82	84	86	89	91	93	95	98
78	80	82	85	87	89	91	93	96	98
79	81	83	85	87	90	92	94	96	98
80	82	84	86	88	90	92	94	96	98
81	83	85	87	89	91	92	94	96	98
82	84	86	87	89	91	93	95	96	98
83	85	86	88	90	92	93	95	97	98
84	86	87	89	90	92	94	95	97	98
85	87	88	90	91	93	94	96	97	99
86	87	89	90	92	93	94	96	97	99
87	88	90	91	92	94	95	96	97	99
88	89	90	92	93	94	95	96	98	99
89	90	91	92	93	95	96	97	98	99
90	91	92	93	94	95	96	97	98	99
91	92	93	94	95	96	96	97	98	99
92	93	94	94	95	96	97	98	98	99
93	94	94	95	96	97	97	98	99	99
94	95	95	96	96	97	98	98	99	99
	ı		1		1		1	I	l

(Authority: 38 U.S.C. 1155)

[41 FR 11293, Mar. 18, 1976, as amended at 54 FR 27161, June 28, 1989; 54 FR 36029, Aug. 31, 1989; 83 FR 17756, Apr. 24, 2018]

§ 4.26 Bilateral factor.

When a partial disability results from disease or injury of both arms, or of both legs, or of paired skeletal muscles, the ratings for the disabilities of the right and left sides will be combined as usual, and 10 percent of this value will be added (i.e., not combined) before proceeding with further combinations, or converting to degree of disability. The bilateral factor will be applied to such bilateral disabilities before other combinations are carried out and the rating for such disabilities including the bilateral factor in this section will be treated as 1 disability for the purpose of arranging in order of severity and for all further combinations. For example, with disabilities evaluated at 60 percent, 20 percent, 10 percent and 10 percent (the two 10's representing bilateral disabilities), the order of severity would be 60, 21 and 20. The 60 and 21 combine to 68 percent and the 68 and 20 to 74 percent, converted to 70 percent as the final degree of disability.

- (a) The use of the terms "arms" and "legs" is not intended to distinguish between the arm, forearm and hand, or the thigh, leg, and foot, but relates to the upper extremities and lower extremities as a whole. Thus with a compensable disability of the right thigh, for example, amputation, and one of the left foot, for example, pes planus, the bilateral factor applies, and similarly whenever there are compensable disabilities affecting use of paired extremities regardless of location or specified type of impairment.
- (b) The correct procedure when applying the bilateral factor to disabilities affecting both upper extremities and both lower extremities is to combine the ratings of the disabilities affecting the 4 extremities in the order of their individual severity and apply the bilateral factor by adding, not combining, 10 percent of the combined value thus attained.
- (c) The bilateral factor is not applicable unless there is partial disability of compensable degree in each of 2 paired extremities, or paired skeletal muscles.

§4.27 Use of diagnostic code numbers.

The diagnostic code numbers appearing opposite the listed ratable disabilities are arbitrary numbers for the purpose of showing the basis of the evaluation assigned and for statistical analysis in the Department of Veterans Affairs, and as will be observed, extend from 5000 to a possible 9999. Great care will be exercised in the selection of the applicable code number and in its citation on the rating sheet. No other numbers than these listed or hereafter furnished are to be employed for rating purposes, with an exception as described in this section, as to unlisted conditions. When an unlisted disease, injury, or residual condition is encountered, requiring rating by analogy, the diagnostic code number will be "builtup" as follows: The first 2 digits will be selected from that part of the schedule most closely identifying the part, or system, of the body involved; the last 2 digits will be "99" for all unlisted conditions. This procedure will facilitate a close check of new and unlisted conditions, rated by analogy. In the selection of code numbers, injuries will generally be represented by the number assigned to the residual condition on the basis of which the rating is determined. With diseases, preference is to be given to the number assigned to the disease itself; if the rating is determined on the basis of residual conditions, the number appropriate to the residual condition will be added, preceded by a hyphen. Thus, rheumatoid (atrophic) arthritis rated as ankylosis of the lumbar spine should be coded "5002-5240." In this way, the exact source of each rating can be easily identified. In the citation of disabilities on rating sheets, the diagnostic terminology will be that of the medical examiner, with no attempt to translate the terms into schedule nomenclature. Residuals of diseases or therapeutic procedures will not be cited without reference to the basic disease.

[41 FR 11293, Mar. 18, 1976, as amended at 70 FR 75399, Dec. 20, 2005]

§ 4.28 Prestabilization rating from date of discharge from service.

The following ratings may be assigned, in lieu of ratings prescribed

elsewhere, under the conditions stated for disability from any disease or injury. The prestabilization rating is not to be assigned in any case in which a total rating is immediately assignable under the regular provisions of the schedule or on the basis of individual unemployability. The prestabilization 50-percent rating is not to be used in any case in which a rating of 50 percent or more is immediately assignable under the regular provisions.

	Rating
Unstabilized condition with severe disability— Substantially gainful employment is not fea- sible or advisable	100
juries— Material impairment of employability likely	50

Note (1): Department of Veterans Affairs examination is not required prior to assignment of prestabilization ratings; however, the fact that examination was accomplished will not preclude assignment of these benefits. Prestabilization ratings are for assignment in the immediate postdischarge period. They will continue for a 12-month period following discharge from service. However, prestabilization ratings may be changed to a regular schedular total rating or one authorizing a greater benefit at any time. In each prestabilization rating an examination will be requested to be accomplished not earlier than 6 months nor more than 12 months following discharge. In those prestabilization ratings in which following examination reduction in evaluation is found to be warranted, the higher evaluation will be continued to the end of the 12th month following discharge or to the end of the period provided under §3.105(e) of this chapter, whichever is later. Special monthly compensation should be assigned concurrently in these cases whenever records are adequate to establish entitlement.

NOTE (2): Diagnosis of disease, injury, or residuals will be cited, with diagnostic code number assigned from this rating schedule for conditions listed therein.

[35 FR 11906, July 24, 1970]

§ 4.29 Ratings for service-connected disabilities requiring hospital treatment or observation.

A total disability rating (100 percent) will be assigned without regard to other provisions of the rating schedule when it is established that a service-connected disability has required hospital treatment in a Department of

Veterans Affairs or an approved hospital for a period in excess of 21 days or hospital observation at Department of Veterans Affairs expense for a service-connected disability for a period in excess of 21 days.

- (a) Subject to the provisions of paragraphs (d), (e), and (f) of this section this increased rating will be effective the first day of continuous hospitalization and will be terminated effective the last day of the month of hospital discharge (regular discharge or release to non-bed care) or effective the last day of the month of termination of treatment or observation for the service-connected disability. A temporary release which is approved by an attending Department of Veterans Affairs physician as part of the treatment plan will not be considered an absence.
- (1) An authorized absence in excess of 4 days which begins during the first 21 days of hospitalization will be regarded as the equivalent of hospital discharge effective the first day of such authorized absence. An authorized absence of 4 days or less which results in a total of more than 8 days of authorized absence during the first 21 days of hospitalization will be regarded as the equivalent of hospital discharge effective the ninth day of authorized absence.
- (2) Following a period of hospitalization in excess of 21 days, an authorized absence in excess of 14 days or a third consecutive authorized absence of 14 days will be regarded as the equivalent of hospital discharge and will interrupt hospitalization effective on the last day of the month in which either the authorized absence in excess of 14 days or the third 14 day period begins, except where there is a finding that convalescence is required as provided by paragraph (e) or (f) of this section. The termination of these total ratings will not be subject to §3.105(e) of this chapter.
- (b) Notwithstanding that hospital admission was for disability not connected with service, if during such hospitalization, hospital treatment for a service-connected disability is instituted and continued for a period in excess of 21 days, the increase to a total rating will be granted from the first day of such treatment. If service connection for the disability under treat-

ment is granted after hospital admission, the rating will be from the first day of hospitalization if otherwise in order.

- (c) The assignment of a total disability rating on the basis of hospital treatment or observation will not preclude the assignment of a total disability rating otherwise in order under other provisions of the rating schedule. and consideration will be given to the propriety of such a rating in all instances and to the propriety of its continuance after discharge. Particular attention, with a view to proper rating under the rating schedule, is to be given to the claims of veterans discharged from hospital, regardless of length of hospitalization, with indications on the final summary of expected confinement to bed or house, or to inability to work with requirement of frequent care of physician or nurse at
- (d) On these total ratings Department of Veterans Affairs regulations governing effective dates for increased benefits will control.
- (e) The total hospital rating if convalescence is required may be continued for periods of 1, 2, or 3 months in addition to the period provided in paragraph (a) of this section.
- (f) Extension of periods of 1, 2 or 3 months beyond the initial 3 months may be made upon approval of the Veterans Service Center Manager.
- (g) Meritorious claims of veterans who are discharged from the hospital with less than the required number of days but need post-hospital care and a prolonged period of convalescence will be referred to the Director, Compensation Service, under §3.321(b)(1) of this chapter.

[29 FR 6718, May 22, 1964, as amended at 41 FR 11294, Mar. 18, 1976; 41 FR 34256, Aug. 13, 1976; 54 FR 4281, Jan. 30, 1989; 54 FR 34981, Aug. 23, 1989; 71 FR 28586, May 17, 2006; 79 FR 2100, Jan. 13, 2014]

§ 4.30 Convalescent ratings.

A total disability rating (100 percent) will be assigned without regard to other provisions of the rating schedule when it is established by report at hospital discharge (regular discharge or release to non-bed care) or outpatient release that entitlement is warranted

under paragraph (a) (1), (2) or (3) of this section effective the date of hospital admission or outpatient treatment and continuing for a period of 1, 2, or 3 months from the first day of the month following such hospital discharge or outpatient release. The termination of these total ratings will not be subject to §3.105(e) of this chapter. Such total rating will be followed by appropriate schedular evaluations. When the evidence is inadequate to assign a schedular evaluation, a physical examination will be scheduled and considered prior to the termination of a total rating under this section.

- (a) Total ratings will be assigned under this section if treatment of a service-connected disability resulted in:
- (1) Surgery necessitating at least one month of convalescence (Effective as to outpatient surgery March 1, 1989.)
- (2) Surgery with severe postoperative residuals such as incompletely healed surgical wounds, stumps of recent amputations, therapeutic immobilization of one major joint or more, application of a body cast, or the necessity for house confinement, or the necessity for continued use of a wheelchair or crutches (regular weight-bearing prohibited). (Effective as to outpatient surgery March 1, 1989.)
- (3) Immobilization by cast, without surgery, of one major joint or more. (Effective as to outpatient treatment March 10, 1976.)

A reduction in the total rating will not be subject to §3.105(e) of this chapter. The total rating will be followed by an open rating reflecting the appropriate schedular evaluation; where the evidence is inadequate to assign the schedular evaluation, a physcial examination will be scheduled prior to the end of the total rating period.

- (b) A total rating under this section will require full justification on the rating sheet and may be extended as follows:
- (1) Extensions of 1, 2 or 3 months beyond the initial 3 months may be made under paragraph (a) (1), (2) or (3) of this section.
- (2) Extensions of 1 or more months up to 6 months beyond the initial 6 months period may be made under paragraph (a) (2) or (3) of this section

upon approval of the Veterans Service Center Manager.

[41 FR 34256, Aug. 13, 1976, as amended at 54 FR 4281, Jan. 30, 1989; 71 FR 28586, May 17, 2006]

§4.31 Zero percent evaluations.

In every instance where the schedule does not provide a zero percent evaluation for a diagnostic code, a zero percent evaluation shall be assigned when the requirements for a compensable evaluation are not met.

[58 FR 52018, Oct. 6, 1993]

Subpart B—Disability Ratings

THE MUSCULOSKELETAL SYSTEM

§4.40 Functional loss.

Disability of the musculoskeletal system is primarily the inability, due to damage or infection in parts of the system, to perform the normal working movements of the body with normal excursion, strength, speed, coordination and endurance. It is essential that the examination on which ratings are based adequately portray the anatomical damage, and the functional loss, with respect to all these elements. The functional loss may be due to absence of part, or all, of the necessary bones. joints and muscles, or associated structures, or to deformity, adhesions, defective innervation, or other pathology, or it may be due to pain, supported by adequate pathology and evidenced by the visible behavior of the claimant undertaking the motion. Weakness is as important as limitation of motion, and a part which becomes painful on use must be regarded as seriously disabled. A little used part of the musculoskeletal system may be expected to show evidence of disuse, either through atrophy, the condition of the skin, absence of normal callosity or the like.

§4.41 History of injury.

In considering the residuals of injury, it is essential to trace the medical-industrial history of the disabled person from the original injury, considering

the nature of the injury and the attendant circumstances, and the requirements for, and the effect of, treatment over past periods, and the course of the recovery to date. The duration of the initial, and any subsequent, period of total incapacity, especially periods reflecting delayed union, inflammation, swelling, drainage, or operative intervention, should be given close attention. This consideration, or the absence of clear cut evidence of injury, may result in classifying the disability as not of traumatic origin, either reflecting congenital or developmental etiology, or the effects of healed dis-

§ 4.42 Complete medical examination of injury cases.

The importance of complete medical examination of injury cases at the time of first medical examination by the Department of Veterans Affairs cannot be overemphasized. When possible, this should include complete neurological and psychiatric examination, and other special examinations indicated by the physical condition, in addition to the required general and orthopedic or surgical examinations. When complete examinations are not conducted covering all systems of the body affected by disease or injury, it is impossible to visualize the nature and extent of the service connected disability. Incomplete examination is a common cause of incorrect diagnosis, especially in the neurological and psychiatric fields, and frequently leaves the Department of Veterans Affairs in doubt as to the presence or absence of disabling conditions at the time of the examination.

§ 4.43 Osteomyelitis.

Chronic, or recurring, suppurative osteomyelitis, once clinically identified, including chronic inflammation of bone marrow, cortex, or periosteum, should be considered as a continuously disabling process, whether or not an actively discharging sinus or other obvious evidence of infection is manifest from time to time, and unless the focus is entirely removed by amputation will entitle to a permanent rating to be combined with other ratings for residual conditions, however, not exceeding

amputation ratings at the site of elec-

§ 4.44 The bones.

The osseous abnormalities incident to trauma or disease, such as malunion with deformity throwing abnormal stress upon, and causing malalignment of joint surfaces, should be depicted from study and observation of all available data, beginning with inception of injury or disease, its nature, degree of prostration, treatment and duration of convalescence, and progress of recovery with development of permanent residuals. With shortening of a long bone, some degree of angulation is to be expected; the extent and direction should be brought out by X-ray and observation. The direction of angulation and extent of deformity should be carefully related to strain on the neighboring joints, especially those connected with weight-bearing.

§ 4.45 The joints.

As regards the joints the factors of disability reside in reductions of their normal excursion of movements in different planes. Inquiry will be directed to these considerations:

- (a) Less movement than normal (due to ankylosis, limitation or blocking, adhesions, tendon-tie-up, contracted scars, etc.).
- (b) More movement than normal (from flail joint, resections, nonunion of fracture, relaxation of ligaments, etc.).
- (c) Weakened movement (due to muscle injury, disease or injury of peripheral nerves, divided or lengthened tendons, etc.).
 - (d) Excess fatigability.
- (e) Incoordination, impaired ability to execute skilled movements smoothly.
- (f) Pain on movement, swelling, deformity or atrophy of disuse. Instability of station, disturbance of locomotion, interference with sitting, standing and weight-bearing are related considerations. For the purpose of rating disability from arthritis, the shoulder, elbow, wrist, hip, knee, and ankle are considered major joints; multiple involvements of the interphalangeal, metacarpal and carpal joints of

the upper extremities, the interphalangeal, metatarsal and tarsal joints of the lower extremities, the cervical vertebrae, the dorsal vertebrae, and the lumbar vertebrae, are considered groups of minor joints, ratable on a parity with major joints. The lumbosacral articulation and both sacrolliac joints are considered to be a group of minor joints, ratable on disturbance of lumbar spine functions.

§ 4.46 Accurate measurement.

Accurate measurement of the length of stumps, excursion of joints, dimensions and location of scars with respect to landmarks, should be insisted on. The use of a goniometer in the measurement of limitation of motion is indispensable in examinations conducted within the Department of Veterans Affairs. Muscle atrophy must also be accurately measured and reported.

[41 FR 11294, Mar. 18, 1976]

§§ 4.47-4.54 [Reserved]

§ 4.55 Principles of combined ratings for muscle injuries.

- (a) A muscle injury rating will not be combined with a peripheral nerve paralysis rating of the same body part, unless the injuries affect entirely different functions.
- (b) For rating purposes, the skeletal muscles of the body are divided into 23 muscle groups in 5 anatomical regions: 6 muscle groups for the shoulder girdle and arm (diagnostic codes 5301 through 5306); 3 muscle groups for the forearm and hand (diagnostic codes 5307 through 5309); 3 muscle groups for the foot and leg (diagnostic codes 5310 through 5312); 6 muscle groups for the pelvic girdle and thigh (diagnostic codes 5313 through 5318); and 5 muscle groups for the torso and neck (diagnostic codes 5319 through 5323).
- (c) There will be no rating assigned for muscle groups which act upon an ankylosed joint, with the following exceptions:
- (1) In the case of an ankylosed knee, if muscle group XIII is disabled, it will be rated, but at the next lower level than that which would otherwise be assigned.
- (2) In the case of an ankylosed shoulder, if muscle groups I and II are se-

verely disabled, the evaluation of the shoulder joint under diagnostic code 5200 will be elevated to the level for unfavorable ankylosis, if not already assigned, but the muscle groups themselves will not be rated.

- (d) The combined evaluation of muscle groups acting upon a single unankylosed joint must be lower than the evaluation for unfavorable ankylosis of that joint, except in the case of muscle groups I and II acting upon the shoulder.
- (e) For compensable muscle group injuries which are in the same anatomical region but do not act on the same joint, the evaluation for the most severely injured muscle group will be increased by one level and used as the combined evaluation for the affected muscle groups.
- (f) For muscle group injuries in different anatomical regions which do not act upon ankylosed joints, each muscle group injury shall be separately rated and the ratings combined under the provisions of §4.25.

(Authority: 38 U.S.C. 1155) [62 FR 30237, June 3, 1997]

§ 4.56 Evaluation of muscle disabilities.

- (a) An open comminuted fracture with muscle or tendon damage will be rated as a severe injury of the muscle group involved unless, for locations such as in the wrist or over the tibia, evidence establishes that the muscle damage is minimal.
- (b) A through-and-through injury with muscle damage shall be evaluated as no less than a moderate injury for each group of muscles damaged.
- (c) For VA rating purposes, the cardinal signs and symptoms of muscle disability are loss of power, weakness, lowered threshold of fatigue, fatigue-pain, impairment of coordination and uncertainty of movement.
- (d) Under diagnostic codes 5301 through 5323, disabilities resulting from muscle injuries shall be classified as slight, moderate, moderately severe or severe as follows:
- (1) Slight disability of muscles—(i) Type of injury. Simple wound of muscle without debridement or infection.

- (ii) History and complaint. Service department record of superficial wound with brief treatment and return to duty. Healing with good functional results. No cardinal signs or symptoms of muscle disability as defined in paragraph (c) of this section.
- (iii) Objective findings. Minimal scar. No evidence of fascial defect, atrophy, or impaired tonus. No impairment of function or metallic fragments retained in muscle tissue.
- (2) Moderate disability of muscles—(i) Type of injury. Through and through or deep penetrating wound of short track from a single bullet, small shell or shrapnel fragment, without explosive effect of high velocity missile, residuals of debridement, or prolonged infection.
- (ii) History and complaint. Service department record or other evidence of in-service treatment for the wound. Record of consistent complaint of one or more of the cardinal signs and symptoms of muscle disability as defined in paragraph (c) of this section, particularly lowered threshold of fatigue after average use, affecting the particular functions controlled by the injured muscles.
- (iii) Objective findings. Entrance and (if present) exit scars, small or linear, indicating short track of missile through muscle tissue. Some loss of deep fascia or muscle substance or impairment of muscle tonus and loss of power or lowered threshold of fatigue when compared to the sound side.
- (3) Moderately severe disability of muscles—(i) Type of injury. Through and through or deep penetrating wound by small high velocity missile or large low-velocity missile, with debridement, prolonged infection, or sloughing of soft parts, and intermuscular scarring.
- (ii) History and complaint. Service department record or other evidence showing hospitalization for a prolonged period for treatment of wound. Record of consistent complaint of cardinal signs and symptoms of muscle disability as defined in paragraph (c) of this section and, if present, evidence of inability to keep up with work requirements.
- (iii) Objective findings. Entrance and (if present) exit scars indicating track of missile through one or more muscle

- groups. Indications on palpation of loss of deep fascia, muscle substance, or normal firm resistance of muscles compared with sound side. Tests of strength and endurance compared with sound side demonstrate positive evidence of impairment.
- (4) Severe disability of muscles—(i) Type of injury. Through and through or deep penetrating wound due to high-velocity missile, or large or multiple low velocity missiles, or with shattering bone fracture or open comminuted fracture with extensive debridement, prolonged infection, or sloughing of soft parts, intermuscular binding and scarring.
- (ii) History and complaint. Service department record or other evidence showing hospitalization for a prolonged period for treatment of wound. Record of consistent complaint of cardinal signs and symptoms of muscle disability as defined in paragraph (c) of this section, worse than those shown for moderately severe muscle injuries, and, if present, evidence of inability to keep up with work requirements.
- (iii) Objective findings. Ragged, depressed and adherent scars indicating wide damage to muscle groups in missile track. Palpation shows loss of deep fascia or muscle substance, or soft flabby muscles in wound area. Muscles swell and harden abnormally in contraction. Tests of strength, endurance, or coordinated movements compared with the corresponding muscles of the uninjured side indicate severe impairment of function. If present, the following are also signs of severe muscle disability:
- (A) X-ray evidence of minute multiple scattered foreign bodies indicating intermuscular trauma and explosive effect of the missile.
- (B) Adhesion of scar to one of the long bones, scapula, pelvic bones, sacrum or vertebrae, with epithelial sealing over the bone rather than true skin covering in an area where bone is normally protected by muscle.
- (C) Diminished muscle excitability to pulsed electrical current in electrodiagnostic tests.
 - (D) Visible or measurable atrophy.
- (E) Adaptive contraction of an opposing group of muscles.
- (F) Atrophy of muscle groups not in the track of the missile, particularly of

the trapezius and serratus in wounds of the shoulder girdle.

(G) Induration or atrophy of an entire muscle following simple piercing by a projectile.

(Authority: 38 U.S.C. 1155 [62 FR 30238, June 3, 1997]

§ 4.57 Static foot deformities.

It is essential to make an initial distinction between bilateral flatfoot as a congenital or as an acquired condition. The congenital condition, with depression of the arch, but no evidence of abnormal callosities, areas of pressure, strain or demonstrable tenderness, is a congenital abnormality which is not compensable or pensionable. In the acquired condition, it is to be remembered that depression of the longitudinal arch, or the degree of depression, is not the essential feature. The attention should be given to anatomical changes, as compared to normal, in the relationship of the foot and leg. particularly to the inward rotation of the superior portion of the os calcis, medial deviation of the insertion of the Achilles tendon, the medial tilting of the upper border of the astragalus. This is an unfavorable mechanical relationship of the parts. A plumb line dropped from the middle of the patella falls inside of the normal point. The forepart of the foot is abducted, and the foot everted. The plantar surface of the foot is painful and shows demonstrable tenderness, and manipulation of the foot produces spasm of the Achilles tendon, peroneal spasm due to adhesion about the peroneal sheaths, and other evidence of pain and limited motion. The symptoms should be apparent without regard to exercise. In severe cases there is gaping of bones on the inner border of the foot, and rigid valgus position with loss of the power of inversion and adduction. Exercise with undeveloped or unbalanced musculature, producing chronic irritation. can be an aggravating factor. In the absence of trauma or other definite evidence of aggravation, service connection is not in order for pes cavus which is a typically congenital or juvenile disease.

§4.58 Arthritis due to strain.

With service incurred lower extremity amputation or shortening, a disabling arthritis, developing in the same extremity, or in both lower extremities, with indications of earlier, or more severe, arthritis in the injured extremity, including also arthritis of the lumbosacral joints and lumbar spine, if associated with the leg amputation or shortening, will be considered as service incurred, provided, however, that arthritis affecting joints not directly subject to strain as a result of the service incurred amputation will not be granted service connection. This will generally require separate evaluation of the arthritis in the joints directly subject to strain. Amputation, or injury to an upper extremity, is not considered as a causative factor with subsequently developing arthritis, except in joints subject to direct strain or actually injured.

§ 4.59 Painful motion.

With any form of arthritis, painful motion is an important factor of disability, the facial expression, wincing, etc., on pressure or manipulation, should be carefully noted and definitely related to affected joints. Muscle spasm will greatly assist the identification. Sciatic neuritis is not uncommonly caused by arthritis of the spine. The intent of the schedule is to recognize painful motion with joint or periarticular pathology as productive of disability. It is the intention to recognize actually painful, unstable, or malaligned joints, due to healed injury, as entitled to at least the minimum compensable rating for the joint. Crepitation either in the soft tissues such as the tendons or ligaments, or crepitation within the joint structures should be noted carefully as points of contact which are diseased. Flexion elicits such manifestations. The joints involved should be tested for pain on both active and passive motion, in weight-bearing and nonweight-bearing and, if possible, with the range of the opposite undamaged joint.

§4.60 [Reserved]

§ 4.61 Examination.

With any form of arthritis (except traumatic arthritis) it is essential that the examination for rating purposes cover all major joints, with especial reference to Heberden's or Haygarth's nodes.

§ 4.62 Circulatory disturbances.

The circulatory disturbances, especially of the lower extremity following injury in the popliteal space, must not be overlooked, and require rating generally as phlebitis.

§ 4.63 Loss of use of hand or foot.

Loss of use of a hand or a foot, for the purpose of special monthly compensation, will be held to exist when no effective function remains other than that which would be equally well served by an amputation stump at the site of election below elbow or knee with use of a suitable prosthetic appliance. The determination will be made on the basis of the actual remaining function of the hand or foot, whether the acts of grasping, manipulation, etc., in the case of the hand, or of balance and propulsion, etc., in the case of the foot, could be accomplished equally well by an amputation stump with prosthesis.

(a) Extremely unfavorable complete ankylosis of the knee, or complete ankylosis of 2 major joints of an extremity, or shortening of the lower extremity of 3½ inches (8.9 cms.) or more, will be taken as loss of use of the hand or foot involved.

(b) Complete paralysis of the external popliteal nerve (common peroneal) and consequent, footdrop, accompanied by characteristic organic changes including trophic and circulatory disturbances and other concomitants confirmatory of complete paralysis of this nerve, will be taken as loss of use of the foot.

[29 FR 6718, May 22, 1964, as amended at 43 FR 45349, Oct. 2, 1978]

§ 4.64 Loss of use of both buttocks.

Loss of use of both buttocks shall be deemed to exist when there is severe damage to muscle Group XVII, bilat-

eral (diagnostic code number 5317) and additional disability rendering it impossible for the disabled person, without assistance, to rise from a seated position and from a stooped position (fingers to toes position) and to maintain postural stability (the pelvis upon head of femur). The assistance may be rendered by the person's own hands or arms, and, in the matter of postural stability, by a special appliance.

§4.65 [Reserved]

§ 4.66 Sacroiliac joint.

The common cause of disability in this region is arthritis, to be identified in the usual manner. The lumbosacral and sacroiliac joints should be considered as one anatomical segment for rating purposes. X-ray changes from arthritis in this location are decrease or obliteration of the joint space, with the appearance of increased bone density of the sacrum and ilium and sharpening of the margins of the joint. Disability is manifest from erector spinae spasm (not accounted for by other pathology), tenderness on deep palpation and percussion over these joints, loss of normal quickness of motion and resiliency, and postural defects often accompanied by limitation of flexion and extension of the hip. Traumatism is a rare cause of disability in this connection, except when superimposed upon congenital defect or upon an existent arthritis; to permit assumption of pure traumatic origin, objective evidence of damage to the joint, and history of trauma sufficiently severe to injure this extremely strong and practically immovable joint is required. There should be careful consideration of lumbosacral sprain, and the various symptoms of pain and paralysis attributable to disease affecting the lumbar vertebrae and the intervertebral disc.

§ 4.67 Pelvic bones.

The variability of residuals following these fractures necessitates rating on specific residuals, faulty posture, limitation of motion, muscle injury, painful motion of the lumbar spine, manifest by muscle spasm, mild to moderate sciatic neuritis, peripheral nerve injury, or limitation of hip motion.

§ 4.68 Amputation rule.

The combined rating for disabilities of an extremity shall not exceed the rating for the amputation at the elective level, were amputation to be performed. For example, the combined evaluations for disabilities below the knee shall not exceed the 40 percent evaluation, diagnostic code 5165. This 40 percent rating may be further combined with evaluation for disabilities above the knee but not to exceed the above the knee amputation elective level. Painful neuroma of a stump after amputation shall be assigned the evaluation for the elective site of reamputation.

§ 4.69 Dominant hand.

Handedness for the purpose of a dominant rating will be determined by the evidence of record, or by testing on VA examination. Only one hand shall be considered dominant. The injured hand, or the most severely injured hand, of an ambidextrous individual will be considered the dominant hand for rating purposes.

(Authority: 38 U.S.C. 1155) [62 FR 30239, June 3, 1997]

§ 4.70 Inadequate examinations.

If the report of examination is inadequate as a basis for the required consideration of service connection and evaluation, the rating agency may re-

quest a supplementary report from the examiner giving further details as to the limitations of the disabled person's ordinary activity imposed by the disease, injury, or residual condition, the prognosis for return to, or continuance of, useful work. When the best interests of the service will be advanced by personal conference with the examiner, such conference may be arranged through channels.

§ 4.71 Measurement of ankylosis and joint motion.

Plates I and II provide a standardized description of ankylosis and joint motion measurement. The anatomical position is considered as 0°, with two major exceptions: (a) Shoulder rotation—arm abducted to 90°, elbow flexed to 90° with the position of the forearm reflecting the midpoint 0° between internal and external rotation of the shoulder; and (b) supination and pronation—the arm next to the body. elbow flexed to 90°, and the forearm in midposition 0° between supination and pronation. Motion of the thumb and fingers should be described by appropriate reference to the joints (See Plate III) whose movement is limited, with a statement as to how near, in centimeters, the tip of the thumb can approximate the fingers, or how near the tips of the fingers can approximate the proximal transverse crease of palm.




PLATE I




PLATE II

 $[29 \ FR \ 6718, May \ 22, \ 1964, as amended at 43 \ FR \ 45349, Oct. \ 2, \ 1978; \ 67 \ FR \ 48785, July \ 26, \ 2002]$

§ 4.71a Schedule of ratings—musculoskeletal system.

ACUTE, SUBACUTE, OR CHRONIC DISEASES

	Rat- ing
5000 Osteomyelitis, acute, subacute, or chronic: Of the pelvis, vertebrae, or extending into major joints, or with multiple localization or with long history of intractability and debility, anemia, amyloid liver changes, or other continuous	
constitutional symptoms	100
Frequent episodes, with constitutional symptoms	60
With definite involucrum or sequestrum, with or	
without discharging sinus	30
With discharging sinus or other evidence of ac-	
tive infection within the past 5 years	20
Inactive, following repeated episodes, without	
evidence of active infection in past 5 years	10

ACUTE, SUBACUTE, OR CHRONIC DISEASES—Continued

Rat-

NOTE (1): A rating of 10 percent, as an exception to the amputation rule, is to be assigned in any case of active osteomyelitis where the amputation rating for the affected part is no percent. This 10 percent rating and the other partial ratings of 30 percent reless are to be combined with ratings for ankylosis, limited motion, nonunion or malunion, shortening, etc., subject, of course, to the amputation rule. The 60 percent rating, as it is based on constitutional symptoms, is not subject to the amputation rule. A rating for osteomyelitis will not be applied following cure by removal or radical resection of the affected bone.

Department of Veterans Affairs

§4.71a

Acute, Subacute, or Chronic Diseases—Continued

	Rat- ing		Rat- ing
NOTE (2): The 20 percent rating on the basis of		With X-ray evidence of involvement	
activity within the past 5 years is not assign-		of 2 or more major joints or 2 or	
able following the initial infection of active os-		more minor joint groups, with oc-	
teomyelitis with no subsequent reactivation.		casional incapacitating exacer-	
The prerequisite for this historical rating is an		bations	20
established recurrent osteomyelitis. To qualify for the 10 percent rating, 2 or more episodes		With X-ray evidence of involvement of 2 or more major joints or 2 or	
following the initial infection are required. This		more minor joint groups	10
20 percent rating or the 10 percent rating,		Note (1): The 20 pct and 10 pct ratings	
when applicable, will be assigned once only to		based on X-ray findings, above, will not	
cover disability at all sites of previously active		be combined with ratings based on limita-	
infection with a future ending date in the case		tion of motion.	
of the 20 percent rating.		Note (2): The 20 pct and 10 pct ratings	
001 Bones and joints, tuberculosis of, active or inactive:		based on X-ray findings, above, will not	
Active	100	be utilized in rating conditions listed under diagnostic codes 5013 to 5024, inclusive.	
Inactive: See §§ 4.88c and 4.89.	100	5004 Arthritis, gonorrheal.	
Multi-joint arthritis (except post-traumatic and		5005 Arthritis, pneumococcic.	
gout), 2 or more joints, as an active process:		5006 Arthritis, typhoid.	
With constitutional manifestations associ-		5007 Arthritis, syphilitic.	
ated with active joint involvement, totally		5008 Arthritis, streptococcic.	
incapacitating	100	5009 Other specified forms of arthropathy (exclud-	
Less than criteria for 100% but with weight		ing gout).	
loss and anemia productive of severe im-		Note (1): Other specified forms of arthrop-	
pairment of health or severely incapaci-		athy include, but are not limited to,	
tating exacerbations occurring 4 or more times a year or a lesser number over pro-		Charcot neuropathic, hypertrophic, crys-	
longed periods	60	talline, and other autoimmune arthropathies.	
Symptom combinations productive of defi-		Note (2): With the types of arthritis, diag-	
nite impairment of health objectively sup-		nostic codes 5004 through 5009, rate the	
ported by examination findings or inca-		acute phase under diagnostic code 5002;	
pacitating exacerbations occurring 3 or		rate any chronic residuals under diag-	
more times a year	40	nostic code 5003.	
One or two exacerbations a year in a well-	20	5010 Post-traumatic arthritis: Rate as limitation of	
established diagnosis Note (1): Examples of conditions rated	20	motion, dislocation, or other specified instability	
using this diagnostic code include, but are		under the affected joint. If there are 2 or more joints affected, each rating shall be combined in	
not limited to, rheumatoid arthritis, psori-		accordance with § 4.25.	
atic arthritis, and spondyloarthropathies.		5011 Decompression illness: Rate manifestations	
Note (2): For chronic residuals, rate under		under the appropriate diagnostic code within the	
diagnostic code 5003.		affected body system, such as arthritis for mus-	
Note (3): The ratings for the active process		culoskeletal residuals; auditory system for vestib- ular residuals; respiratory system for pulmonary	
will not be combined with the residual rat- ings for limitation of motion, ankylosis, or		barotrauma residuals; and neurologic system for	
diagnostic code 5003. Instead, assign the		cerebrovascular accident residuals.	
higher evaluation.		5012 Bones, neoplasm, malignant, primary or sec-	
03 Degenerative arthritis, other than post-trau-		ondary	10
matic:		Note: The 100 percent rating will be contin-	
Degenerative arthritis established by X-ray		ued for 1 year following the cessation of surgical, X-ray, antineoplastic chemo-	
findings will be rated on the basis of limi-		therapy or other prescribed therapeutic	
tation of motion under the appropriate di- agnostic codes for the specific joint or		procedure. If there has been no local re-	
joints involved (DC 5200 etc.). When how-		currence or metastases, rate based on re-	
ever, the limitation of motion of the spe-		siduals.	
cific joint or joints involved is noncompen-		5013 Osteoporosis, residuals of.	
sable under the appropriate diagnostic		5014 Osteomalacia, residuals of.	
codes, a rating of 10 pct is for application		5015 Bones, neoplasm, benign.	
for each such major joint or group of minor joints affected by limitation of mo-		5016 Osteitis deformans.	
tion, to be combined, not added under di-		5017 Gout.	
agnostic code 5003. Limitation of motion		5018 [Removed]	
must be objectively confirmed by findings		5019 Bursitis. 5020 [Removed]	
such as swelling, muscle spasm, or satis-		5020 [Removed] 5021 Myositis.	
factory evidence of painful motion. In the		5021 Myosius. 5022 [Removed]	
absence of limitation of motion, rate as		5022 [Removed] 5023 Heterotopic ossification.	
below:		5024 Tenosynovitis, tendinitis, tendinosis or	

38 CFR Ch. I (7-1-21 Edition)

Rating Major Minor

§4.71a

ACUTE, SUBACUTE, OR CHRONIC DISEASES— Continued

	Rat- ing
Note to DCs 5013 through 5024: Evaluate the diseases under diagnostic codes 5013 through 5024 as degenerative arthritis, based on limitation of motion of affected parts. 5025 Fibromyalgia (fibrositis, primary fibromyalgia syndrome) With widespread musculoskeletal pain and tender points, with or without associated fatigue, sleep disturbance, stiffness, paresthesias, headache, irritable bowel symptoms, depression, anxiety, or Raynaud's-like symptoms:	
That are constant, or nearly so, and refractory to therapy	40
time That require continuous medication for control	20 10
NOTE: Widespread pain means pain in both the left and right sides of the body, that is both above and below the waist, and that affects both the axial skeleton (<i>i.e.</i> , cervical spine, anterior chest, thoracic spine, or low back) and the extremities.	10

PROSTHETIC IMPLANTS AND RESURFACING

	Rat	ing
	Major	Minor
Note (1): When an evaluation is assigned for joint resurfacing or the prosthetic replacement of a joint under diagnostic codes 5051–5056, an additional rating under §4.71a may not also be assigned for that joint, unless otherwise directed. Note (2): Only evaluate a revision procedure in the same manner as the original procedure under diagnostic codes 5051–5056 if all the original components are replaced. Note (3): The term 'prosthetic replacement' in diagnostic codes 5051–5053 and 5055–5056 means a total replacement of the named joint. However, in DC 5054, "prosthetic replacement" means a total replacement of the head of the femur or of the acetabulum. Note (4): The 100 percent rating for 1 year following implantation of prosthesis will commence after initial grant of the 1-month total rating assigned under §4.30 following hospital discharge. Note (5): The 100 percent rating for 4 months following implantation of prosthesis or resurfacing under DCs 5054 and 5055 will commence after initial grant of the 1-month total rating assigned under §4.30 following hospital discharge. Note (6): Special monthly compensation is assignable during the 100 percent rating period the earliest date permanent use of crutches is established.		

PROSTHETIC IMPLANTS AND RESURFACING— Continued

		Prosthetic replacement of the shoulder joint:
		For 1 year following implantation of
100	100	prosthesis
		With chronic residuals consisting of severe, painful motion or weak-
50	60	ness in the affected extremity
		With intermediate degrees of resid-
		ual weakness, pain or limitation of motion, rate by analogy to di-
		agnostic codes 5200 and 5203.
20	30	Minimum rating
		5052 Elbow replacement (prosthesis).
		Prosthetic replacement of the elbow
		joint: For 1 year following implantation of
100	100	prosthesis
		With chronic residuals consisting of
		severe painful motion or weak-
40	50	ness in the affected extremity With intermediate degrees of resid-
		ual weakness, pain or limitation
		of motion rate by analogy to di-
		agnostic codes 5205 through
20	30	5208. Minimum evaluation
20	50	5053 Wrist replacement (prosthesis).
		Prosthetic replacement of wrist joint:
400	400	For 1 year following implantation of
100	100	prosthesis
		severe, painful motion or weak-
30	40	ness in the affected extremity
		With intermediate degrees of resid-
		ual weakness, pain or limitation of motion, rate by analogy to di-
		agnostic code 5214.
20	20	Minimum rating
		5054 Hip, resurfacing or replacement
		(prosthesis): For 4 months following implantation
100		of prosthesis or resurfacing
		Prosthetic replacement of the head
		of the femur or of the acetab-
		ulum: Following implantation of
		prosthesis with painful
		motion or weakness
		such as to require the
1 90		use of crutches Markedly severe residual
		weakness, pain or limi-
		tation of motion fol-
		lowing implantation of
70		prosthesis
,,		Moderately severe residu- als of weakness, pain or
,		
		limitation of motion
50		Minimum evaluation, total
50		Minimum evaluation, total replacement only
50		Minimum evaluation, total replacement only Note: At the conclusion of the 100 percent
50		Minimum evaluation, total replacement only Note: At the conclusion of the 100 percent evaluation period, evaluate resurfacing
50		Minimum evaluation, total replacement only Note: At the conclusion of the 100 percent evaluation period, evaluate resurfacing under diagnostic codes 5250 through 5255; there is no minimum evaluation for
50		Minimum evaluation, total replacement only
50		Minimum evaluation, total replacement only
50		Minimum evaluation, total replacement only

40

20 10

Department of Veterans Affairs

§4.71a

PROSTHETIC IMPLANTS AND RESURFACING—Continued

PROSTHETIC IMPLANTS AND RESURFACING—Continued

	Rat	ing		Rati	ing
	Major	Minor		Major	Minor
Prosthetic replacement of knee joint: With chronic residuals consisting of severe painful motion or weakness in the affected extremity		60	With chronic residuals consisting of severe painful motion or weakness. With intermediate degrees of residual weakness, pain or limitation of motion rate by analogy to 5270 or 5271. Minimum rating		40
pain or limitation of motion rate by analogy to diagnostic codes 5256, 5261, or 5262. Minimum evaluation, total replacement only Note: At the conclusion of the 100 percent evaluation period, evaluate resurfacing under diagnostic codes 5256 through 5262; there is no minimum evaluation for resurfacing. 5056 Ankle replacement (prosthesis). Prosthetic replacement of ankle joint:		30	COMBINATIONS OF DISABILITIES 5104 Anatomical loss of one hand and loss of use of one foot 5105 Anatomical loss of one foot and loss of use of one hand 5106 Anatomical loss of both hands 5107 Anatomical loss of both feet 5108 Anatomical loss of one hand and one foot 5109 Loss of use of both hands 5110 Loss of use of both feet 5111 Loss of use of one hand and one foot		1100 1100 1100 1100 1100 1100 1100
For 1 year following implantation of prosthesis		100	¹ Also entitled to special monthly compensati	ion.	

TABLE II—RATINGS FOR MULTIPLE LOSSES OF EXTREMITIES WITH DICTATOR'S RATING CODE AND 38 CFR CITATION

			Impairment of	other extremity		
Impairment of one extremity	Anatomical loss or loss of use below elbow	Anatomical loss or loss of use below knee	Anatomical loss or loss of use above elbow (preventing use of prosthesis)	Anatomical loss or loss of use above knee (preventing use of prosthesis)	Anatomical loss near shoulder (preventing use of prosthesis)	Anatomical loss near hip (pre- venting use of prosthesis)
Anatomical loss or loss of use below elbow. Anatomical loss or loss of use below knee.	M Codes M-1 a, b, or c, 38 CFR 3.350 (c)(1)(i).	L Codes L-1 d, e, f, or g, 38 CFR 3.350(b). L Codes L-1 a, b, or c, 38	M½ Code M-5, 38 CFR 3.350 (f)(1)(x). L½ Code L-2 b, 38 CFR 3.350	L ¹ / ₂ Code L–2 c, 38 CFR 3.350 (f)(1)(vi). L ¹ / ₂ Code L–2 a, 38 CFR	N Code N-3, 38 CFR 3.350 (f)(1)(xi). M Code M-3 b, 38 CFR 3.350	M Code M-3 c, 38 CFR 3.350 (f)(1)(viii) M Code M-3 a, 38 CFR
Anatomical loss or loss of use above elbow (preventing use		CFR 3.350(b).	(f)(1)(iii). N Code N–1, 38 CFR 3.350 (d)(1).	3.350 (f)(1)(i). M Code M–2 a, 38 CFR 3.350 (c)(1)(iii).	(f)(1)(iv). N½ Code N-4, 38 CFR 3.350 (f)(1)(ix).	3.350 (f)(1)(ii) M½ Code M-4 c, 38 CFR 3.350 (f)(1)(xi)
of prosthesis). Anatomical loss or loss of use above knee (preventing use of prosthesis).				M Code M-2 a, 38 CFR 3.350 (c)(1)(ii).	M ¹ / ₂ Code M–4 b, 38 CFR 3.350 (f)(1)(vii).	M½ Code M-4 a, 38 CFR 3.350 (f)(1)(v)
Anatomical loss near shoulder (preventing use					O Code O-1, 38 CFR 3.350 (e)(1)(i).	N Code N-2 b, 38 CFR 3.350 (d)(3)
of prosthesis). Anatomical loss near hip (pre- venting use of prosthesis).						N Code N-2 a, 38 CFR 3.350 (d)(2)

Note.—Need for aid attendance or permanently bedridden qualifies for subpar. L. Code L–1 h, i (38 CFR 3.350(b)). Paraplegia with loss of use of both lower extremities and loss of anal and bladder sphincter control qualifies for subpar. O. Code O–2 (38 CFR 3.350(e)(2)). Where there are additional disabilities rated 50% or 100%, or anatomical or loss of use of a third extremity see 38 CFR 3.350(f) (3), (4) or (5).

38 CFR Ch. I (7-1-21 Edition)

§4.71a

(Authority: 38 U.S.C. 1115)

AMPUTATIONS: UPPER EXTREMITY

	Rati	ing	
	Major	Minor	(d
Arm, amputation of:			
5120 Complete amputation, upper extremity:			
Forequarter amputation (involving			
complete removal of the humerus along with any portion of the			
scapula, clavicle, and/or ribs)	¹ 100	¹ 100	(e
Disarticulation (involving complete removal of the humerus only)	190	1 90	
5121 Above insertion of deltoid	190	1 80	
5122 Below insertion of deltoid	180	170	
Forearm, amputation of:			
5123 Above insertion of pronator teres	¹ 80	170	
5124 Below insertion of pronator teres	170	¹ 60	
5125 Hand, loss of use of	170	¹ 60	
MULTIPLE FINGER AMPUTATIONS			(f)
5126 Five digits of one hand, amputation			(*)
of	170	1 60	
Four digits of one hand, amputation of:	'0		
5127 Thumb, index, long and ring	170	¹ 60	
5128 Thumb, index, long and little	170	1 60	
5129 Thumb, index, ring and little	170	¹ 60	
5130 Thumb, long, ring and little	170	¹ 60	
5131 Index, long, ring and little	60	50	5152
Three digits of one hand, amputation of:			W
5132 Thumb, index and long	60	50	A
5133 Thumb, index and ring	60	50	
5134 Thumb, index and little	60	50	A1
5135 Thumb, long and ring	60	50	5153
5136 Thumb, long and little	60	50	W
5137 Thumb, ring and little5138 Index, long and ring	60	50 40	W
5138 Index, long and ring5139 Index, long and little	50 50	40	VV
5140 Index, ring and little	50	40	
5141 Long, ring and little	40	30	TI
Two digits of one hand, amputation of:	10		5154
5142 Thumb and index	50	40	W
5143 Thumb and long	50	40	
5144 Thumb and ring	50	40	W
5145 Thumb and little	50	40	
5146 Index and long	40	30	E4EE
5147 Index and ring	40	30	5155 W
5148 Index and little	40	30	VV
5149 Long and ring	30	20	W
5150 Long and little	30	20	•
5151 Ring and little	30	20	
(a) The ratings for multiple finger ampu-			5156
tations apply to amputations at the proximal interphalangeal joints or			W
through proximal phalanges			
(b) Amputation through middle pha-			W
langes will be rated as prescribed for			
unfavorable ankylosis of the fingers			N.I
(c) Amputations at distal joints, or			N
through distal phalanges, other than			
negligible losses, will be rated as pre-			
scribed for favorable ankylosis of the fingers			¹ En

AMPUTATIONS: UPPER EXTREMITY—Continued

AMPUTATIONS: UPPER EXTREMITY—	-0011111	iueu
	Rati	ing
	Major	Minor
 (d) Amputation or resection of metacarpal bones (more than one-half the bone lost) in multiple fingers injuries will require a rating of 10 percent added to (not combined with) the ratings, multiple finger amputations, subject to the amputation rule applied to the forearm. (e) Combinations of finger amputations at various levels, or finger amputations with ankylosis or limitation of motion of the fingers will be rated on the basis of the grade of disability; i.e., amputation, unfavorable ankylosis, most representative of the levels or combinations. With an even number of fingers involved, and adjacent grades of disability, select the higher of the two grades. (f) Loss of use of the hand will be held to exist when no effective function remains other than that which would be equally well served by an amputation stump with a suitable prosthetic application. 		
ance. SINGLE FINGER AMPUTATIONS		
5152 Thumb, amputation of:		
With metacarpal resection At metacarpophalangeal joint or through	40	30
proximal phalanxAt distal joint or through distal phalanx 5153 Index finger, amputation of	30 20	20 20
With metacarpal resection (more than one-half the bone lost)	30	20
mal interphalangeal joint or proximal thereto	20 10	20 10
5154 Long finger, amputation of: With metacarpal resection (more than one-half the bone lost) Without metacarpal resection, at proxi-	20	20
mal interphalangeal joint or proximal thereto	10	10
With metacarpal resection (more than one-half the bone lost)	20	20
mal interphalangeal joint or proximal thereto	10	10
With metacarpal resection (more than one-half the bone lost)	20	20
mal interphalangeal joint or proximal thereto NOTE: The single finger amputation ratings are the only applicable ratings for amputations of whole or part of single fingers.	10	10

¹ Entitled to special monthly compensation.

SINGLE FINGER AMPUTATIONS




PLATE III

AMPUTATIONS: LOWER EXTREMITY

AMPUTATIONS: LOWER EXTREMITY—Continued

	Rat- ing
Thigh, amputation of:	

		Rat- ing
5160	Complete amputation, lower extremity:	

38 CFR Ch. I (7-1-21 Edition)

AMPUTATIONS: LOWER EXTREMITY—Continued

	Rat- ing
Trans-pelvic amputation (involving complete	
removal of the femur and intrinsic pelvic	
musculature along with any portion of the	2400
pelvic bones) Disarticulation (involving complete removal	2100
of the femur and intrinsic pelvic muscula-	
ture only)	290
Note: Separately evaluate residuals involving other	30
body systems (e.g., bowel impairment, bladder im-	
pairment) under the appropriate diagnostic code.	
5161 Upper third, one-third of the distance from	
perineum to knee joint measured from perineum	280
5162 Middle or lower thirds	² 60
Leg, amputation of:	
5163 With defective stump, thigh amputation rec-	
ommended	² 60
5164 Amputation not improvable by prosthesis con-	
trolled by natural knee action	² 60
5165 At a lower level, permitting prosthesis	2 40

AMPUTATIONS: LOWER EXTREMITY—Continued

	Rat- ing
5166 Forefoot, amputation proximal to metatarsal	
bones (more than one-half of metatarsal loss)	² 40
5167 Foot, loss of use of	² 40
5170 Toes, all, amputation of, without metatarsal	
loss or transmetatarsal, amputation of, with up to	
half of metatarsal loss	30
5171 Toe, great, amputation of:	
With removal of metatarsal head	30
Without metatarsal involvement	10
5172 Toes, other than great, amputation of, with re-	
moval of metatarsal head:	
One or two	20
Without metatarsal involvement	(
5173 Toes, three or four, amputation of, without	
metatarsal involvement:	
Including great toe	20
Not including great toe	10

² Also entitled to special monthly compensation.

AMPUTATIONS: LOWER EXTREMITY




PLATE IV

38 CFR Ch. I (7-1-21 Edition)

§4.71a

THE SHOULDER AND ARM

THE ELBOW AND FOREARM—Continued

	Rating	
	Major	Minor
5200 Scapulohumeral articulation, anky-		
losis of:		
NOTE: The scapula and humerus move as one piece.		
Unfavorable, abduction limited to 25°		
from side	50	40
Intermediate between favorable and un-		
favorable	40	30
Favorable, abduction to 60°, can reach		
mouth and head	30	20
5201 Arm, limitation of motion of: Flexion and/or abduction limited to		
25° from side	40	30
Midway between side and shoulder	10	00
level (flexion and/or abduction		
limited to 45°)	30	20
At shoulder level (flexion and/or ab-		
duction limited to 90°)	20	20
5202 Humerus, other impairment of: Loss of head of (flail shoulder)		70
Nonunion of (false flail joint)	80 60	70 50
Fibrous union of	50	40
Recurrent dislocation of at		
scapulohumeral joint:		
With frequent episodes		
and guarding of all arm		
movements	30	20
With infrequent episodes and guarding of move-		
ment only at shoulder		
level (flexion and/or ab-		
duction at 90°)	20	20
Malunion of:		
Marked deformity	30	20
Moderate deformity	20	20
5203 Clavicle or scapula, impairment of: Dislocation of	20	20
Nonunion of:	20	20
With loose movement	20	20
Without loose movement	10	10
Malunion of	10	10
Or rate on impairment of function of		
contiguous joint.		

THE ELBOW AND FOREARM

	Rating	
	Major	Minor
5205 Elbow, ankylosis of:		
Unfavorable, at an angle of less than		
50° or with complete loss of		
supination or pronation	60	50
Intermediate, at an angle of more than		
90°, or between 70° and 50°	50	40
Favorable, at an angle between 90° and		
70°	40	30
5206 Forearm, limitation of flexion of:		
Flexion limited to 45°	50	40
Flexion limited to 55°	40	30
Flexion limited to 70°	30	20
Flexion limited to 90°	20	20
Flexion limited to 100°	10	10
Flexion limited to 110°	0	0
5207 Forearm, limitation of extension of:		
Extension limited to 110°	50	40
Extension limited to 100°	40	30
Extension limited to 90°	30	20
Extension limited to 75°	20	20

	Rating	
	Major	Minor
Extension limited to 60° Extension limited to 45°	10 10	10 10
extension to 45°	20 60	20 50
dius5210 Radius and ulna, nonunion of, with	20	20
flail false joint	50	40
deformity	40	30
deformity	30	20
Nonunion in lower half	20	20
Malunion of, with bad alignment 5212 Radius, impairment of: Nonunion in lower half, with false movement: With loss of bone substance (1 inch	10	10
(2.5 cms.) or more) and marked deformity	40	30
deformity	30	20
Nonunion in upper half	20	20
Malunion of, with bad alignment 5213 Supination and pronation, impairment of:	10	10
Loss of (bone fusion):		
The hand fixed in supination or		
hyperpronation The hand fixed in full pronation	40 30	30 20
The hand fixed near the middle of the arc or moderate pronation	20	20
Limitation of pronation: Motion lost beyond middle of arc Motion lost beyond last quarter of arc, the hand does not approach	30	20
full pronationLimitation of supination:	20	20
To 30° or less	10	10

THE WRIST

	Rating	
	Major	Minor
5214 Wrist, ankylosis of: Unfavorable, in any degree of palmar flexion, or with ulnar or radial deviation	50 40 30	40 30 20

THE WRIST—Continued

	Rating	
	Major	Minor
Dorsiflexion less than 15°	10	10
Palmar flexion limited in line with fore- arm	10	10

EVALUATION OF ANKYLOSIS OR LIMITATION OF MOTION OF SINGLE OR MULTIPLE DIGITS OF THE HAND

	Rati	Rating	
	Major	Minor	
(1) For the index, long, ring, and little fingers (digits II, III, IV, and V), zero degrees of flexion represents the fingers fully extended, making a straight line with the rest of the hand. The position of function of the hand is with the wrist dorsiflexed 20 to 30 degrees, the metacarpophalangeal and proximal interphalangeal joints flexed to 30 degrees, and the thumb (digit I) abducted and rotated so that the thumb pad faces the finger pads. Only joints in these positions are considered to be in favorable position. For digits II through V, the metacarpophalangeal joint has a range of zero to 90 degrees of flexion, the proximal interphalangeal joint has a range of zero to 100 degrees of flexion, and the distal (terminal) interphalangeal joint has a range of zero to 70 or 80 degrees of flex-			
ion			
the next higher level			
thereto (ii) If both the metacarpophalangeal and proximal interphalangeal joints of a digit are ankylosed, evaluate as unfavorable ankylosis, even if each joint is individually fixed in a favorable position.			

EVALUATION OF ANKYLOSIS OR LIMITATION OF MOTION OF SINGLE OR MULTIPLE DIGITS OF THE HAND—Continued

	Rating	
	Major	Minor
(iii) If only the metacarpophalangeal or proximal interphalangeal joint is ankylosed, and there is a gap of more than two inches (5.1 cm.) between the fingertip(s) and the proximal transverse crease of the palm, with the finger(s) flexed to the extent possible, evaluate as unfavorable ankylosis		
proximal transverse crease of the palm, with the finger(s) flexed to the extent possible, evaluate as favorable ankylosis		
ankylosed, and either is in extension or full flexion, or there is rotation or angulation of a bone, evaluate as amputation at metacarpophalangeal joint or through proximal phalanx		
is individually fixed in a favorable position		
interphalangeal joint is ankylosed, and there is a gap of more than two inches (5.1 cm.) between the thumb pad and the fingers, with the thumb attempting to oppose the fingers, evaluate as unfavorable ankylosis (iv) If only the carpometacarpal or interphalangeal joint is ankylosed, and there is a gap of two inches (5.1 cm.) or less between the thumb pad and the fingers, with the thumb attempting to oppose the fingers, evaluate		
as favorable ankylosis(5) If there is limitation of motion of two or more digits, evaluate each digit separately		
and combine the evaluations		
I. Multiple Digits: Unfavorable Ank	ylosis	
5216 Five digits of one hand, unfavorable ankylosis of	60	50
ankylosis of: Thumb and any three fingers Index, long, ring, and little fingers	60 50	50 40

38 CFR Ch. I (7-1-21 Edition)

§4.71a

EVALUATION OF ANKYLOSIS OR LIMITATION OF MOTION OF SINGLE OR MULTIPLE DIGITS OF THE HAND—Continued

THE HAND—Continued		
	Rating	
	Major	Minor
Note: Also consider whether evaluation as		
amputation is warranted. 5218 Three digits of one hand, unfavorable		
ankylosis of:		
Thumb and any two fingers	50	40
Index, long, and ring; index, long,		
and little; or index, ring, and little	40	
fingers Long, ring, and little fingers	40 30	30 20
Note: Also consider whether evaluation as		
amputation is warranted.		
5219 Two digits of one hand, unfavorable		
ankylosis of:	40	20
Thumb and any finger	40	30
index and little fingers	30	20
Long and ring; long and little; or		
ring and little fingers	20	20
Note: Also consider whether evaluation as amputation is warranted.		
	deele	
II. Multiple Digits: Favorable Anky	yiosis	
5220 Five digits of one hand, favorable an-		40
kylosis of5221 Four digits of one hand, favorable	50	40
ankylosis of:		
Thumb and any three fingers	50	40
Index, long, ring, and little fingers	40	30
5222 Three digits of one hand, favorable an-		
kylosis of: Thumb and any two fingers	40	30
Index, long, and ring; index, long,		
and little; or index, ring, and little		
fingers	30	20
Long, ring and little fingers	20	20
5223 Two digits of one hand, favorable ankylosis of:		
Thumb and any finger	30	20
Index and long; index and ring; or		
index and little fingers	20	20
Long and ring; long and little; or	10	10
ring and little fingers	10	10
III. Ankylosis of Individual Dig	its	
5224 Thumb, ankylosis of:		
Unfavorable	20	20
Favorable Note: Also consider whether evaluation as	10	10
amputation is warranted and whether an		
additional evaluation is warranted for re-		
sulting limitation of motion of other digits		
or interference with overall function of the hand.		
5225 Index finger, ankylosis of:		
Unfavorable or favorable	10	10
Note: Also consider whether evaluation as		
amputation is warranted and whether an		
additional evaluation is warranted for re- sulting limitation of motion of other digits		
or interference with overall function of the		
hand.		

5226 Long finger, ankylosis of:
Unfavorable or favorable

EVALUATION OF ANKYLOSIS OR LIMITATION OF MOTION OF SINGLE OR MULTIPLE DIGITS OF THE HAND—Continued

		Rati	ing
		Major	Minor
Note: Also consider whether eval amputation is warranted and will additional evaluation is warrante sulting limitation of motion of ot or interference with overall functionand. 5227 Ring or little finger, ankylosis Unfavorable or favorable. Note: Also consider whether eval amputation is warranted and will additional evaluation is warrante sulting limitation of motion of ot or interference with overall functional.	hether an ed for re- ther digits ion of the s of:	0	(
IV. Limitation of Motion o	f Individua	l Digits	
5228 Thumb, limitation of motion: With a gap of more than to (5.1 cm.) between the th and the fingers, with th attempting to oppose the With a gap of one to tw (2.5 to 5.1 cm.) between thumb pad and the fing	humb pad he thumb e fingers vo inches ween the	20	20
the thumb attempting to the fingers	o oppose one inch	10	10
and the fingers, with the attempting to oppose the second	he thumb e fingers	0	(
crease of the palm, wit ger flexed to the extent	gertip and crease of r flexed to ; with exe than 30 one inch e fingertip ransverse th the finpossible,	10	10
and; extension is limite more than 30 degrees 5230 Ring or little finger, limitation		0	(
tion: Any limitation of motion		0	

10

10

Department of Veterans Affairs

§4.71a

Rating

	Rat- ing
General Rating Formula for Diseases and Injuries of the Spine	
(For diagnostic codes 5235 to 5243 unless 5243 is evaluated under the Formula for Rating Intervertebral Disc Syndrome Based on Incapacitating Episodes):	

THE SPINE

With or without symptoms such as pain (wither or not it radiates), stiffness, or aching in the area of the spine affected by residuals of injury or disease

Unfavorable ankylosis of the entire spine

100

50

40

30

20

Unfavorable ankylosis of the entire thoracolumbar spine

Unfavorable ankylosis of the entire cervical spine; or, forward flexion of the thoracolumbar spine 30 degrees or less; or, favorable ankylosis of the entire thoracolumbar spine

Forward flexion of the cervical spine 15 degrees or less; or, favorable ankylosis of the entire cervical spine

Forward flexion thoracolumbar spine greater than 30 degrees but not greater than 60 degrees; or, forward flexion of the cervical spine greater than 15 degrees but not greater than 30 degrees; or, the combined range of motion of the thoracolumbar spine not greater than 120 degrees; or, the combined range of motion of the cervical spine not greater than 170 degrees; or, muscle spasm or quarding severe enough to result in an abnormal gait or abnormal spinal contour such as scoliosis reversed lordosis, or abnormal kyphosis

Forward flexion thoracolumbar spine greater than 60 degrees but not greater than 85 degrees; or, forward flexion of the cervical spine greater than 30 degrees but not greater than 40 degrees; or, combined range of motion of the thoracolumbar spine greater than 120 degrees but not greater than 235 degrees; or, combined range of motion of the cervical spine greater than 170 degrees but not greater than 335 degrees: or. muscle spasm, guarding, or localized tenderness not resulting in abnormal gait or abnormal spi-nal contour; or, vertebral body fracture with loss of 50 percent or more of the height

Note (1): Evaluate any associated objective neurologic abnormalities, including, but not limited to, bowel or bladder impairment, separately, under an appropriate diagnostic code.

THE SPINE—Continued

Note (2): (See also Plate V.) For VA compensation purposes, normal forward flexion of the cervical spine is zero to 45 degrees, extension is zero to 45 degrees, extension is zero to 45 degrees, and left and right lateral rotation are zero to 80 degrees. Normal forward flexion of the thoracolumbar spine is zero to 90 degrees, extension is zero to 30 degrees, left and right lateral flexion are zero to 30 degrees, and left and right lateral rotation are zero to 30 degrees. The combined range of motion refers to the sum of the range of forward flexion, extension, left and right lateral flexion, and left and right rotation. The normal combined range of motion of the cervical spine is 340 degrees and of the thoracolumbar spine is 240 degrees. The normal ranges of motion for each component of spinal motion provided in this note are the maximum that can be used for calculation of the combined range of motion.

Note (3): In exceptional cases, an examiner may state that because of age, body habitus, neurologic disease, or other factors not the result of disease or injury of the spine, the range of motion of the spine in a particular individual should be considered normal for that individual, even though it does not conform to the normal range of motion stated in Note (2). Provided that the examiner supplies an explanation, the examiner's assessment that the range of motion is normal for that individual will be accepted.

Note (4): Round each range of motion measurement to the nearest five degrees.

Note (5): For VA compensation purposes, unfavorable ankylosis is a condition in which the entire cervical spine, the entire thoracolumbar spine, or the entire spine is fixed in flexion or extension, and the ankylosis results in one or more of the following: difficulty walking because of a limited line of vision; restricted opening of the mouth and chewing; breathing limited to diaphragmatic respiration; gastrointestinal symptoms due to pressure of the costal margin on the abdomen; dyspnea or dysphagia; atlantoaxial or cervical subluxation or dislocation; or neurologic symptoms due to nerve root stretching. Fixation of a spinal segment in neutral position (zero degrees) always represents favorable ankylosis.

Note (6): Separately evaluate disability of the thoracolumbar and cervical spine segments, except when there is unfavorable ankylosis of both segments, which will be rated as a single disability.

5235 Vertebral fracture or dislocation

5236 Sacroiliac injury and weakness

5237 Lumbosacral or cervical strain

5238 Spinal stenosis

5239 Spondylolisthesis or segmental instability

5240 Ankylosing spondylitis

5241 Spinal fusion

5242 Degenerative arthritis, degenerative disc disease other than intervertebral disc syndrome (also, see either DC 5003 or 5010)

10 5243 Intervertebral disc syndrome: Assign this diagnostic code only when there is disc herniation with compression and/or irritation of the adjacent nerve root; assign diagnostic code 5242 for all other disc diagnoses.


431

38 CFR Ch. I (7-1-21 Edition)

THE SPINE—Continued

THE SPINE—Continued

	Rat- ing		Rat- ing
Evaluate intervertebral disc syndrome (preoperatively or postoperatively) either under the General Rating Formula for Diseases and Injuries of the Spine or under the Formula for Rating Intervertebral Disc Syndrome Based on Incapacitating Episodes, whichever method results in the higher evaluation when all disabilities are combined under § 4.25. 5244 Traumatic paralysis, complete: Paraplegia: Rate under diagnostic code 5110. Quadriplegia: Rate separately under diagnostic codes 5109 and 5110 and combine evaluations in accordance with § 4.25. Note: If traumatic paralysis does not cause loss of use of both hands or both feet, it is incomplete paralysis. Evaluate residuals of incomplete traumatic paralysis under the appropriate diagnostic code (e.g., § 4.124a, Diseases of the Peripheral Nerves). Formula for Rating Intervertebral Disc Syndrome Based on Incapacitating Episodes With incapacitating episodes having a total duration of at least 6 weeks during the past 12 months	60	With incapacitating episodes having a total duration of at least 4 weeks but less than 6 weeks during the past 12 months With incapacitating episodes having a total duration of at least 2 weeks but less than 4 weeks during the past 12 months With incapacitating episodes having a total duration of at least one week but less than 2 weeks during the past 12 months Note (1): For purposes of evaluations under diagnostic code 5243, an incapacitating episode is a period of acute signs and symptoms due to intervertebral disc syndrome that requires bed rest prescribed by a physician and treatment by a physician. Note (2): If intervertebral disc syndrome is present in more than one spinal segment, provided that the effects in each spinal segment are clearly distinct, evaluate each segment on the basis of incapacitating episodes or under the General Rating Formula for Diseases and Injuries of the Spine, whichever method results in a higher evaluation for that segment.	40 20 10






PLATE V
RANGE OF MOTION OF CERVICAL AND THORACOLUMBAR SPINE

38 CFR Ch. I (7-1-21 Edition)

THE			

	Rat- ing
5250 Hip, ankylosis of:	
Unfavorable, extremely unfavorable ankylosis,	
the foot not reaching ground, crutches neces-	
sitated	³ 90
Intermediate	70
Favorable, in flexion at an angle between 20°	
and 40°, and slight adduction or abduction	60
5251 Thigh, limitation of extension of:	
Extension limited to 5°	10
5252 Thigh, limitation of flexion of:	
Flexion limited to 10°	40
Flexion limited to 20°	30
Flexion limited to 30°	20
Flexion limited to 45°	10
5253 Thigh, impairment of:	
Limitation of abduction of, motion lost beyond	
10°	20
Limitation of adduction of, cannot cross legs	10
Limitation of rotation of, cannot toe-out more	
than 15°, affected leg	10
5254 Hip, flail joint	80
5255 Femur, impairment of:	
Fracture of shaft or anatomical neck of: With nonunion, with loose motion	
	00
(spiral or oblique fracture)	80
With nonunion, without loose mo- tion, weight bearing preserved	
with aid of brace	60
Fracture of surgical neck of, with	60
false joint	60
Malunion of:	00
Evaluate under diagnostic codes	
5256, 5257, 5260, or 5261 for	
the knee, or 5250–5254 for the	
hip, whichever results in the	
highest evaluation.	

 $^{^{\}rm 3}\,\mbox{Entitled}$ to special monthly compensation.

THE KNEE AND LEG

	Rat- ing	(
		No
5256 Knee, ankylosis of:		i
Extremely unfavorable, in flexion at an angle of		
45° or more	60	t
In flexion between 20° and 45°	50	i
In flexion between 10° and 20°	40	(
Favorable angle in full extension, or in slight		t
flexion between 0° and 10°	30	t
5257 Knee, other impairment of:		5258 Carti
Recurrent subluxation or instability:		episodes
Unrepaired or failed repair of com-		joint
plete ligament tear causing per-		5259 Carti
sistent instability, and a medical		5260 Leg,
provider prescribes both an as-		Flexion
sistive device (e.g., cane(s),		Flexion
crutch(es), walker) and bracing for ambulation	30	Flexion Flexion
One of the following:	30	5261 Leg,
(a) Sprain, incomplete lig-		Extensi
ament tear, or repaired		Extensi
complete ligament tear		Extensi
causing persistent insta-		Extensi
bility, and a medical		Extensi
provider prescribes a		Extensi
brace and/or assistive		5262 Tibia
device (e.g., cane(s),		No.
crutch(es), walker) for		
ambulation.		Ma

THE KNEE AND LEG—Continued

	Rat- ing
(b) Unrepaired or failed re-	
pair of complete liga-	
ment tear causing per-	
sistent instability, and a	
medical provider pre-	
scribes either an assist-	
ive device (e.g.,	
cane(s), crutch(es),	
walker) or bracing for	
ambulation	20
Sprain, incomplete ligament tear, or	
complete ligament tear (repaired,	
unrepaired, or failed repair) caus- ing persistent instability, without	
a prescription from a medical	
provider for an assistive device	
(e.g., cane(s), crutch(es), walker)	
or bracing for ambulation	10
Patellar instability:	
A diagnosed condition involving the	
patellofemoral complex with re-	
current instability after surgical	
repair that requires a prescription	
by a medical provider for a brace	
and either a cane or a walker	30
A diagnosed condition involving the	
patellofemoral complex with re-	
current instability after surgical	
repair that requires a prescription	
by a medical provider for one of the following: A brace, cane, or	
walker	20
A diagnosed condition involving the	20
patellofemoral complex with re-	
current instability (with or without	
history of surgical repair) that	
does not require a prescription	
from a medical provider for a	
brace, cane, or walker	10
Note (1): For patellar instability, the	
patellofemoral complex consists of the	
quadriceps tendon, the patella, and the	
patellar tendon. Note (2): A surgical procedure that does not	
involve repair of one or more	
patellofemoral components that contribute	
to the underlying instability shall not qual-	
ify as surgical repair for patellar instability	
(including, but not limited to, arthroscopy	
to remove loose bodies and joint aspira-	
tion).	
8 Cartilage, semilunar, dislocated, with frequent	
pisodes of "locking," pain, and effusion into the	
int	20
9 Cartilage, semilunar, removal of, symptomatic 0 Leg, limitation of flexion of:	10
0 Leg, limitation of flexion of: Flexion limited to 15°	30
Flexion limited to 30°	20
Flexion limited to 45°	10
Flexion limited to 60°	0
1 Leg, limitation of extension of:	
Extension limited to 45°	50
Extension limited to 30°	40
Extension limited to 20°	30
Extension limited to 15°	20
Extension limited to 10°	10
Extension limited to 5°	0
2 Tibia and fibula, impairment of:	
Nonunion of, with loose motion, requiring	
brace	40
Malunion of:	

Department of Veterans Affairs

§4.71a

THE KNEE AND LEG—Continued

	Rat- ing
Evaluate under diagnostic codes 5256, 5257, 5260, or 5261 for the knee, or 5270 or 5271 for the ankle, whichever results in the highest evaluation. Medial tibial stress syndrome (MTSS), or shin splints:	
Requiring treatment for no less than 12 consecutive months, and unresponsive to surgery and ei- ther shoe orthotics or other con-	
servative treatment, both lower extremities	30
servative treatment, one lower extremity	20
treatment less than 12 consecutive months, one or both lower ex-	10
tremities	C
tively demonstrated)	10

THE ANKLE

	Rat- ing
5270 Ankle, ankylosis of:	
In plantar flexion at more than 40°, or in	
dorsiflexion at more than 10° or with abduc-	
tion, adduction, inversion or eversion deformity	40
In plantar flexion, between 30° and 40°, or in	
dorsiflexion, between 0° and 10°	30
In plantar flexion, less than 30°	20
5271 Ankle, limited motion of:	
Marked (less than 5 degrees dorsiflexion or	
less than 10 degrees plantar flexion)	20
Moderate (less than 15 degrees dorsiflexion	
or less than 30 degrees plantar flexion)	10
5272 Subastragalar or tarsal joint, ankylosis of:	
In poor weight-bearing position	20
In good weight-bearing position	10
5273 Os calcis or astragalus, malunion of:	
Marked deformity	20
Moderate deformity	10
5274 Astragalectomy	20

SHORTENING OF THE LOWER EXTREMITY

	Rat- ing
5275 Bones, of the lower extremity, shortening of: Over 4 inches (10.2 cms.)	³ 60 ³ 50 40 30 20 10

SHORTENING OF THE LOWER EXTREMITY—Continued

	Rat- ing
NOTE: Measure both lower extremities from ante- rior superior spine of the ilium to the internal malleolus of the tibia. Not to be combined with other ratings for fracture or faulty union in the same extremity.	

³ Also entitled to special monthly compensation.

THE FOOT

	Rat- ing
5269 Plantar fasciitis: No relief from both non-surgical and surgical	
treatment, bilateral	30
treatment, unilateral	20
Otherwise, unilateral or bilateral	10
Note (1): With actual loss of use of the foot, rate 40 percent.	
Note (2): If a veteran has been	
recommended for surgical inter-	
vention, but is not a surgical can-	
didate, evaluate under the 20 percent or 30 percent criteria,	
whichever is applicable.	
5276 Flatfoot, acquired:	
Pronounced; marked pronation, extreme tender-	
ness of plantar surfaces of the feet, marked	
inward displacement and severe spasm of the tendo achillis on manipulation, not improved	
by orthopedic shoes or appliances.	
Bilateral	50
Unilateral	30
Severe; objective evidence of marked deformity	
(pronation, abduction, etc.), pain on manipula- tion and use accentuated, indication of swell-	
ing on use, characteristic callosities:	
Bilateral	30
Unilateral	20
Moderate; weight-bearing line over or medial to great toe, inward bowing of the tendo achillis,	
pain on manipulation and use of the feet, bilat-	
eral or unilateral	10
Mild; symptoms relieved by built-up shoe or arch	
support	0
5277 Weak foot, bilateral: A symptomatic condition secondary to many	
constitutional conditions, characterized by at-	
rophy of the musculature, disturbed circulation,	
and weakness:	
Rate the underlying condition, minimum rat-	10
ing5278 Claw foot (pes cavus), acquired:	10
Marked contraction of plantar fascia with	
dropped forefoot, all toes hammer toes, very	
painful callosities, marked varus deformity:	
BilateralUnilateral	50 30
All toes tending to dorsiflexion, limitation of	30
dorsiflexion at ankle to right angle, shortened	
plantar fascia, and marked tenderness under	
metatarsal heads:	
Bilateral Unilateral	30 20
Great toe dorsiflexed, some limitation of	20
dorsiflexion at ankle, definite tenderness under	
metatarsal heads:	
Bilateral	10

38 CFR Ch. I (7-1-21 Edition)

THE FOOT—Continued

	Rat- ing
Unilateral	10
Slight	0
5279 Metatarsalgia, anterior (Morton's disease), unilateral, or bilateral	10
5280 Hallux valgus, unilateral:	
Operated with resection of metatarsal head	10
Severe, if equivalent to amputation of great toe	10
5281 Hallux rigidus, unilateral, severe:	
Rate as hallux valgus, severe.	
Note: Not to be combined with claw foot	
ratings. 5282 Hammer toe:	
All toes, unilateral without claw foot	10
Single toes	0
5283 Tarsal, or metatarsal bones, malunion of, or	U
nonunion of:	
Severe	30
Moderately severe	20
Moderate	10
NOTE: With actual loss of use of the foot, rate 40	
percent.	
5284 Foot injuries, other:	
Severe	30
Moderately severe	20
Moderate	10
NOTE: With actual loss of use of the foot, rate 40 percent.	

THE SKULL

	Rat- ing
5296 Skull, loss of part of, both inner and outer ta- bles:	
With brain hernia Without brain hernia:	80
Area larger than size of a 50-cent piece or 1.140 in ² (7.355 cm ²)	50
Area intermediate	30
piece or 0.716 in 2 (4.619 cm ²)	10
plications.	

THE RIBS

	Rat- ing
5297 Ribs, removal of:	
More than six	50
Five or six	40
Three or four	30
Two	20
One or resection of two or more ribs without re-	
generation	10
NOTE (1): The rating for rib resection or removal	
is not to be applied with ratings for purrulent	
pleurisy, lobectomy, pneumonectomy or inju-	
ries of pleural cavity.	
NOTE (2): However, rib resection will be consid-	
ered as rib removal in thoracoplasty performed	
for collapse therapy or to accomplish oblitera-	
tion of space and will be combined with the	
rating for lung collapse, or with the rating for	
lobectomy, pneumonectomy or the graduated	
ratings for pulmonary tuberculosis.	

THE COCCYX

	Rat- ing
5298 Coccyx, removal of: Partial or complete, with painful residuals Without painful residuals	10

(Authority: 38 U.S.C. 1155)

[29 FR 6718, May 22, 1964, as amended at 34 FR 5062, Mar. 11, 1969; 40 FR 42536, Sept. 15, 1975; 41 FR 11294, Mar. 18, 1976; 43 FR 45350, Oct. 2, 1978; 51 FR 6411, Feb. 24, 1986; 61 FR 20439, May 7, 1996; 67 FR 48785, July 26, 2002; 67 FR 54349, Aug. 22, 2002; 68 FR 51456, Aug. 27, 2003; 69 FR 32450, June 10, 2004; 80 FR 42041, July 16, 2015; 85 FR 76460, Nov. 30, 2020, 85 FR 85523, Dec. 29, 2020, 86 FR 8142, Feb. 4, 2021]

§4.72 [Reserved]

§ 4.73 Schedule of ratings—muscle injuries.

NOTE (1): When evaluating any claim involving muscle injuries resulting in loss of use of any extremity or loss of use of both buttocks (diagnostic code 5317, Muscle Group XVII), refer to §3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation.

NOTE (2): Ratings of slight, moderate, moderately severe, or severe for diagnostic codes 5301 through 5323 will be determined based upon the criteria contained in §4.56.

THE SHOULDER GIRDLE AND ARM

	Rating	
	Domi- nant	Non- domi- nant
5301 Group I. Function: Upward rotation of scapula; elevation of arm above shoulder level. Extrinsic muscles of shoulder girdle: (1) Trapezius; (2) levator scapulae; (3) serratus magnus. Severe Moderately Severe Moderate Slight 5302 Group II. Function: Depression of arm from vertical overhead to hanging at side (1, 2); downward rotation of scapula (3, 4); 1 and 2 act with Group III in forward and backward swing of arm. Extrinsic muscles of shoulder girdle: (1) Pectoralis major II (costosternal); (2) latissimus dorsi and teres major (teres major, although technically an intrinsic muscle, is included with latissimus dorsi);	40 30 10 0	30 20 10 0
(3) pectoralis minor; (4) rhomboid. Severe	40	30
Moderately Severe	30	20
Moderate	20	20
Slight	0	0

THE SHOULDER GIRDLE AND ARM—Continued

	Rating	
	Domi- nant	Non- domi- nant
5303 Group III. Function: Elevation and abduction of arm to level of shoulder; act with 1 and 2 of Group II in forward and backward swing of arm. Intrinsic muscles of shoulder girdle: (1) Pectoralis major I (clavicular); (2) deltoid. Severe	40	30
Moderately Severe	30 20 0	20 20 0
shoulder against injury in strong move- ments, holding head of humerus in socket; abduction; outward rotation and inward ro- tation of arm. Intrinsic muscles of shoulder girdle: (1) Supraspinatus; (2) infraspinatus and teres minor; (3) subscapularis; (4) coracobrachialis.		
Severe	30	20
Moderately Severe Moderate	20 10	20 10
Slight	0	0
5305 Group V. Function: Elbow supination (1) (long head of biceps is stabilizer of shoulder joint); flexion of elbow (1, 2, 3). Flexor muscles of elbow: (1) Biceps; (2) brachialis; (3) brachioradialis.	0	U
Severe	40	30
Moderately Severe	30	20
Moderate	10	10
Slight	0	0
Severe	40	30
Moderately Severe	30	20
Moderate	10	10
Slight	0	0

THE FOREARM AND HAND

	Rating	
	Domi- nant	Non- domi- nant
5907 Group VII. Function: Flexion of wrist and fingers. Muscles arising from internal condyle of humerus: Flexors of the carpus and long flexors of fingers and thumb; pronator. Severe	40 30	30 20
Moderately Severe Moderate	10	10
Slight	0	10
5308 Group VIII. Function: Extension of wrist, fingers, and thumb; abduction of thumb. Muscles arising mainly from external condyle of humerus: Extensors of carpus, fingers, and thumb; supinator.	· ·	0
Severe	30	20
Moderately Severe	20	20
Moderate	10	10
Slight	0	0

THE FOREARM AND HAND—Continued

	Rating	
	Domi- nant	Non- domi- nant
5309 Group IX. Function: The forearm muscles act in strong grasping movements and are supplemented by the intrinsic muscles in delicate manipulative movements. Intrinsic muscles of hand: Thenar eminence; short flexor, opponens, abductor and adductor of thumb; hypothenar eminence; short flexor, opponens and abductor of little finger; 4 lumbricales; 4 dorsal and 3 palmar interossei. NOTE: The hand is so compact a structure that isolated muscle injuries are rare, being nearly always complicated with injuries of bones, joints, tendons, etc. Rate on limitation of motion, minimum 10 percent.		
THE FOOT AND LEG		

	9
5310 Group X. Function: Movements of forefoot and toes; propulsion thrust in walking. Intrinsic muscles of the foot: Plantar: (1) Flexor digitorum brevis; (2) abductor hallucis; (3) abductor digiti minimi; (4) quadratus plantae; (5) lumbricales; (6) flexor hallucis brevis; (7) adductor hallucis; (8) flexor digiti minimi brevis; (9) dorsal and plantar interossei. Other important plantar structures: Plantar aponeurosis, long plantar and calcaneonavicular ligament, tendons of posterior tibial, peroneus longus, and long flexors of great and little toes.	
Severe Moderately Severe Moderately Severe Moderate Slight Sight Dorsal: (1) Extensor hallucis brevis; (2) extensor digitorum brevis. Other important dorsal structures: cruciate, crural, deltoid, and other ligaments; tendons of long extensors of toes and peronei mus-	30 20 10 0
cles. Severe	20 10 10 0
wounds of the foot—10. 5311 Group XI. Function: Propulsion, plantar flexion of foot (1); stabilization of arch (2, 3); flexion of toes (4, 5); Flexion of knee (6). Posterior and lateral crural muscles, and muscles of the calf: (1) Triceps surae (gastrocnemius and soleus); (2) tibialis posterior; (3) peroneus longus; (4) peroneus brevis; (5) flexor hallucis longus; (6) flexor digitorum longus; (7) popliteus; (8) plantaris. Severe	30 20 10
Slight Slight Function: Dorsiflexion (1); extension of toes (2); stabilization of arch (3). Anterior muscles of the leg: (1) Tibialis anterior; (2) extensor digitorum longus; (3) extensor hallucis longus; (4) peroneus tertius. Severe	30

38 CFR Ch. I (7-1-21 Edition)

THE FOOT AND LEG—Continued

	Rat- ing
Moderately Severe	20 10 0

THE PELVIC GIRDLE AND THIGH

THE PELVIC GIRDLE AND THIGH	
	Ra in
fexion of knee; outward and inward rotation of flexed knee; acting with rectus femoris and sartorius (see XIV, 1, 2) synchronizing simultaneous flexion of hip and knee and extension of hip and knee by belt-over-pulley action at knee joint. Posterior thigh group, Hamstring complex of 2-joint muscles: (1) Biceps femoris; (2) semimembranosus; (3) semitendinosus. Severe Moderately Severe Moderately Severe Moderately Severe Slight 5314 Group XIV. Function: Extension of knee (2, 3, 4, 5); simultaneous flexion of hip and flexion of knee (1); tension of fascia lata and iliotibial (Maissiat's) band, acting with XVII (1) in postural support of body (6); acting with hamstrings in synchronizing hip and knee (1, 2). Anterior thigh group: (1) Sartorius; (2) rectus femoris; (3) vastus externus; (4) vastus intermedius; (5) vastus intermus; (6) tensor vaginae femoris. Severe Moderately Severe	
Moderate Slight ————————————————————————————————————	
5316 Group XVI. Function: Flexion of hip (1, 2, 3). Pelvic girdle group 1: (1) Psoas; (2) iliacus; (3) pectineus. Severe Moderately Severe Moderate Slight	
5317 Group XVII. Function: Extension of hip (1); abduction of thigh; elevation of opposite side of pelvis (2, 3); tension of fascia lata and iliotibial (Maissiat's) band, acting with XIV (6) in postural support of body steadying pelvis upon head of femur and condyles of femur on tibia (1). Pelvic girdle group 2: (1) Gluteus maximus; (2) gluteus medius; (3) gluteus minimus. Severe Moderately Severe Moderate	*
Slight	

THE PELVIC GIRDLE AND THIGH—Continued

	Rat- ing
Slight	C
* If hilatoral cap & 2 350(a)(2) of this chapter to do	tormino

 * If bilateral, see §3.350(a)(3) of this chapter to determine whether the veteran may be entitled to special monthly compensation.

THE TORSO AND NECK

5319	Group XIX. Function: Support and compres-
sion	of abdominal wall and lower thorax; flexion
	lateral motions of spine; synergists in strong
	nward movements of arm (1). Muscles of the
	ominal wall: (1) Rectus abdominis; (2) external
oblid	que; (3) internal oblique; (4) transversalis; (5)
qua	dratus lumborum.
	Severe
	Moderately Severe
	Moderate
	Slight
5320	Group XX. Function: Postural support of body;
	ension and lateral movements of spine. Spinal
	scles: Sacrospinalis (erector spinae and its pro-
	pations in thoracic and cervical regions).
	Cervical and thoracic region:.
	Severe
	Moderately Severe
	Moderate
	Slight
	Lumbar region:.
	Severe
	Moderately Severe
	Moderate
	Slight
5321	Group XXI. Function: Respiration. Muscles of
resp	piration: Thoracic muscle group.
	Severe or Moderately Severe
	Moderate
	Slight
5322	Group XXII. Function: Rotary and forward
	rements of the head; respiration; deglutition.
	scles of the front of the neck: (Lateral, supra-,
	infrahyoid group.) (1) Trapezius I (clavicular in-
	ion); (2) sternocleidomastoid; (3) the "hyoid"
mus	scles; (4) sternothyroid; (5) digastric.
	Severe
	Moderately Severe
	Moderate
	Slight
5323	Group XXIII. Function: Movements of the
hea	d; fixation of shoulder movements. Muscles of
the	side and back of the neck: Suboccipital; lateral
vert	ebral and anterior vertebral muscles.
	Severe
	Moderately Severe
	Moderate
Climba	

	Rat- ing
5324 Diaphragm, rupture of, with herniation. Rate under diagnostic code 7346.	

MISCELLANEOUS—Continued

Rat-
ina

- 5325 Muscle injury, facial muscles. Evaluate functional impairment as seventh (facial) cranial nerve neuropathy (diagnostic code 8207), disfiguring scar (diagnostic code 7800), etc. Minimum, if interfering to any extent with mastication-10.
- 5326 Muscle hernia, extensive. Without other injury to the muscle-10.
- Muscle, neoplasm of, malignant (excluding soft tissue sarcoma)-100.
- NOTE: A rating of 100 percent shall continue beyond the cessation of any surgery, radiation treatment, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residual impairment of function.
- 5328 Muscle, neoplasm of, benign, postoperative Rate on impairment of function, *i.e.*, limitation of motion, or scars, diagnostic code 7805, etc.
- 5329 Sarcoma, soft tissue (of muscle, fat, or fibrous connective tissue)-100.
- NOTE: A rating of 100 percent shall continue beyond the cessation of any surgery, radiation treatment. antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis. rate on residual impairment of function.
- - Rhabdomyolysis, residuals of:
 Rate each affected muscle group separately and combine in accordance with § 4.25. Note: Separately evaluate any chronic renal complications within the appropriate body system.
- 5331 Compartment syndrome:
 - Rate each affected muscle group separately and combine in accordance with § 4.25.

(Authority: 38 U.S.C. 1155)

[62 FR 30239, June 3, 1997, as amemded 85 FR 76464, Nov. 30, 20201

THE ORGANS OF SPECIAL SENSE

§4.75 General considerations for evaluating visual impairment.

- (a) Visual impairment. The evaluation of visual impairment is based on impairment of visual acuity (excluding developmental errors of refraction), visual field, and muscle function.
- (b) Examination for visual impairment. The examination must be conducted by a licensed optometrist or by a licensed ophthalmologist. The examiner must identify the disease, injury, or other

- pathologic process responsible for any visual impairment found. Examinations of visual fields or muscle function will be conducted only when there is a medical indication of disease or injury that may be associated with visual field defect or impaired muscle function. Unless medically contraindicated, the fundus must be examined with the claimant's pupils dilated.
- (c) Service-connected visual impairment of only one eye. Subject to the provisions of 38 CFR 3.383(a), if visual impairment of only one eye is serviceconnected, the visual acuity of the other eye will be considered to be 20/40 for purposes of evaluating the serviceconnected visual impairment.
- (d) Maximum evaluation for visual impairment of one eye. The evaluation for visual impairment of one eye must not exceed 30 percent unless there is anatomical loss of the eye. Combine the evaluation for visual impairment of one eye with evaluations for other disabilities of the same eye that are not based on visual impairment (e.g., disfigurement under diagnostic code 7800).
- (e) Anatomical loss of one eye with inability to wear a prosthesis. When the claimant has anatomical loss of one eye and is unable to wear a prosthesis, increase the evaluation for visual acuity under diagnostic code 6063 by 10 percent, but the maximum evaluation for visual impairment of both eyes must not exceed 100 percent. A 10-percent increase under this paragraph precludes an evaluation under diagnostic code 7800 based on gross distortion or asymmetry of the eye but not an evaluation under diagnostic code 7800 based on other characteristics of disfigurement.
- (f) Special monthly compensation. When evaluating visual impairment, refer to 38 CFR 3.350 to determine whether the claimant may be entitled to special monthly compensation. Footnotes in the schedule indicate levels of visual impairment that potentially establish entitlement to special monthly compensation; however, other levels of visual impairment combined with disabilities of other body systems may also establish entitlement.

(Authority: 38 U.S.C. 1114 and 1155)

[73 FR 66549, Nov. 10, 2008]

§4.76 Visual acuity.

- (a) Examination of visual acuity. Examination of visual acuity must include the central uncorrected and corrected visual acuity for distance and near vision using Snellen's test type or its equivalent.
- (b) Evaluation of visual acuity. (1) Evaluate central visual acuity on the basis of corrected distance vision with central fixation, even if a central scotoma is present. However, when the lens required to correct distance vision in the poorer eye differs by more than three diopters from the lens required to correct distance vision in the better eye (and the difference is not due to congenital or developmental refractive error), and either the poorer eye or both eves are service connected, evaluate the visual acuity of the poorer eye using either its uncorrected or corrected visual acuity, whichever results in better combined visual acuity.
- (2) Provided that he or she customarily wears contact lenses, evaluate the visual acuity of any individual affected by a corneal disorder that results in severe irregular astigmatism that can be improved more by contact lenses than by eyeglass lenses, as corrected by contact lenses.
- (3) In any case where the examiner reports that there is a difference equal to two or more scheduled steps between

near and distance corrected vision, with the near vision being worse, the examination report must include at least two recordings of near and distance corrected vision and an explanation of the reason for the difference. In these cases, evaluate based on corrected distance vision adjusted to one step poorer than measured.


(4) To evaluate the impairment of visual acuity where a claimant has a reported visual acuity that is between two sequentially listed visual acuities, use the visual acuity which permits the higher evaluation.

(Authority: 38 U.S.C. 1155) [73 FR 66549, Nov. 10, 2008]

§ 4.76a Computation of average concentric contraction of visual fields.

TABLE III—NORMAL VISUAL FIELD EXTENT AT 8
PRINCIPAL MERIDIANS

Meridian	Normal de- grees
Temporally	85
Down temporally	85
Down	65
Down nasally	50
Nasally	60
Up nasally	55
Up	45
Up temporally	55
Total	500



TS-19 52a

Example of computation of concentric contraction under the schedule with abnormal findings taken from Figure 1.

Loss	Degrees
Temporally	55
Down temporally	55
Down	45

Loss	Degrees
Down nasally Nasally Up nasally Up temporally	30 40 35 25 35
Total loss	320

Remaining field 500° minus 320° = 180°. 180° ÷ 8 = 22½° average concentric contraction.

(Authority: 38 U.S.C. 1155)

 $[43\ \mathrm{FR}\ 45352,\ \mathrm{Oct.}\ 2,\ 1978,\ \mathrm{as}\ \mathrm{amended}\ \mathrm{at}\ 73\ \mathrm{FR}\ 66549,\ \mathrm{Nov.}\ 10,\ 2008]$

§ 4.77 Visual fields.

(a) Examination of visual fields. Examiners must use either Goldmann kinetic perimetry or automated perimetry using Humphrey Model 750, Octopus Model 101, or later versions of these perimetric devices with simulated kinetic Goldmann testing capability. For phakic (normal) individuals, as well as for pseudophakic or aphakic individuals who are well adapted to intraocular lens implant or contact lens correction, visual field examinations must be conducted using a standard target size and luminance, Goldmann's equivalent which III/4e. For aphakic individuals not well adapted to contact correction lens Or pseudophakic individuals not well adapted to intraocular lens implant. visual field examinations must be conducted using Goldmann's equivalent

IV/4e. The examiner must document the results for at least 16 meridians 22½ degrees apart for each eye and indicate the Goldmann equivalent used. See Table III for the normal extent (in degrees) of the visual fields at the 8 principal meridians (45 degrees apart). When the examiner indicates that additional testing is necessary to evaluate visual fields, the additional testing must be conducted using either a tangent screen or a 30-degree threshold visual field with the Goldmann III stimulus size. The examination report must document the results of either the tangent screen or of the 30-degree threshold visual field with the Goldmann III stimulus size.

- (b) Evaluation of visual fields. Determine the average concentric contraction of the visual field of each eye by measuring the remaining visual field (in degrees) at each of eight principal meridians 45 degrees apart, adding them, and dividing the sum by eight.
- (c) Combination of visual field defect and decreased visual acuity. To determine the evaluation for visual impairment when both decreased visual acuity and visual field defect are present in one or both eyes and are service connected, separately evaluate the visual acuity and visual field defect (expressed as a level of visual acuity), and combine them under the provisions of § 4.25.




Figure 2. Goldmann Perimeter Chart

52c

(Authority: 38 U.S.C. 1155)

 $[53 \ \mathrm{FR} \ 30262, \ \mathrm{Aug.} \ 11, \ 1988, \ \mathrm{as} \ \mathrm{amended} \ \mathrm{at} \ 73 \ \mathrm{FR} \ 66549, \ \mathrm{Nov.} \ 10, \ 2008; \ 74 \ \mathrm{FR} \ 7648, \ \mathrm{Feb.} \ 19, \ 2009; \ 83 \ \mathrm{FR} \ 15320, \ \mathrm{Apr.} \ 10, \ 2018]$

§4.78 Muscle function.

(a) Examination of muscle function. The examiner must use a Goldmann perimeter chart or the Tangent Screen method that identifies the four major quadrants (upward, downward, left, and right lateral) and the central field (20 degrees or less) (see Figure 2). The examiner must document the results of muscle function testing by identifying the quadrant(s) and range(s) of degrees in which diplopia exists.

(b) Evaluation of muscle function. (1) An evaluation for diplopia will be assigned to only one eye. When a claimant has both diplopia and decreased

visual acuity or visual field defect, assign a level of corrected visual acuity for the poorer eye (or the affected eye, if disability of only one eye is serviceconnected) that is: one step poorer than it would otherwise warrant if the evaluation for diplopia under diagnostic code 6090 is 20/70 or 20/100; two steps poorer if the evaluation under diagnostic code 6090 is 20/200 or 15/200; or three steps poorer if the evaluation under diagnostic code 6090 is 5/200. This adjusted level of corrected visual acuity, however, must not exceed a level of 5/200. Use the adjusted visual acuity for the poorer eye (or the affected eye, if

38 CFR Ch. I (7-1-21 Edition)

§4.79

disability of only one eye is service-connected), and the corrected visual acuity for the better eye (or visual acuity of 20/40 for the other eye, if only one eye is service-connected) to determine the percentage evaluation for visual impairment under diagnostic codes 6065 through 6066.

(2) When diplopia extends beyond more than one quadrant or range of degrees, evaluate diplopia based on the quadrant and degree range that provides the highest evaluation.

(3) When diplopia exists in two separate areas of the same eye, increase the equivalent visual acuity under diagnostic code 6090 to the next poorer level of visual acuity, not to exceed 5/200

(Authority: 38 U.S.C. 1155)

[73 FR 66550, Nov. 10, 2008, as amended at 83 FR 15321, Apr. 10, 2018]

§4.79 Schedule of ratings—eye.

DISEASES OF THE EYE

al Rating Formula for Diseases of the Eye: Evaluate on the basis of either visual impairment due to the particular condition or on incapacitating episodes, whichever results in a higher evaluation	
With documented incapacitating episodes requiring 7 or more treatment visits for an eye condition during the past 12 months With documented incapacitating episodes requiring at least 5 but less than 7 treatment visits for an eye condition during the past 12 months With documented incapacitating episodes requiring at least 3 but less than 5 treatment visits for an eye condition during the past 12 months With documented incapacitating episodes requiring at least 3 but less than 5 treatment visits for an eye condition during the past 12 months Note (1): For the purposes of evaluation under 38 CFR 4.79, an incapacitating episode is an eye condition severe enough to require a clinic visit to a provider specifically for treatment purposes. Note (2): Examples of treatment may include but are not limited to: Systemic immunosuppressants or biologic agents; intravitreal or periocular injections; laser treatments; or other surgical interventions. Note (3): For the purposes of evaluating visual impairment due to the particular condition, refer to 38 CFR 4.75-4.78 and to § 4.79, diagnostic codes 6061–6091.	60 40 20 10
Choroidopathy, including uveitis, iritis, cyclitis, or choroiditis. Keratopathy.	
Unhealed eye injury.	
Note: This code includes orbital trauma, as well as penetrating or non-penetrating eye injury	
Tuberculosis of eye:	
	100
Localized scars, atrophy, or irregularities of the retina, unilateral or bilateral, that are centrally located and that result in an irregular, duplicated, enlarged, or diminished image	10
medication is required	10
medication is required	10
Malignant neoplasms of the eye, orbit, and adnexa (excluding skin): Malignant neoplasms of the eye, orbit, and adnexa (excluding skin) that require therapy that is comparable to those used for systemic malignancies, i.e., systemic chemotherapy, X-ray therapy more extensive than the the case of the control of the contro	100
to the area of the eye, or surgery more extensive than enucleation Note: Continue the 100 percent rating beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy, or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating will be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination will be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, evaluate based on residuals Malignant neoplasms of the eye, orbit, and adnexa (excluding skin) that do not require therapy comparable to that for systemic malignancies: Separately evaluate visual and nonvisual impairment, e.g., disfigurement (diagnostic code 7800), and combine the evaluations.	100
	with documented incapacitating episodes requiring at least 1 but less than 3 treatment visits for an eye condition during the past 12 months

DISEASES OF THE EYE—Continued

	DISEASES OF THE LTE—Continued	
		Rating
	Separately evaluate visual and nonvisual impairment, e.g., disfigurement (diagnostic code 7800), and combine the evaluations	
	Nystagmus, central	10
	Trachomatous conjunctivitis: Active: Evaluate under the General Rating Formula for Diseases of the Eye, minimum rating	30
	Active: Evaluate under the General Rating Formula for Diseases of the Eye, minimum rating	10
6019	Ptosis, unilateral or bilateral: Evaluate based on visual impairment or, in the absence of visual impairment, on disfigurement (diagnostic code 7800).	
6020	Ectropion: Bilateral	20 10
6021	Unilateral Entropion: Bilateral	20
6022	Unilateral Lagophthalmos:	10
	Bilateral Unilateral	20 10
6023	Loss of eyebrows, complete, unilateral or bilateral	10
	Loss of eyelashes, complete, unilateral or bilateral	10
6025	Disorders of the lacrimal apparatus (epiphora, dacryocystitis, etc.): Bilateral	20 10
6026	Optic neuropathy	
6027	Cataract: Preoperative: Evaluate under the General Rating Formula for Diseases of the Eye Postoperative: If a replacement lens is present (pseudophakia), evaluate under the General Rating Formula for Diseases of the Eye. If there is no replacement lens, evaluate based on aphakia (diagnostic code 6029)	
6029	Aphakia or dislocation of crystalline lens: Evaluate based on visual impairment, and elevate the resulting level of visual impairment one step. Minimum (unilateral or bilateral)	30
	Paralysis of accommodation (due to neuropathy of the Oculomotor Nerve (cranial nerve III)). Loss of eyelids, partial or complete:	20
6034	Separately evaluate both visual impairment due to eyelid loss and nonvisual impairment, e.g., disfigurement (diagnostic code 7800), and combine the evaluations. Pteryglum:	
	Evaluate under the General Rating Formula for Diseases of the Eye, disfigurement (diagnostic code 7800), conjunctivitis (diagnostic code 6018), etc., depending on the particular findings, and combine in accordance with § 4.25	
	Keratoconus Status post corneal transplant: Evaluate under the General Rating Formula for Diseases of the Eye. Minimum, if there is pain,	
6037	photophobia, and glare sensitivity	10
	Diabetic retinopathy	
	Retinal dystrophy (including retinitis pigmentosa, wet or dry macular degeneration, early-onset macular de- eration, rod and/or cone dystrophy)	
	Post-chiasmal disorders	
	Impairment of Central Visual Acuity	
6061	Anatomical loss of both eyes ¹	100
6062	Anatomical loss of one eye: 1 Anatomical loss of one eye: 1	100
	In the other eye 5/200 (1.5/60)	100
	In the other eye 10/200 (3/60)	90 80
	In the other eye 13/200 (4:3/00)	70
	In the other eye 20/100 (6/30)	60
	In the other eye 20/70 (6/21)	60
	In the other eye 20/50 (6/15)	50 40
6064	No more than light perception in one eye: 1	10
	In the other eye 5/200 (1.5/60)	100
	In the other eye 10/200 (3/60)	90 80
	,,	50

DISEASES OF THE EYE—Continued

	Rating
In the other eye 20/200 (6/60)	70
In the other eye 20/100 (6/30)	60
In the other eye 20/70 (6/21)	50
In the other eye 20/50 (6/15)	40
In the other eye 20/40 (6/12)	
6065 Vision in one eye 5/200 (1.5/60):	
In the other eye 5/200 (1.5/60)	1100
In the other eye 10/200 (3/60)	
In the other eye 15/200 (4.5/60)	
In the other eye 20/200 (6/60)	
In the other eye 20/100 (6/30)	
In the other eye 20/70 (6/21)	
In the other eye 20/50 (6/15)	
In the other eye 20/40 (6/12)	
6066 Visual acuity in one eye 10/200 (3/60) or better:	
Vision in one eye 10/200 (3/60):	
In the other eye 10/200 (3/60)	90
	1
In the other eye 15/200 (4.5/60)	
In the other eye 20/200 (6/60)	
In the other eye 20/100 (6/30)	
In the other eye 20/70 (6/21)	
In the other eye 20/50 (6/15)	
In the other eye 20/40 (6/12)	30
Vision in one eye 15/200 (4.5/60):	
In the other eye 15/200 (4.5/60)	80
In the other eye 20/200 (6/60)	70
In the other eye 20/100 (6/30)	60
In the other eye 20/70 (6/21)	40
In the other eye 20/50 (6/15)	30
In the other eye 20/40 (6/12)	20
Vision in one eye 20/200 (6/60):	
In the other eye 20/200 (6/60)	70
In the other eye 20/100 (6/30)	60
In the other eye 20/70 (6/21)	
In the other eye 20/50 (6/15)	
In the other eye 20/40 (6/12)	
Vision in one eye 20/100 (6/30):	
In the other eye 20/100 (6/30)	50
In the other eye 20/70 (6/21)	
In the other eye 20/50 (6/15)	
In the other eye 20/40 (6/12)	
Vision in one eye 20/70 (6/21):	"
	30
In the other eye 20/70 (6/21)	
In the other eye 20/40 (6/12)	1 "
Vision in one eye 20/50 (6/15):	
In the other eye 20/50 (6/15)	
In the other eye 20/40 (6/12)	10
Vision in one eye 20/40 (6/12):	
In the other eye 20/40 (6/12)	(

¹ Review for entitlement to special monthly compensation under 38 CFR 3.350.

RATINGS FOR IMPAIRMENT OF VISUAL FIELDS

	Rating
6080 Visual field defects:	
Homonymous hemianopsia	. 30
Loss of temporal half of visual field:	
Bilateral	. 30
Unilateral	. 10
Or evaluate each affected eye as 20/70 (6/21).	
Loss of nasal half of visual field:	
Bilateral	. 10
Unilateral	. 10
Or evaluate each affected eye as 20/50 (6/15).	
Loss of inferior half of visual field:	
Bilateral	. 30
Unilateral	. 10
Or evaluate each affected eye as 20/70 (6/21).	
Loss of superior half of visual field:	

RATINGS FOR IMPAIRMENT OF VISUAL FIELDS—Continued

		Rating
	Bilateral	10
	Or evaluate each affected eye as 20/50 (6/15).	10
Conce	entric contraction of visual field:	
001100	With remaining field of 5 degrees: 1	
	Bilateral	100
	Unilateral	30
	Or evaluate each affected eye as 5/200 (1.5/60).	
With r	emaining field of 6 to 15 degrees:	
	Bilateral	70
	Unilateral	20
	Or evaluate each affected eye as 20/200 (6/60).	
With r	emaining field of 16 to 30 degrees:	
	Bilateral	50
	Unilateral	10
	Or evaluate each affected eye as 20/100 (6/30).	
With r	emaining field of 31 to 45 degrees:	
	Bilateral	30
	Unilateral	10
	Or evaluate each affected eye as 20/70 (6/21).	
With r	emaining field of 46 to 60 degrees:	
	Bilateral	10
	Unilateral	10
	Or evaluate each affected eye as 20/50 (6/15).	
6081	Scotoma, unilateral:	
	Minimum, with scotoma affecting at least one-quarter of the visual field (quadrantanopsia) or with centrally located scotoma of any size	10
	Alternatively, evaluate based on visual impairment due to scotoma, if that would result in a higher evaluation.	

 $^{^{\}rm 1}\,\mbox{Review}$ for entitlement to special monthly compensation under 38 CFR 3.350.

RATINGS FOR IMPAIRMENT OF MUSCLE FUNCTION

Degree of diplopia	Equivalent visual acuity
6090 Diplopia (double vision):	
(a) Central 20 degrees	5/200 (1.5/60)
(b) 21 degrees to 30 degrees	
(1) Down	15/200 (4.5/60)
(2) Lateral	20/100 (6/30)
(3) Up	20/70 (6/21)
(c) 31 degrees to 40 degrees	
(1) Down	20/200 (6/60)
(2) Lateral	20/70 (6/21)
(3) Up	20/40 (6/12)
Note: In accordance with 38 CFR 4.31, diplopia that is occasional or that is correctable with spectacles is	
evaluated at 0 percent.	
6091 Symblepharon:	
Evaluate under the General Rating Formula for Diseases of the Eye, lagophthalmos (diagnostic code 6022), disfigurement (diagnostic code 7800), etc., depending on the particular findings, and combine in accordance with § 4.25	

(Authority: 38 U.S.C. 1155)

 $[73\;\mathrm{FR}\;66550,\,\mathrm{Nov}.\;10,\,2008,\,\mathrm{as}\;\mathrm{amended}\;\mathrm{at}\;83\;\mathrm{FR}\;15321,\,\mathrm{Apr}.\;10,\,2018]$

$\S\S4.80-4.84$ [Reserved]

IMPAIRMENT OF AUDITORY ACUITY

\$4.85 Evaluation of hearing impairment.

(a) An examination for hearing impairment for VA purposes must be conducted by a state-licensed audiologist

and must include a controlled speech discrimination test (Maryland CNC) and a puretone audiometry test. Examinations will be conducted without the use of hearing aids.

(b) Table VI, "Numeric Designation of Hearing Impairment Based on Puretone Threshold Average and

§ 4.85

Speech Discrimination," is used to determine a Roman numeral designation (I through XI) for hearing impairment based on a combination of the percent of speech discrimination (horizontal rows) and the puretone threshold average (vertical columns). The Roman numeral designation is located at the point where the percentage of speech discrimination and puretone threshold average intersect.

- (c) Table VIa, "Numeric Designation of Hearing Impairment Based Only on Puretone Threshold Average," is used to determine a Roman numeral designation (I through XI) for hearing impairment based only on the puretone threshold average. Table VIa will be used when the examiner certifies that use of the speech discrimination test is not appropriate because of language difficulties, inconsistent speech discrimination scores, etc., or when indicated under the provisions of §4.86.
- (d) "Puretone threshold average," as used in Tables VI and VIa, is the sum of the puretone thresholds at 1000, 2000, 3000 and 4000 Hertz, divided by four. This average is used in all cases (including those in §4.86) to determine the

Roman numeral designation for hearing impairment from Table VI or VIa.

- (e) Table VII, "Percentage Evaluations for Hearing Impairment," is used to determine the percentage evaluation by combining the Roman numeral designations for hearing impairment of each ear. The horizontal rows represent the ear having the better hearing and the vertical columns the ear having the poorer hearing. The percentage evaluation is located at the point where the row and column intersect.
- (f) If impaired hearing is service-connected in only one ear, in order to determine the percentage evaluation from Table VII, the non-service-connected ear will be assigned a Roman Numeral designation for hearing impairment of I, subject to the provisions of § 3.383 of this chapter.
- (g) When evaluating any claim for impaired hearing, refer to §3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation due either to deafness, or to deafness in combination with other specified disabilities.
 - (h) Numeric tables VI, VIA*, and VII.

TABLE VI

NUMERIC DESIGNATION OF HEARING IMPAIRMENT BASED ON PURETONE THRESHOLD AVERAGE AND SPEECH DISCRIMINATION

Puretone Threshold Average

% of discrim- ination	0-41	42-49	50-57	58-65	66-73	74-81	82-89	90-97	98+
92-100	I	I	I	II	II	II	III	III	IV
84-90	II	II	II	III	III	III	IV	IV	IV
76-82	III	III	IV	IV	IV	V	V	V	V
68-74	IV	IV	V	V	VI	VI	VII	VII	VII
60-66	V	V	VI	VI	VII	VII	VIII	VIII	VIII
52-58	VI	VI	VII	VII	VIII	VIII	VIII	VIII	IX
44-50	VII	VII	VIII	VIII	VIII	IX	IX	IX	X
36-42	VIII	VIII	VIII	IX	IX	IX	X	X	X
0-34	IX	X	XI	XI	XI	XI	XI	XI	XI

TABLE VIA*

NUMERIC DESIGNATION OF HEARING IMPAIRMENT BASED ONLY ON PURETONE THRESHOLD AVERAGE

Puretone Threshold Average

0-41	42-48	49-55	56-62	63-69	70-76	77-83	84-90	91-97	98-104	105+
I	II	III	IV	V	VI	VII	VIII	IX	X	XI

^{*} This table is for use only as specified in §§ 4.85 and 4.86.

TABLE VII

PERCENTAGE EVALUATION FOR HEARING IMPAIRMENT (DIAGNOSTIC CODE 6100)

Poorer Ear

	XI	100*										
	X	90	80									
	IX	80	70	60								
	VIII	70	60	50	50							
<u> </u>	VII	60	60	50	40	40						
Better Ear	VI	50	50	40	40	30	30					
Be	V	40	40	40	30	30	20	20				
	IV	30	30	30	20	20	20	10	10			
	III	20	20	20	20	20	10	10	10	0		
	II	10	10	10	10	10	10	10	0 -	0	0	
	I	10	10	0	0	0	0	0	0	. 0	0	0
		XI	X	IX	VIII	VII	VI	v	IV	III	II	I

^{*} Review for entitlement to special monthly compensation under §3.350 of this chapter.

 $[64~{\rm FR}~25206,~{\rm May}~11,~1999]$

§ 4.86 Exceptional patterns of hearing impairment.

(a) When the puretone threshold at each of the four specified frequencies (1000, 2000, 3000, and 4000 Hertz) is 55 decibels or more, the rating specialist will determine the Roman numeral designation for hearing impairment from either Table VI or Table VIa, whichever results in the higher nu-

meral. Each ear will be evaluated separately.

(b) When the puretone threshold is 30 decibels or less at 1000 Hertz, and 70 decibels or more at 2000 Hertz, the rating specialist will determine the Roman numeral designation for hearing impairment from either Table VI or Table VIa, whichever results in the higher numeral. That numeral will then be elevated to the next higher

§4.88a

Roman numeral. Each ear will be evaluated separately.

(Authority: 38 U.S.C. 1155) [64 FR 25209, May 11, 1999]

§ 4.87 Schedule of ratings—ear.

DISEASES OF THE EAR

Rating

10

30

100

60 30

10

6200 Chronic suppurative otitis media, mastoiditis, or cholesteatoma (or any combination): During suppuration, or with aural polyps NOTE: Evaluate hearing impairment, and complications such as labyrinthitis, tinnitus, facial nerve paralysis, or bone loss of skull, separately.
6201 Chronic nonsuppurative otitis media with effusion (serous otitis media): Rate hearing impairment 6202 Otosclerosis:
Rate hearing impairment 6204 Peripheral vestibular disorders: Dizziness and occasional staggering Occasional dizziness NoTE: Objective findings supporting the diagnosis of vestibular disequilibrium are required before a compensable evaluation can be assigned under this code. Hearing impairment or suppuration shall be separately rated and
combined. 6205 Meniere's syndrome (endolymphatic hydrops): Hearing impairment with attacks of vertigo and cerebellar gait occurring more than once weekly, with or without tinnitus
Complete loss of both Complete loss of one Deformity of one, with loss of one-third or more of the substance
6208 Malignant neoplasm of the ear (other than skin only)

Rate on impairment of function.

6210 Chronic otitis externa:

DISEASES OF THE EAR—Continued

	Rat- ing
Swelling, dry and scaly or serous discharge, and itching requiring frequent and prolonged treatment Serous State	10 0 10

(Authority: 38 U.S.C. 1155)

 $[64\ FR\ 25210,\ May\ 11,\ 1999,\ as\ amended\ at\ 68\ FR\ 25823,\ May\ 14,\ 2003]$

§4.87a Schedule ratings-other sense organs.

	Rat- ing
6275 Sense of smell, complete loss	10 10

(Authority: 38 U.S.C. 1155)

[64 FR 25210, May 11, 1999]

INFECTIOUS DISEASES, IMMUNE DIS-ORDERS AND NUTRITIONAL Defi-CIENCIES

§4.88 [Reserved]

§ 4.88a Chronic fatigue syndrome.

(a) For VA purposes, the diagnosis of chronic fatigue syndrome requires:

- (1) new onset of debilitating fatigue severe enough to reduce daily activity to less than 50 percent of the usual level for at least six months; and
- (2) the exclusion, by history, physical examination, and laboratory tests, of all other clinical conditions that may produce similar symptoms; and
 - (3) six or more of the following: (i) acute onset of the condition,

 - (ii) low grade fever,
 - (iii) nonexudative pharyngitis,

451

38 CFR Ch. I (7-1-21 Edition)

§4.88b

- (iv) palpable or tender cervical or axillary lymph nodes,
- (v) generalized muscle aches or weakness,
- (vi) fatigue lasting 24 hours or longer after exercise,

(vii) headaches (of a type, severity, or pattern that is different from headaches in the pre-morbid state),

- (viii) migratory joint pains,
- (ix) neuropsychologic symptoms,
- (x) sleep disturbance.
- (b) [Reserved]

[59 FR 60902, Nov. 29, 1994]

§4.88b Schedule of ratings—infectious diseases, immune disorders and nutritional deficiencies.

NOTE: Rate any residual disability of infection within the appropriate body system as indicated by the notes in the evaluation criteria. As applicable, consider the long-term health effects potentially associated with infectious diseases as listed in §3.317(d) of this

Evaluate under the General Rating Formula.

chapter, specifically Brucellosis, Campylobacter jejuni, Coxiella burnetii (Q fever), Malaria, Mycobacterium Tuberculosis, Nontyphoid Salmonella, Shigella, Visceral Leishmaniasis, and West Nile virus.

Rating

		Rating
Gene	ral Rating Formula for Infectious Diseases: For active disease	100
	After active disease has resolved, rate at 0 percent for infection. Rate any residual disability of infection within the appropriate body system.	
6300	Vibriosis (Cholera, Non-cholera):	
	Evaluate under the General Rating Formula.	
	Note: Rate residuals of cholera and non-cholera vibrio infections, such as renal failure, skin, and musculo- skeletal conditions, within the appropriate body system.	
6301	Visceral leishmaniasis:	
	As active disease	100
	Note 1: Continue a 100 percent evaluation beyond the cessation of treatment for active disease. Six months after discontinuance of such treatment, determine the appropriate disability rating by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. Thereafter, rate under the appropriate body system any residual disability of infection, which includes, but is not limited to liver damage and bone marrow disease.	
	Note 2: Confirm the recurrence of active infection by culture, histopathology, or other diagnostic laboratory testing.	
6302	Leprosy (Hansen's disease):	
	As active disease	100
	Note: Continue a 100 percent evaluation beyond the cessation of treatment for active disease. Six months after discontinuance of such treatment, determine the appropriate disability rating by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. Thereafter, rate under the appropriate body system any residual disability of infection, which includes, but is not limited to, skin lesions, peripheral neuropathy, or amputations.	
6304	Malaria:	
	Evaluate under the General Rating Formula. Note 1: The diagnosis of malaria, both initially and during relapse, depends on the identification of the malarial parasites in blood smears or other specific diagnostic laboratory tests such as antigen detection,	
	immunologic (immunochromatographic) tests, and molecular testing such as polymerase chain reaction tests.	
	Note 2: Rate under the appropriate body system any residual disability of infection, which includes, but is not limited to, liver or splenic damage, and central nervous system conditions.	
6305	Lymphatic filariasis, to include elephantiasis:	
0000	Evaluate under the General Rating Formula.	
	Note: Rate under the appropriate body system any residual disability of infection, which includes, but is not limited to, epididymitis, lymphangitis, lymphatic obstruction, or lymphedema affecting extremities, genitals, and/or breasts.	
6306	Bartonellosis:	
	Evaluate under the General Rating Formula.	
	Note: Rate under the appropriate body system any residual disability of infection, which includes, but is not limited to, endocarditis or skin lesions.	
6307	Plague:	
	Evaluate under the General Rating Formula. Note: Rate under the appropriate body system any residual disability of infection.	
6308	Relansing Fever:	1

	Note: Rate under the appropriate hody system any residual disability of infaction, which includes, but is not	Rati
	Note: Rate under the appropriate body system any residual disability of infection, which includes, but is not limited to, liver or spleen damage, iritis, uveitis, or central nervous system involvement.	
309	Rheumatic fever:	
	Evaluate under the General Rating Formula.	
	Note: Rate under the appropriate body system any residual disability of infection, which includes, but is not limited to, heart damage.	
310	Syphilis, and other treponema infections:	
	Note: Rate under the appropriate body system any residual disability of infection, which includes, but is not	
	limited to, diseases of the nervous system, vascular system, eyes, or ears (see DC 7004, DC 8013, DC	
2011	8014, DC 8015, and DC 9301).	
311	Tuberculosis, miliary: As active disease	
	Inactive disease: See §§ 4.88c and 4.89.	
	Note 1: Confirm the recurrence of active infection by culture, histopathology, or other diagnostic laboratory	
	testing.	
	Note 2: Rate under the appropriate body system any residual disability of infection which includes, but is not limited to, skin conditions and conditions of the respiratory, central nervous, musculoskeletal, ocular, gastro-	
	intestinal, and genitourinary systems and those residuals listed in §4.88c.	
312	Nontuberculosis mycobacterium infection:	
	As active disease	
	Note 1: Continue the rating of 100 percent for the duration of treatment for active disease followed by a man-	
	datory VA exam. If there is no relapse, rate on residuals. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.	
	Note 2: Confirm the recurrence of active infection by culture, histopathology, or other diagnostic laboratory	
	testing.	
	Note 3: Rate under the appropriate body system any residual disability of infection which includes, but is not limited to, skin conditions and conditions of the respiratory, central nervous, musculoskeletal, ocular, gastro-	
	inteed to, skin conditions and conditions of the respiratory, central nervous, musculoskeletal, ocular, gastro- intestinal, and genitourinary systems and those residuals listed in §4.88c.	
313	Avitaminosis:	
	Marked mental changes, moist dermatitis, inability to retain adequate nourishment, exhaustion, and cachexia	
	With all of the symptoms listed below, plus mental symptoms and impaired bodily vigor	
	With stomatitis, diarrhea, and symmetrical dermatitis	
	Confirmed diagnosis with nonspecific symptoms such as: decreased appetite, weight loss, abdominal discom-	
	fort, weakness, inability to concentrate and irritability	
314	Beriberi: As active disease:	
	With congestive heart failure, anasarca, or Wernicke-Korsakoff syndrome	
	With cardiomegaly, or; with peripheral neuropathy with footdrop or atrophy of thigh or calf muscles	
	With peripheral neuropathy with absent knee or ankle jerks and loss of sensation, or; with symptoms such as	
	weakness, fatigue, anorexia, dizziness, heaviness and stiffness of legs, headache or sleep disturbance	
315	Thereafter rate residuals under the appropriate body system. Pellagra:	
	Marked mental changes, moist dermatitis, inability to retain adequate nourishment, exhaustion, and cachexia	
	With all of the symptoms listed below, plus mental symptoms and impaired bodily vigor	
	With stomatitis, diarrhea, and symmetrical dermatitis	
	With stomatitis, or achlorhydria, or diarrhea	
	fort, weakness, inability to concentrate and irritability	
316	Brucellosis:	
	Evaluate under the General Rating Formula.	
	Note 1: Culture, serologic testing, or both must confirm the initial diagnosis and recurrence of active infection. Note 2: Rate under the appropriate body system any residual disability of infection, which includes, but is not	
	limited to, meningitis, liver, spleen and musculoskeletal conditions.	
317	Rickettsial, enrlichia, and anaplasma infections:	
	Evaluate under the General Rating Formula.	
	Note 1: Rate under the appropriate body system any residual disability of infection, which includes, but is not limited to, bone marrow, spleen, central nervous system, and skin conditions.	
	Note 2: This diagnostic code includes, but is not limited to, scrub typhus, Rickettsial pox, African tick-borne	
	fever, Rocky Mountain spotted fever, ehrlichiosis, or anaplasmosis.	
318	Melioidosis:	
	Evaluate under the General Rating Formula.	
	Note 1: Confirm by culture or other specific diagnostic laboratory tests the initial diagnosis and any relapse or	
	chronic activity of infection. Note 2: Rate under the appropriate body system any residual disability of infection, which includes, but is not	
	limited to, arthritis, lung lesions, or meningitis.	
319	Lyme disease:	
	Evaluate under the General Rating Formula.	
	Note: Rate under the appropriate body system any residual disability of infection, which includes, but is not limited to, arthritis, Bell's palsy, radiculopathy, ocular, or cognitive dysfunction.	
320	Parasitic diseases otherwise not specified:	
	Evaluate under the General Rating Formula.	
	Note: Rate under the appropriate body system any residual disability of infection.	
325	Hyperinfection syndrome or disseminated strongyloidiasis:	

		Rating
	As active disease Note: Continue the rating of 100 percent through active disease followed by a mandatory VA exam. If there is no relapse, rate on residual disability. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.	10
326	Schistosomiasis:	
	As acute or asymptomatic chronic disease	
329	ous system. Hemorrhagic fevers, including dengue, yellow fever, and others:	
	Evaluate under the General Rating Formula. Note: Rate under the appropriate body system any residual disability of infection, which includes, but is not	
30	limited to, conditions of the central nervous system, liver, or kidney. Campylobacter jejuni infection:	
	Evaluate under the General Rating Formula.	
121	Note: Rate under the appropriate body system any residual disability of infection, which includes, but is not limited to, Guillain-Barre syndrome, reactive arthritis, or uveitis. Coxiella burnetii infection (Q fever):	
100	Evaluate under the General Rating Formula.	
	Note: Rate under the appropriate body system any residual disability of infection, which includes, but is not limited to, chronic hepatitis, endocarditis, osteomyelitis, post Q-fever chronic fatigue syndrome, or vascular infections.	
33	Nontyphoid salmonella infections:	
	Evaluate under the General Rating Formula.	
	Note: Rate under the appropriate body system any residual disability of infection, which includes, but is not limited to, reactive arthritis.	
34	Shigella infections:	
	Evaluate under the General Rating Formula. Note: Rate under the appropriate body system any residual disability of infection, which includes, but is not	
	limited to, hemolytic-uremic syndrome or reactive arthritis.	
35	West Nile virus infection: Evaluate under the General Rating Formula.	
	Note: Rate under the appropriate body system any residual disability of infection, which includes, but is not	
50	limited to, variable physical, functional, or cognitive disabilities. Lupus erythematosus, systemic (disseminated):	
50	Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impair-	
	ment of health	1
	Exacerbations once or twice a year or symptomatic during the past 2 years	
	NOTE: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.	
151	HIV-related illness: AIDS with recurrent opportunistic infections (see Note 3) or with secondary diseases afflicting multiple body	
	systems; HIV-related illness with debility and progressive weight loss	1
	Refractory constitutional symptoms, diarrhea, and pathological weight loss; or minimum rating following devel-	
	opment of AIDS-related opportunistic infection or neoplasm	
	ing with T4 cell count less than 200	
	Following development of HIV-related constitutional symptoms; T4 cell count between 200 and 500; use of approved medication(s); or with evidence of depression or memory loss with employment limitations	
	Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4	
	cell count	
	regimens and medications prescribed as part of a research protocol at an accredited medical institution.	
	Note 2: Diagnosed psychiatric illness, central nervous system manifestations, opportunistic infections, and	
	neoplasms may be rated separately under the appropriate diagnostic codes if a higher overall evaluation results, provided the disability symptoms do not overlap with evaluations otherwise assignable above.	
	Note 3: The following list of opportunistic infections are considered AIDS-defining conditions, that is, a diag-	
	nosis of AIDS follows if a person has HIV and one more of these infections, regardless of the CD4 count—candidiasis of the bronchi, trachea, esophagus, or lungs; invasive cervical cancer; coccidioidomycosis;	
	cryptococcosis; cryptosporidiosis; cytomegalovirus (particularly CMV retinitis); HIV-related encephalopathy;	
	herpes simplex-chronic ulcers for greater than one month, or bronchitis, pneumonia, or esophagitis;	
	histoplasmosis; isosporiasis (chronic intestinal); Kaposi's sarcoma; lymphoma; mycobacterium avium com- plex; tuberculosis; pneumocystis jirovecii (carinii) pneumonia; pneumonia, recurrent; progressive multifocal	
	leukoencephalopathy; salmonella septicemia, recurrent; toxoplasmosis of the brain; and wasting syndrome	
54	due to HIV. Chronic fatique syndrome (CFS):	
J +	Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, or confusion), or a	
	combination of other signs and symptoms:	
	Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care	1
	Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness	·
	level; or which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year	
	non del vegi	

	Rating
Which are nearly constant and restrict routine daily activities from 50 to 75 percent of the pre-illness level; or which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year	40 20 10

 $[61~{\rm FR}~39875,\,{\rm July}~31,\,1996,\,{\rm as~amended~at}~84~{\rm FR}~28230,\,{\rm June}~18,\,2019]$

§4.88c Ratings for inactive nonpulmonary tuberculosis initially entitled after August 19, 1968.

	Rat- ing
For 1 year after date of inactivity, following active tuberculosis	100

 $[34\ FR\ 5062,\ Mar.\ 11,\ 1969.\ Redesignated\ at\ 59\ FR\ 60902,\ Nov.\ 29,\ 1994]$

§ 4.89 Ratings for inactive nonpulmonary tuberculosis in effect on August 19, 1968.

Public Law 90–493 repealed section 356 of title 38, United States Code which provided graduated ratings for inactive tuberculosis. The repealed section, however, still applies to the case of any veteran who on August 19, 1968, was receiving or entitled to receive compensation for tuberculosis. The use of the protective provisions of Pub. L. 90–493 should be mentioned in the discussion portion of all ratings in which these provisions are applied. For use in

rating cases in which the protective provisions of Pub. L. 90–493 apply, the former evaluations are retained in this section.

	Rat- ing
For 2 years after date of inactivity, following active tuberculosis, which was clinically identified during service or subsequently. Thereafter, for 4 years, or in any event, to 6 years after date of inactivity	100 50 30 0

[29 FR 6718, May 22, 1964, as amended at 34 FR 5062, Mar. 11, 1969; 43 FR 45361, Oct. 2, 1978]

THE RESPIRATORY SYSTEM

§ 4.96 Special provisions regarding evaluation of respiratory conditions.

- (a) Rating coexisting respiratory conditions. Ratings under diagnostic codes 6600 through 6817 and 6822 through 6847 will not be combined with each other. Where there is lung or pleural involvement, ratings under diagnostic codes 6819 and 6820 will not be combined with each other or with diagnostic codes 6600 through 6817 or 6822 through 6847. A single rating will be assigned under the diagnostic code which reflects the predominant disability with elevation to the next higher evaluation where the severity of the overall disability warrants such elevation. However, in cases protected by the provisions of Pub. L. 90-493, the graduated ratings of 50 and 30 percent for inactive tuberculosis will not be elevated.
- (b) Rating "protected" tuberculosis cases. Public Law 90-493 repealed section 356 of title 38. United States Code which had provided graduated ratings for inactive tuberculosis. The repealed section, however, still applies to the case of any veteran who on August 19, 1968, was receiving or entitled to receive compensation for tuberculosis. The use of the protective provisions of Pub. L. 90-493 should be mentioned in the discussion portion of all ratings in which these provisions are applied. For application in rating cases in which the protective provisions of Pub. L. 90-493 apply the former evaluations pertaining to pulmonary tuberculosis are retained in §4.97.
- (c) Special monthly compensation. When evaluating any claim involving complete organic aphonia, refer to §3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation. Footnotes in the schedule indicate conditions which potentially establish entitlement to special monthly compensation; however, there are other conditions in this section which under certain circumstances also establish entitlement to special monthly compensation.
- (d) Special provisions for the application of evaluation criteria for diagnostic codes 6600, 6603, 6604, 6825–6833, and 6840–

- 6845. (1) Pulmonary function tests (PFT's) are required to evaluate these conditions except:
- (i) When the results of a maximum exercise capacity test are of record and are 20 ml/kg/min or less. If a maximum exercise capacity test is not of record, evaluate based on alternative criteria.
- (ii) When pulmonary hypertension (documented by an echocardiogram or cardiac catheterization), cor pulmonale, or right ventricular hypertrophy has been diagnosed.
- (iii) When there have been one or more episodes of acute respiratory fail-
- (iv) When outpatient oxygen therapy is required.
- (2) If the DLCO (SB) (Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method) test is not of record, evaluate based on alternative criteria as long as the examiner states why the test would not be useful or valid in a particular case.
- (3) When the PFT's are not consistent with clinical findings, evaluate based on the PFT's unless the examiner states why they are not a valid indication of respiratory functional impairment in a particular case.
- (4) Post-bronchodilator studies are required when PFT's are done for disability evaluation purposes except when the results of pre-bronchodilator pulmonary function tests are normal or when the examiner determines that post-bronchodilator studies should not be done and states why.
- (5) When evaluating based on PFT's, use post-bronchodilator results in applying the evaluation criteria in the rating schedule unless the post-bronchodilator results were poorer than the pre-bronchodilator results. In those cases, use the pre-bronchodilator values for rating purposes.
- (6) When there is a disparity between the results of different PFT's (FEV-1 (Forced Expiratory Volume in one second), FVC (Forced Vital Capacity), etc.), so that the level of evaluation would differ depending on which test result is used, use the test result that the examiner states most accurately reflects the level of disability.
- (7) If the FEV-1 and the FVC are both greater than 100 percent, do not assign

Department of Veterans Affairs

a compensable evaluation based on a decreased FEV–1/FVC ratio.

(Authority: 38 U.S.C. 1155)

 $[34\ {\rm FR}\ 5062,\ {\rm Mar}.\ 11,\ 1969,\ {\rm as}\ {\rm amended}\ {\rm at}\ 61\ {\rm FR}\ 46727,\ {\rm Sept.}\ 5,\ 1996;\ 71\ {\rm FR}\ 52459,\ {\rm Sept.}\ 6,\ 2006]$

§4.97 Schedule of ratings—respiratory system.

		Ratin
	DISEASES OF THE NOSE AND THROAT	
	Septum, nasal, deviation of: Traumatic only, With 50-percent obstruction of the nasal passage on both sides or complete obstruction on one side Nose, loss of part of, or scars: Exposing both nasal passages Loss of part of one ala, or other obvious disfigurement	:
lote:	Or evaluate as DC 7800, scars, disfiguring, head, face, or neck.	
5510 5511 5512 5513	Sinusitis, pansinusitis, chronic. Sinusitis, ethmoid, chronic. Sinusitis, frontal, chronic. Sinusitis, maxillary, chronic. Sinusitis, sphenoid, chronic. Sinusitis, sphenoid, chronic. General Rating Formula for Sinusitis (DC's 6510 through 6514): Following radical surgery with chronic osteomyelitis, or; near constant sinusitis characterized by headaches, pain and tenderness of affected sinus, and purulent discharge or crusting after re-	
	peated surgeries Three or more incapacitating episodes per year of sinusitis requiring prolonged (lasting four to six weeks) antibiotic treatment, or; more than six non-incapacitating episodes per year of sinusitis characterized by headaches, pain, and purulent discharge or crusting One or two incapacitating episodes per year of sinusitis requiring prolonged (lasting four to six weeks) antibiotic treatment, or; three to six non-incapacitating episodes per year of sinusitis characterized by headaches, pain, and purulent discharge or crusting	
	Detected by X-ray only	
	Laryngitis, tuberculous, active or inactive. Rate under §§ 4.88c or 4.89, whichever is appropriate.	
516	Laryngitis, chronic: Hoarseness, with thickening or nodules of cords, polyps, submucous infiltration, or pre-malignant changes on biopsy	
518	Hoarseness, with inflammation of cords or mucous membrane Laryngectomy, total. Rate the residuals of partial laryngectomy as laryngitis (DC 6516), aphonia (DC 6519), or stenosis of larynx (DC 6520).	11
519	Aphonia, complete organic: Constant inability to communicate by speech Constant inability to speak above a whisper	11
	Note: Evaluate incomplete aphonia as laryngitis, chronic (DC 6516).	
6520	Larynx, stenosis of, including residuals of laryngeal trauma (unilateral or bilateral): Forced expiratory volume in one second (FEV-1) less than 40 percent of predicted value, with Flow-Volume Loop compatible with upper airway obstruction, or; permanent tracheostomy FEV-1 of 40- to 55-percent predicted, with Flow-Volume Loop compatible with upper airway obstruction FEV-1 of 56- to 70-percent predicted, with Flow-Volume Loop compatible with upper airway obstruction FEV-1 of 71- to 80-percent predicted, with Flow-Volume Loop compatible with upper airway obstruction	1
	Note: Or evaluate as aphonia (DC 6519).	
521	Pharynx, injuries to: Stricture or obstruction of pharynx or nasopharynx, or; absence of soft palate secondary to trauma, chemical burn, or granulomatous disease, or; paralysis of soft palate with swallowing difficulty (nasal regurgitation) and speech impairment	
522	Allergic or vasomotor rhinitis: With polyps	
523	Without polyps, but with greater than 50-percent obstruction of nasal passage on both sides or complete obstruction on one side	
	Rhinoscleroma	
524	both sides or complete obstruction on one side	

		Ratin
	Wegener's granulomatosis, lethal midline granuloma	10
	DISEASES OF THE TRACHEA AND BRONCHI	
6600	Bronchitis, chronic: FEV-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to Forced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor pulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo or cardiac catheterization), or; episode(s) of acute respiratory failure, or; requires outpatient oxygen therapy FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)	10
	FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent	
6601	predicted Bronchiectasis: With incapacitating episodes of infection of at least six weeks total duration per year With incapacitating episodes of infection of four to six weeks total duration per year, or; near constant findings of cough with purulent sputum associated with anorexia, weight loss, and frank hemoptysis and requiring antibiotic usage almost continuously	10
	cough with sputum that is at times purulent or blood-tinged and that requires prolonged (lasting four to six weeks) antibiotic usage more than twice a year	:
	Note: An incapacitating episode is one that requires bedrest and treatment by a physician.	
6602	Asthma, bronchial: FEV-1 less than 40-percent predicted, or; FEV-1/FVC less than 40 percent, or; more than one attack per week with episodes of respiratory failure, or; requires daily use of systemic (oral or parenteral) high dose corticosteroids or immuno-suppressive medications FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; at least monthly visits to a physician for required care of exacerbations, or; intermittent (at least three per year) courses of systemic (oral or parenteral) corticosteroids FEV-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; daily inhalational or oral bronchodilator therapy, or; inhalational anti-inflammatory medication FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; intermittent inhalational or oral bronchodilator therapy	10
	Note: In the absence of clinical findings of asthma at time of examination, a verified history of asthmatic attacks must be of record.	
	Emphysema, pulmonary: FEV-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to Forced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor pulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo or cardiac catheterization), or; episode(s) of acute respiratory failure, or; requires outpatient oxygen therapy. FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit). FEV-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65-percent predicted FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent predicted	10 6 3
8604	Chronic obstructive pulmonary disease: FEV-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to Forced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor pulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo or cardiac catheterization), or; episode(s) of acute respiratory failure, or; requires outpatient oxygen therapy. FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)	10
	F	

		Rating
trea repo bero 1 to Vet Note inac thou	Tuberculosis, pulmonary, chronic, moderately advanced, active Tuberculosis, pulmonary, chronic, minimal, active Tuberculosis, pulmonary, chronic, active, advancement unspecified Tuberculosis, pulmonary, chronic, mactive, advanced, inactive. Tuberculosis, pulmonary, chronic, moderately advanced, inactive. Tuberculosis, pulmonary, chronic, minimal, inactive. Tuberculosis, pulmonary, chronic, inactive, advancement unspecified. General Rating Formula for Inactive Pulmonary Tuberculosis: For two years after date of inactivity, following active tuberculosis, which was clinically identified during service or subsequently Thereafter for four years, or in any event, to six years after date of inactivity Thereafter, for five years, or to eleven years after date of inactivity Following far advanced lesions diagnosed at any time while the disease process was active, minimum Following moderately advanced lesions, provided there is continued disability, emphysema, dyspnea on exertion, impairment of health, etc Otherwise (1): The 100-percent rating under codes 6701 through 6724 is not subject to a requirement of precedent hospital thment. It will be reduced to 50 percent for failure to submit to examination or to follow prescribed treatment upon or to that effect from the medical authorities. When a veteran is placed on the 100-percent rating for inactive tubulosis, the medical authorities will be appropriately notified of the fact, and of the necessity, as given in footnote as Service Center in the event of failure to submit to examination or to follow treatment. (2): The graduated 50-percent and 30-percent ratings and the permanent 30 percent and 20 percent ratings for inaccive tubulosis, the rating will be for removal of ribs combined with ratings for other respiratory disabilities. Following tracoplasty the rating will be for removal of ribs combined with ratings for other respiratory disabilities.	Rating 100 100 100 100 100 20 0
	Ratings for Pulmonary Tuberculosis Initially Evaluated After August 19, 1968	
	Tuberculosis, pulmonary, chronic, active	100
	NONTUBERCULOUS DISEASES	
6817	Pulmonary Vascular Disease: Primary pulmonary hypertension, or; chronic pulmonary thromboembolism with evidence of pulmonary hypertension, right ventricular hypertrophy, or cor pulmonale, or; pulmonary hypertension secondary to other obstructive disease of pulmonary arteries or veins with evidence of right ventricular hypertrophy or cor pulmonale	100 60 30 0
	that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals.	

		Rating
	Bacterial Infections of the Lung	
	Actinomycosis.	
6823 6824	Nocardiosis. Chronic lung abscess. General Rating Formula for Bacterial Infections of the Lung (diagnostic codes 6822 through 6824): Active infection with systemic symptoms such as fever, night sweats, weight loss, or hemoptysis Depending on the specific findings, rate residuals as interstitial lung disease, restrictive lung disease, or, when obstructive lung disease is the major residual, as chronic bronchitis (DC 6600).	100
	Interstitial Lung Disease	
6825	3	
6826 6827		
6828	Eosinophilic granuloma of lung.	
6829 6830		
6831	Hypersensitivity pneumonitis (extrinsic allergic alveolitis).	
6832 6833		
	General Rating Formula for Interstitial Lung Disease (diagnostic codes 6825 through 6833): Forced Vital Capacity (FVC) less than 50-percent predicted, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption with cardiorespiratory limitation, or; cor pulmonale or pulmonary hypertension, or; requires outpatient oxygen therapy FVC of 50- to 64-percent predicted, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum exercise capacity of 15 to 20 ml/kg/min oxygen consumption with cardiorespiratory limitation FVC of 65- to 74-percent predicted, or; DLCO (SB) of 56- to 65-percent predicted FVC of 75- to 80-percent predicted, or; DLCO (SB) of 66- to 80-percent predicted	100 60 30 10
	Mycotic Lung Disease	
6834 6835 6836 6837 6838 6839	Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis. General Rating Formula for Mycotic Lung Disease (diagnostic codes 6834 through 6839): Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis Chronic pulmonary mycosis requiring suppressive therapy with no more than minimal symptoms such as occasional minor hemoptysis or productive cough	100 50 30 0
	to many years after the initial infection which may have been unrecognized. Accordingly, when service con- nection is under consideration in the absence of record or other evidence of the disease in service, service in southwestern United States where the disease is endemic and absence of prolonged residence in this lo- cality before or after service will be the deciding factor.	
	Restrictive Lung Disease	
6840 6841 6842 6843 6844 6845	Traumatic chest wall defect, pneumothorax, hernia, etc. Post-surgical residual (lobectomy, pneumonectomy, etc.). Chronic pleural effusion or fibrosis. General Rating Formula for Restrictive Lung Disease (diagnostic codes 6840 through 6845): FEV-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to Forced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor pulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo or cardiac catheterization), or; episode(s) of acute respiratory failure, or; requires outpatient oxygen therapy FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)	100
	FEV-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65- percent predicted	30
	pototik produced	. 50

		Rating
	FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent predicted	10
	Note (1): A 100-percent rating shall be assigned for pleurisy with empyema, with or without pleurocutaneous fistula, until resolved.	
	Note (2): Following episodes of total spontaneous pneumothorax, a rating of 100 percent shall be assigned as of the date of hospital admission and shall continue for three months from the first day of the month after hospital discharge.	
	Note (3): Gunshot wounds of the pleural cavity with bullet or missile retained in lung, pain or discomfort on exertion, or with scattered rales or some limitation of excursion of diaphragm or of lower chest expansion shall be rated at least 20-percent disabling. Disabling injuries of shoulder girdle muscles (Groups I to IV) shall be separately rated and combined with ratings for respiratory involvement. Involvement of Muscle Group XXI (DC 5321), however, will not be separately rated.	
6846	Sarcoidosis: Cor pulmonale, or; cardiac involvement with congestive heart failure, or; progressive pulmonary disease with fever, night sweats, and weight loss despite treatment. Pulmonary involvement requiring systemic high dose (therapeutic) corticosteroids for control	100 60 30
	Chronic hilar adenopathy or stable lung infiltrates without symptoms or physiologic impairment	0
6847	Sleep Apnea Syndromes (Obstructive, Central, Mixed): Chronic respiratory failure with carbon dioxide retention or cor pulmonale, or; requires tracheostomy Requires use of breathing assistance device such as continuous airway pressure (CPAP) machine Persistent day-time hypersomnolence Asymptomatic but with documented sleep disorder breathing	100 50 30

 $^{^{\}rm 1}\,\mbox{Review}$ for entitlement to special monthly compensation under $\S\,3.350$ of this chapter.

[61 FR 46728, Sept. 5, 1996, as amended at 71 FR 28586, May 17, 2006]

THE CARDIOVASCULAR SYSTEM

§ 4.100 Application of the evaluation criteria for diagnostic codes 7000– 7007, 7011, and 7015–7020.

- (a) Whether or not cardiac hypertrophy or dilatation (documented by electrocardiogram, echocardiogram, or X-ray) is present and whether or not there is a need for continuous medication must be ascertained in all cases.
- (b) Even if the requirement for a 10% (based on the need for continuous medication) or 30% (based on the presence of cardiac hypertrophy or dilatation) evaluation is met, METs testing is required in all cases except:
- (1) When there is a medical contraindication.
- (2) When the left ventricular ejection fraction has been measured and is 50% or less.
- (3) When chronic congestive heart failure is present or there has been more than one episode of congestive heart failure within the past year.
- (4) When a 100% evaluation can be assigned on another basis.
- (c) If left ventricular ejection fraction (LVEF) testing is not of record,

evaluate based on the alternative criteria unless the examiner states that the LVEF test is needed in a particular case because the available medical information does not sufficiently reflect the severity of the veteran's cardiovascular disability.

[71 FR 52460, Sept. 6, 2006]

§§ 4.101-4.103 [Reserved]

§ 4.104 Schedule of ratings—cardiovascular system.

DISEASES OF THE HEART

Rating

NOTE (1): Evaluate cor pulmonale, which is a form of secondary heart disease, as part of the pulmonary condition that causes it.

NOTE (2): One MET (metabolic equivalent) is the energy cost of standing quietly at rest and represents an oxygen uptake of 3.5 millillers per kilogram of body weight per minute. When the level of METs at which dyspnea, fatigue, angina, dizziness, or syncope develops is required for evaluation, and a laboratory determination of METs by exercise testing cannot be done for medical reasons, an estimation by a medical examiner of the level of activity (expressed in METs and supported by specific examples, such as slow stair climbing or shoveling snow) that results in dyspnea, fatigue, angina, dizziness, or syncope may be used.

38 CFR Ch. I (7-1-21 Edition)

DISEASES OF THE HEART—Continued

		Rat- ing			Rat- ing
7000 Valvular heart disease (including rheumatic heart disease):				Thereafter, with documented pericarditis resulting in:	
	During active infection with valvular heart damage and for three months following cessation of therapy for the active infec- tion	100		Chronic congestive heart failure, or; work- load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection	
	Thereafter, with valvular heart disease (doc- umented by findings on physical examina- tion and either echocardiogram, Doppler echocardiogram, or cardiac catheteriza-			fraction of less than 30 percent. More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater	100
	tion) resulting in: Chronic congestive heart failure, or; work- load of 3 METs or less results in dyspnea,			than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ven- tricular dysfunction with an ejection frac- tion of 30 to 50 percent	60
	fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent	100		Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or;	60
	More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue,			evidence of cardiac hypertrophy or dilata- tion on electro-cardiogram, echocardio- gram, or X-ray	30
	angina, dizziness, or syncope, or; left ven- tricular dysfunction with an ejection frac- tion of 30 to 50 percent	60		Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or;	
	Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or;		7003	continuous medication required Pericardial adhesions: Chronic congestive heart failure, or; work-	10
	evidence of cardiac hypertrophy or dilata- tion on electro-cardiogram, echocardio- gram, or X-ray	30		load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent	100
	Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required	10		More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater	100
7001	Endocarditis: For three months following cessation of therapy for active infection with cardiac in-	10		than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection frac-	
	volvement	100		tion of 30 to 50 percent	60
	cardiogram, or cardiac catheterization) resulting in: Chronic congestive heart failure, or; work-			tion on electro-cardiogram, echocardiogram, or X-ray	30
	load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent	100	7004	greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required	10
	More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ven-	.00	7004	Chronic congestive heart failure, or; work- load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection	100
	tricular dysfunction with an ejection frac- tion of 30 to 50 percent	60		fraction of less than 30 percent	100
	greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilata-			than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ven- tricular dysfunction with an ejection frac-	
	tion on electrocardiogram, echocardio- gram, or X-ray	30		tion of 30 to 50 percent	60
7002		10		evidence of cardiac hypertrophy or dilata- tion on electrocardiogram, echocardio- gram, or X-ray	30
	For three months following cessation of therapy for active infection with cardiac involvement	100		Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required	10

§4.104

DISEASES OF THE HEART—Continued

	Rat- ing		Rat- ing
NOTE: Evaluate syphilitic aortic aneurysms under DC 7110 (aortic aneurysm).		More than one episode of acute congestive heart failure in the past year, or; workload	
005 Arteriosclerotic heart disease (Coronary artery disease):		of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue,	
With documented coronary artery disease resulting in:		angina, dizziness, or syncope, or; left ven- tricular dysfunction with an ejection frac-	
Chronic congestive heart failure, or; work-		tion of 30 to 50 percent	6
load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or;		greater than 7 METs results in dyspnea,	
left ventricular dysfunction with an ejection fraction of less than 30 percent	100	fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilata-	
More than one episode of acute congestive	100	tion on electrocardiogram, echocardio- gram, or X-ray	;
heart failure in the past year, or; workload of greater than 3 METs but not greater		Workload of greater than 7 METs but not	
than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ven-		greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or;	
tricular dysfunction with an ejection frac-		continuous medication required	
tion of 30 to 50 percent	60	Rate under the appropriate cardiovascular diag-	
greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or;		nostic code, depending on particular findings. 7010 Supraventricular arrhythmias:	
evidence of cardiac hypertrophy or dilata-		Paroxysmal atrial fibrillation or other supra-	
tion on electrocardiogram, echocardiogram, or X-ray	30	ventricular tachycardia, with more than four episodes per year documented by	
Workload of greater than 7 METs but not greater than 10 METs results in dyspnea,		ECG or Holter monitor Permanent atrial fibrillation (lone atrial fibril-	
fatigue, angina, dizziness, or syncope, or;		lation), or; one to four episodes per year	
continuous medication required DTE: If nonservice-connected arteriosclerotic heart	10	of paroxysmal atrial fibrillation or other su- praventricular tachycardia documented by	
disease is superimposed on service-connected val-		ECG or Holter monitor	
vular or other non-arteriosclerotic heart disease, request a medical opinion as to which condition is		For indefinite period from date of hospital	
causing the current signs and symptoms.		admission for initial evaluation and med- ical therapy for a sustained ventricular ar-	
006 Myocardial infarction: During and for three months following myo-		rhythmia, or; for indefinite period from date of hospital admission for ventricular	
cardial infarction, documented by labora- tory tests	100	aneurysmectomy, or; with an automatic	
Thereafter:		implantable Cardioverter-Defibrillator (AICD) in place	1
With history of documented myocardial in- farction, resulting in:		Chronic congestive heart failure, or; work- load of 3 METs or less results in dyspnea,	
Chronic congestive heart failure, or; work-		fatigue, angina, dizziness, or syncope, or;	
load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or;		left ventricular dysfunction with an ejection fraction of less than 30 percent	1
left ventricular dysfunction with an ejection fraction of less than 30 percent	100	More than one episode of acute congestive heart failure in the past year, or; workload	
More than one episode of acute congestive heart failure in the past year, or; workload		of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue,	
of greater than 3 METs but not greater		angina, dizziness, or syncope, or; left ven-	
than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ven-		tricular dysfunction with an ejection fraction of 30 to 50 percent	
tricular dysfunction with an ejection frac- tion of 30 to 50 percent	60	Workload of greater than 5 METs but not greater than 7 METs results in dyspnea,	
Workload of greater than 5 METs but not greater than 7 METs results in dyspnea,		fatigue, angina, dizziness, or syncope, or;	
fatigue, angina, dizziness, or syncope, or;		evidence of cardiac hypertrophy or dilata- tion on electrocardiogram, echocardio-	
evidence of cardiac hypertrophy or dilata- tion on electrocardiogram, echocardio-		gram, or X-ray Workload of greater than 7 METs but not	
gram, or X-ray	30	greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or;	
Workload of greater than 7 METs but not greater than 10 METs results in dyspnea,		continuous medication required	
fatigue, angina, dizziness, or syncope, or; continuous medication required	10		
007 Hypertensive heart disease:			
Chronic congestive heart failure, or; work- load of 3 METs or less results in dyspnea,			
fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection			
fraction of less than 30 percent	100		

38 CFR Ch. I (7-1-21 Edition)

§4.104

DISEASES OF THE HEART—Continued

	Rat- ing		Rat- ing
NOTE: A rating of 100 percent shall be assigned from the date of hospital admission for initial evaluation and medical therapy for a sustained ventricular arrhythmia or for ventricular aneurysmectomy. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter.		NOTE: A rating of 100 percent shall be assigned as of the date of hospital admission for valve replacement. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. 7017 Coronary bypass surgery: For three months following hospital admission for surgery.	
7015 Atrioventricular block: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection		sion for surgery	100
fraction of less than 30 percent	100	left ventricular dysfunction with an ejection fraction of less than 30 percent	100
tion of 30 to 50 percent	60	tion of 30 to 50 percent	60
gram, or X-ray	30	gram, or X-ray	30 10
quired	10	7018 Implantable cardiac pacemakers: For two months following hospital admission for implantation or reimplantation Thereafter: Evaluate as supraventricular arrhythmias (DC 7010), ventricular arrhythmias (DC 7011), or atrioventricular block (DC 7015).	100
a disability. 7016 Heart valve replacement (prosthesis): For indefinite period following date of hos-		Minimum	10
pital admission for valve replacement Thereafter: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection	100	7019 Cardiac transplantation: For an indefinite period from date of hospital admission for cardiac transplantation Thereafter: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea,	100
fraction of less than 30 percent	100	fatigue, angina, dizziness, or syncope, or, left ventricular dysfunction with an ejection fraction of less than 30 percent	100
tricular dysfunction with an ejection frac- tion of 30 to 50 percent	60	angina, dizziness, or syncope, or; left ven- tricular dysfunction with an ejection frac- tion of 30 to 50 percent Minimum	60 30
evidence of cardiac hypertrophy or dilata- tion on electrocardiogram, echocardio- gram, or X-ray	30	Note: A rating of 100 percent shall be assigned as of the date of hospital admission for cardiac trans- plantation. One year following discharge, the ap- propriate disability rating shall be determined by mandatory VA examination. Any change in evalua- tion based upon that or any subsequent examina- tion shall be subject to the provisions of §3.105(e) of this chapter.	

§4.104

DISEASES OF THE HEART—Continued

	Rat-		Rat
Chronic congestive heart failure, or; work-load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent	Rating 100 60 30 10 40 20	Evaluate residuals of surgical correction according to organ systems affected. NOTE: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. 7111 Aneurysm, any large artery: If symptomatic, or; for indefinite period from date of hospital admission for surgical correction	Rating
nantly 90mm. or greater, and isolated systolic hy- pertension means that the systolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm. IOTE (2): Evaluate hypertension due to aortic insuffi-		tion based upon that or any subsequent examina- tion shall be subject to the provisions of §3.105(e) of this chapter. 7112 Aneurysm, any small artery: Asymptomatic	
ciency or hyperthyroidism, which is usually the iso- lated systolic type, as part of the condition causing it rather than by a separate evaluation. IOTE (3): Evaluate hypertension separately from hy- pertensive heart disease and other types of heart		Note: If symptomatic, evaluate according to body system affected. Following surgery, evaluate re- siduals under the body system affected. 7113 Arteriovenous fistula, traumatic:	
disease. 110 Aortic aneurysm:		With high output heart failure Without heart failure but with enlarged heart, wide pulse pressure, and tachycardia	10
If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction (including any type of part in		Without cardiac involvement but with edema, stasis dermatitis, and either ulceration or cellulitis:	
correction (including any type of graft in- sertion) Precluding exertion	100 60	Lower extremity Upper extremity With edema or stasis dermatitis:	
· · · · · · · · · · · · · · · · · · ·		Lower extremity	1

38 CFR Ch. I (7-1-21 Edition)

DISEASES OF THE HEART—Continued

	Rat- ing		Rat- ing
Claudication on walking less than 25 yards on a level grade at 2 miles per hour, and; either persistent coldness of the extremity or ankle/brachial index of 0.5 or less		NOTE: For purposes of this section, characteristic attacks consist of sequential color changes of the digits of one or more extremities lasting minutes to hours, sometimes with pain and paresthesias, and precipitated by exposure to cold or by emotional upsets. These evaluations are for the disease as a whole, regardless of the number of extremities involved or whether the nose and ears are involved. 7118 Angioneurotic edema: Attacks without laryngeal involvement lasting one to seven days or longer and occurring more than eight times a year, or, attacks with laryngeal involvement of any duration occurring more than twice a year. Attacks without laryngeal involvement lasting one to seven days and occurring five to eight times a year, or; attacks with laryngeal involvement lasting one or even days and occurring two to four times a year. Attacks without laryngeal involvement lasting one to seven days and occurring two to four times a year. 7119 Erythromelalgia: Characteristic attacks that occur more than once a day, last an average of more than two hours each, respond poorly to treatment, and that restrict most routine daily activities. Characteristic attacks that occur more than once a day, last an average of more than two hours each, and respond poorly to treatment, but that do not restrict most routine daily activities. Characteristic attacks that occur daily or more often but that respond to treatment Characteristic attacks that occur less than daily but at least three times a week and that respond to treatment.	
hour, and; trophic changes (thin skin, absence of hair, dystrophic nails) or ankle/brachial index of 0.7 or less	40	Note: For purposes of this section, a characteristic attack of erythromelalgia consists of burning pain in the hands, feet, or both, usually bilateral and symmetrical, with increased skin temperature and redness, occurring at warm ambient temperatures. These evaluations are for the disease as a whole, regardless of the number of extremities involved.	10
systolic blood pressure at the ankle (determined by Doppler study) divided by the simultaneous brachial artery systolic blood pressure. The normal index is 1.0 or greater. NOTE (2): These evaluations are for involvement of a single extremity. If more than one extremity is affected, evaluate each extremity separately and		7120 Varicose veins: With the following findings attributed to the effects of varicose veins: Massive board-like edema with constant pain at rest Persistent edema or subcutaneous induration, stasis pigmentation or eczema, and persistent ulceration	100
combine (under $\S 4.25$), using the bilateral factor ($\S 4.26$), if applicable.		Persistent edema and stasis pigmentation or eczema, with or without intermittent ulcer-	
7117 Raynaud's syndrome: With two or more digital ulcers plus autoamputation of one or more digits and		ation Persistent edema, incompletely relieved by elevation of extremity, with or without be-	40
history of characteristic attacks	100 60	ginning stasis pigmentation or eczema Intermittent edema of extremity or aching and fatigue in leg after prolonged standing or walking, with symptoms relieved by	20
Characteristic attacks occurring at least daily Characteristic attacks occurring four to six	40	elevation of extremity or compression ho- siery	10
times a week	20	veins	0
times a week	10	single extremity. If more than one extremity is involved, evaluate each extremity separately and combine (under §4.25), using the bilateral factor (§4.26), if applicable.	

DISEASES OF THE HEART—Continued

Rating

100

60

7121	Post-phlebitic syndrome of any etiology:
	With the following findings attributed to venous disease:
	Massive board-like edema with constant pain at rest
	Persistent edema or subcutaneous induration, stasis pigmentation or eczema, and persistent ulcera- tion
	Persistent edema and stasis pig- mentation or eczema, with or without intermittent ulceration
	Persistent edema, incompletely re- lieved by elevation of extremity, with or without beginning stasis pigmentation or eczema
	Intermittent edema of extremity or aching and fatigue in leg after prolonged standing or walking, with symptoms relieved by elevation of extremity or compression hosiery
	Asymptomatic palpable or visible varicose veins

NOTE: These evaluations are for involvement of a single extremity. If more than one extremity is involved, evaluate each extremity separately and combine (under §4.25), using the bilateral factor (§4.26), if applicable.

7122 Cold injury residuals:

With the following in affected parts:

Arthralgia or other pain, numbness, or cold sensitivity plus two or more of the following: tissue loss, nail abnormalities, color changes, locally impaired sensation, hyperhidrosis, X-ray abnormalities (osteoporosis, subarticular punched out lesions, or osteo-arthritis)

Arthralgia or other pain, numbness, or cold sensitivity plus tissue loss, nail abnormalities, color changes, locally impaired sensation, hyperhidrosis, or X-ray abnormalities (osteoporosis, subarticular punched out lesions, or osteoarthritis)

Arthralgia or other pain, numbness, or cold sensitivity

NOTE (1): Separately evaluate amputations of fingers or toes, and complications such as squamous cell carcinoma at the site of a cold injury scar or peripheral neuropathy, under other diagnostic codes. Separately evaluate other disabilities that have been diagnosed as the residual effects of cold injury, such as Raynaud's phenomenon, muscle atrophy, etc., unless they are used to support an evaluation under diagnostic code 7122.

NOTE (2): Evaluate each affected part (e.g., hand, foot, ear, nose) separately and combine the ratings in accordance with §§ 4.25 and 4.26.

7123 Soft tissue sarcoma (of vascular origin)

DISEASES OF THE HEART—Continued

N A 11 (400) 1 1 1 1	
NOTE: A rating of 100 percent shall continue beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals.	

40 (Authority: 38 U.S.C. 1155)

[62 FR 65219, Dec. 11, 1997, as amended at 63 FR 37779, July 14, 1998; 71 FR 52460, Sept. 6, 2006; 79 FR 2100, Jan. 13, 2014; 82 FR 50804, Nov. 2, 2017]

THE DIGESTIVE SYSTEM

§ 4.110 Ulcers.

Experience has shown that the term "peptic ulcer" is not sufficiently specific for rating purposes. Manifest differences in ulcers of the stomach or duodenum in comparison with those at an anastomotic stoma are sufficiently recognized as to warrant two separate graduated descriptions. In evaluating the ulcer, care should be taken that the findings adequately identify the particular location.

§4.111 Postgastrectomy syndromes.

There are various postgastrectomy symptoms which may occur following anastomotic operations of the stomach. When present, those occurring during or immediately after eating and known as the "dumping syndrome" are characterized by gastrointestinal complaints and generalized symptoms simulating hypoglycemia; those occurring from 1 to 3 hours after eating usually present definite manifestations of hypoglycemia.

§ 4.112 Weight loss.

For purposes of evaluating conditions in §4.114, the term "substantial weight loss" means a loss of greater than 20 percent of the individual's baseline weight, sustained for three months or longer; and the term "minor weight loss" means a weight loss of 10 to 20 percent of the individual's baseline weight, sustained for three months or longer. The term "inability to gain

weight" means that there has been substantial weight loss with inability to regain it despite appropriate therapy. "Baseline weight" means the average weight for the two-year-period preceding onset of the disease.

(Authority: 38 U.S.C. 1155) [66 FR 29488, May 31, 2001]

§ 4.113 Coexisting abdominal conditions.

There are diseases of the digestive system, particularly within the abdomen, which, while differing in the site of pathology, produce a common disability picture characterized in the main by varying degrees of abdominal distress or pain, anemia and disturbances in nutrition. Consequently, certain coexisting diseases in this area, as indicated in the instruction under the title "Diseases of the Digestive System," do not lend themselves to distinct and separate disability evaluations without violating the fundamental principle relating pyramiding as outlined in §4.14.

\$4.114 Schedule of ratings—digestive system.

Ratings under diagnostic codes 7301 to 7329, inclusive, 7331, 7342, and 7345 to 7348 inclusive will not be combined with each other. A single evaluation will be assigned under the diagnostic code which reflects the predominant disability picture, with elevation to the next higher evaluation where the severity of the overall disability warrants such elevation.

	Rat- ing
7200 Mouth, injuries of. Rate as for disfigurement and impairment of function of mastication. 7201 Lips, injuries of. Rate as for disfigurement of face. 7202 Tongue, loss of whole or part: With inability to communicate by speech One-half or more	100 60 30
impairment of general health Severe, permitting liquids only Moderate 7204 Esophagus, spasm of (cardiospasm). If not amenable to dilation, rate as for the degree of obstruction (stricture). 7205 Esophagus, diverticulum of, acquired. Rate as for obstruction (stricture). 7301 Pertiponeum adhesions of	80 50 30

38 CFR Ch. I (7-1-21 Edi	tion)
	Rat- ing
Severe; definite partial obstruction shown by X-	
ray, with frequent and prolonged episodes of	
severe colic distension, nausea or vomiting, following severe peritonitis, ruptured appendix,	
perforated ulcer, or operation with drainage	50
Moderately severe; partial obstruction mani-	
fested by delayed motility of barium meal and	
less frequent and less prolonged episodes of pain	30
Moderate; pulling pain on attempting work or ag-	
gravated by movements of the body, or occa-	
sional episodes of colic pain, nausea, con-	
stipation (perhaps alternating with diarrhea) or abdominal distension	10
Mild	
NOTE: Ratings for adhesions will be considered	
when there is history of operative or other traumatic or infectious (intraabdominal) proc-	
ess, and at least two of the following: disturb-	
ance of motility, actual partial obstruction, re-	
flex disturbances, presence of pain.	
7304 Ulcer, gastric. 7305 Ulcer, duodenal:	
Severe; pain only partially relieved by standard	
ulcer therapy, periodic vomiting, recurrent hematemesis or melena, with manifestations	
hematemesis or melena, with manifestations	
of anemia and weight loss productive of defi- nite impairment of health	60
Moderately severe; less than severe but with im-	
pairment of health manifested by anemia and	
weight loss; or recurrent incapacitating epi-	
sodes averaging 10 days or more in duration at least four or more times a year	40
Moderate; recurring episodes of severe symp-	
toms two or three times a year averaging 10	
days in duration; or with continuous moderate manifestations	20
Mild; with recurring symptoms once or twice	_`
yearly	10
7306 Ulcer, marginal (gastrojejunal): Pronounced; periodic or continuous pain	
unrelieved by standard ulcer therapy with peri-	
odic vomiting, recurring melena or	
hematemesis, and weight loss. Totally inca-	
pacitating	100
nounced and less continuous symptoms with	
definite impairment of health	60
Moderately severe; intercurrent episodes of ab- dominal pain at least once a month partially or	
completely relieved by ulcer therapy, mild and	
transient episodes of vomiting or melena	40
Moderate; with episodes of recurring symptoms	
several times a year	20
once or twice yearly	10
'307 Gastritis, hypertrophic (identified by	
gastroscope):	
Chronic; with severe hemorrhages, or large ul- cerated or eroded areas	60
Chronic; with multiple small eroded or ulcerated	
areas, and symptoms	30
Chronic; with small nodular lesions, and symptoms	10
Gastritis, atrophic.	10
A complication of a number of diseases, includ-	
ing pernicious anemia.	
Rate the underlying condition. 7308 Postgastrectomy syndromes:	
Severe; associated with nausea, sweating, cir-	
culatory disturbance after meals, diarrhea,	
hypoglycemic symptoms, and weight loss with	
malnutrition and anemia	60

malnutrition and anemia

	Rat-		Rat-
	ing		ing
Moderate; less frequent episodes of epigastric disorders with characteristic mild circulatory		Asymptomatic	0
symptoms after meals but with diarrhea and	40	parallel in symptomatology with ulcerative coli-	
weight loss	40	tis and should be rated on the scale provided for the latter. Similarly, lung abscess due to	
with characteristic mild circulatory symptoms		amebiasis will be rated under the respiratory	
or continuous mild manifestations	20	system schedule, diagnostic code 6809.	
7309 Stomach, stenosis of.		7322 Dysentery, bacillary.	
Rate as for gastric ulcer.		Rate as for ulcerative colitis	
7310 Stomach, injury of, residuals.		7323 Colitis, ulcerative:	
Rate as peritoneal adhesions.		Pronounced; resulting in marked malnutrition,	
7311 Residuals of injury of the liver: Depending on the specific residuals, separately		anemia, and general debility, or with serious complication as liver abscess	100
evaluate as adhesions of peritoneum (diag-		Severe; with numerous attacks a year and mal-	100
nostic code 7301), cirrhosis of liver (diagnostic		nutrition, the health only fair during remissions	60
code 7312), and chronic liver disease without		Moderately severe; with frequent exacerbations	30
cirrhosis (diagnostic code 7345).		Moderate; with infrequent exacerbations	10
7312 Cirrhosis of the liver, primary biliary cirrhosis,		7324 Distomiasis, intestinal or hepatic: Severe symptoms	30
or cirrhotic phase of sclerosing cholangitis:		Moderate symptoms	10
Generalized weakness, substantial weight loss, and persistent jaundice, or; with one of the fol-		Mild or no symptoms	0
lowing refractory to treatment: ascites, hepatic		7325 Enteritis, chronic.	
encephalopathy, hemorrhage from varices or		Rate as for irritable colon syndrome.	
portal gastropathy (erosive gastritis)	100	7326 Enterocolitis, chronic.	
History of two or more episodes of ascites, he-		Rate as for irritable colon syndrome. 7327 Diverticulitis.	
patic encephalopathy, or hemorrhage from		Rate as for irritable colon syndrome, peritoneal	
varices or portal gastropathy (erosive gas-		adhesions, or colitis, ulcerative, depending	
tritis), but with periods of remission between attacks	70	upon the predominant disability picture.	
History of one episode of ascites, hepatic	70	7328 Intestine, small, resection of:	
encephalopathy, or hemorrhage from varices		With marked interference with absorption and	
or portal gastropathy (erosive gastritis)	50	nutrition, manifested by severe impairment of	
Portal hypertension and splenomegaly, with		health objectively supported by examination findings including material weight loss	60
weakness, anorexia, abdominal pain, malaise,		With definite interference with absorption and	00
and at least minor weight loss	30	nutrition, manifested by impairment of health	
Symptoms such as weakness, anorexia, abdom- inal pain, and malaise	10	objectively supported by examination findings	
Note: For evaluation under diagnostic code	10	including definite weight loss	40
7312, documentation of cirrhosis (by biopsy or		Symptomatic with diarrhea, anemia and inability	
imaging) and abnormal liver function tests		to gain weight	20
must be present.		predominant disability, rate under diagnostic	
7314 Cholecystitis, chronic:	30	code 7301.	
Severe; frequent attacks of gall bladder colic Moderate; gall bladder dyspepsia, confirmed by	30	7329 Intestine, large, resection of:	
X-ray technique, and with infrequent attacks		With severe symptoms, objectively supported by	
(not over two or three a year) of gall bladder		examination findings	40
colic, with or without jaundice	10	With moderate symptoms With slight symptoms	20 10
Mild	0	Note: Where residual adhesions constitute the	10
7315 Cholelithiasis, chronic.		predominant disability, rate under diagnostic	
Rate as for chronic cholecystitis. 7316 Cholangitis, chronic.		code 7301.	
Rate as for chronic cholecystitis.		7330 Intestine, fistula of, persistent, or after attempt	
7317 Gall bladder, injury of.		at operative closure:	100
Rate as for peritoneal adhesions.		Copious and frequent, fecal discharge Constant or frequent, fecal discharge	100 60
7318 Gall bladder, removal of:		Slight infrequent, fecal discharge	30
With severe symptoms	30	Healed; rate for peritoneal adhesions.	
With mild symptoms	10	7331 Peritonitis, tuberculous, active or inactive:	
Nonsymptomatic	0	Active	100
See Hemic and Lymphatic Systems.		Inactive: See §§ 4.88b and 4.89.	
7319 Irritable colon syndrome (spastic colitis, mu-		7332 Rectum and anus, impairment of sphincter control:	
cous colitis, etc.):		Complete loss of sphincter control	100
Severe; diarrhea, or alternating diarrhea and		Extensive leakage and fairly frequent involuntary	
constipation, with more or less constant ab-		bowel movements	60
dominal distress	30	Occasional involuntary bowel movements, ne-	
Moderate; frequent episodes of bowel disturb-	10	cessitating wearing of pad	30
ance with abdominal distress	10	Constant slight, or occasional moderate leakage Healed or slight, without leakage	10 0
sional episodes of abdominal distress	0	7333 Rectum and anus, stricture of:	U
7321 Amebiasis:	•	Requiring colostomy	100
Mild gastrointestinal disturbances, lower abdom-		Great reduction of lumen, or extensive leakage	50
inal cramps, nausea, gaseous distention,		Moderate reduction of lumen, or moderate con-	
chronic constipation interrupted by diarrhea	10	stant leakage	30

	Rat- ing		Rat- ing
7334 Rectum, prolapse of:		Near-constant debilitating symptoms (such as fa-	
Severe (or complete), persistent	50	tigue, malaise, nausea, vomiting, anorexia,	
Moderate, persistent or frequently recurring	30	arthralgia, and right upper quadrant pain)	100
Mild with constant slight or occasional moderate		Daily fatigue, malaise, and anorexia, with sub-	
leakage	10	stantial weight loss (or other indication of mal-	
7335 Ano, fistula in.		nutrition), and hepatomegaly, or; incapacitating	
Rate as for impairment of sphincter control.		episodes (with symptoms such as fatigue,	
7336 Hemorrhoids, external or internal:		malaise, nausea, vomiting, anorexia,	
With persistent bleeding and with secondary	20	arthralgia, and right upper quadrant pain) hav-	
anemia, or with fissuresLarge or thrombotic, irreducible, with excessive	20	ing a total duration of at least six weeks during	
redundant tissue, evidencing frequent		the past 12-month period, but not occurring	
recurrences	10	constantly	60
Mild or moderate	0	Daily fatigue, malaise, and anorexia, with minor	
7337 Pruritus ani.	-	weight loss and hepatomegaly, or; incapaci- tating episodes (with symptoms such as fa-	
Rate for the underlying condition.		tique, malaise, nausea, vomiting, anorexia,	
7338 Hernia, inguinal:		arthralgia, and right upper quadrant pain) hav-	
Large, postoperative, recurrent, not well sup-		ing a total duration of at least four weeks, but	
ported under ordinary conditions and not read-		less than six weeks, during the past 12-month	
ily reducible, when considered inoperable	60	period	40
Small, postoperative recurrent, or unoperated ir-		Daily fatigue, malaise, and anorexia (without	
remediable, not well supported by truss, or not		weight loss or hepatomegaly), requiring dietary	
readily reducible	30	restriction or continuous medication, or; inca-	
Postoperative recurrent, readily reducible and	10	pacitating episodes (with symptoms such as	
well supported by truss or belt	10	fatigue, malaise, nausea, vomiting, anorexia,	
Not operated, but remediableSmall, reducible, or without true hernia protru-	0	arthralgia, and right upper quadrant pain) hav-	
sion	0	ing a total duration of at least two weeks, but	
NOTE: Add 10 percent for bilateral involvement,	U	less than four weeks, during the past 12-	
provided the second hernia is compensable.		month period	20
This means that the more severely disabling		Intermittent fatigue, malaise, and anorexia, or;	
hernia is to be evaluated, and 10 percent,		incapacitating episodes (with symptoms such	
only, added for the second hernia, if the latter		as fatigue, malaise, nausea, vomiting, ano-	
is of compensable degree.		rexia, arthralgia, and right upper quadrant	
7339 Hernia, ventral, postoperative:		pain) having a total duration of at least one week, but less than two weeks, during the	
Massive, persistent, severe diastasis of recti		past 12-month period	10
muscles or extensive diffuse destruction or		Nonsymptomatic	0
weakening of muscular and fascial support of		NOTE (1): Evaluate sequelae, such as cirrhosis	Ü
abdominal wall so as to be inoperable	100	or malignancy of the liver, under an appro-	
Large, not well supported by belt under ordinary	40	priate diagnostic code, but do not use the	
conditions	40	same signs and symptoms as the basis for	
Small, not well supported by belt under ordinary		evaluation under DC 7354 and under a diag-	
conditions, or healed ventral hernia or post-op- erative wounds with weakening of abdominal		nostic code for sequelae. (See § 4.14.).	
wall and indication for a supporting belt	20	NOTE (2): For purposes of evaluating conditions	
Wounds, postoperative, healed, no disability, belt		under diagnostic code 7345, "incapacitating	
not indicated	0	episode" means a period of acute signs and	
7340 Hernia, femoral.		symptoms severe enough to require bed rest	
Rate as for inguinal hernia.		and treatment by a physician.	
7342 Visceroptosis, symptomatic, marked	10	NOTE (3): Hepatitis B infection must be con-	
7343 Malignant neoplasms of the digestive system,		firmed by serologic testing in order to evaluate	
exclusive of skin growths	100	it under diagnostic code 7345.	
NOTE: A rating of 100 percent shall continue be-		7346 Hernia hiatal:	
yond the cessation of any surgical, X-ray,		Symptoms of pain, vomiting, material weight loss	
antineoplastic chemotherapy or other thera-		and hematemesis or melena with moderate anemia; or other symptom combinations pro-	
peutic procedure. Six months after discontinu-		ductive of severe impairment of health	60
ance of such treatment, the appropriate dis-		Persistently recurrent epigastric distress with	00
ability rating shall be determined by mandatory		dysphagia, pyrosis, and regurgitation, accom-	
VA examination. Any change in evaluation based upon that or any subsequent examina-		panied by substernal or arm or shoulder pain,	
tion shall be subject to the provisions of		productive of considerable impairment of	
§ 3.105(e) of this chapter. If there has been no		health	30
local recurrence or metastasis, rate on residu-		With two or more of the symptoms for the 30	
als.		percent evaluation of less severity	10
7344 Benign neoplasms, exclusive of skin growths:		7347 Pancreatitis:	
Evaluate under an appropriate diagnostic code,		With frequently recurrent disabling attacks of ab-	
depending on the predominant disability or the		dominal pain with few pain free intermissions	
specific residuals after treatment.		and with steatorrhea, malabsorption, diarrhea	
7345 Chronic liver disease without cirrhosis (includ-		and severe malnutrition	100
ing hepatitis B, chronic active hepatitis, auto-		With frequent attacks of abdominal pain, loss of	
immune hepatitis, hemochromatosis, drug-induced		normal body weight and other findings show-	
hepatitis, etc., but excluding bile duct disorders		ing continuing pancreatic insufficiency be-	
and hepatitis C):		tween acute attacks	60

Department of Veterans Affairs		§4.115	
	Rat- ing		Rat- ing
Moderately severe; with at least 4–7 typical attacks of abdominal pain per year with good remission between attacks	30	Intermittent fatigue, malaise, and anorexia, or; incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least one week, but less than two weeks, during the past 12-month period	10 0
tinuing gastric retention With symptoms and confirmed diagnosis of alkaline gastritis, or of confirmed persisting diarrhea	40 30	under diagnostic code 7354, "incapacitating episode" means a period of acute signs and symptoms severe enough to require bed rest and treatment by a physician.	
Recurrent ulcer with incomplete vagotomy NOTE: Rate recurrent ulcer following complete vagotomy under diagnostic code 7305, min-	20	(Authority: 38 U.S.C. 1155)	
imum rating 20 percent; and rate dumping syndrome under diagnostic code 7308. 7351 Liver transplant: For an indefinite period from the date of hospital admission for transplant surgery	100 30	[29 FR 6718, May 22, 1964, as amended FR 5063, Mar. 11, 1969; 40 FR 42540, Sep 1975; 41 FR 11301, Mar. 18, 1976; 66 FR May 31, 2001]	ot. 15,
as of the date of hospital admission for trans- plant surgery and shall continue. One year fol-		§ 4.115 Nephritis.	
lowing discharge, the appropriate disability rat- ing shall be determined by mandatory VA ex- amination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. 7354 Hepatitis C (or non-A, non-B hepatitis):		Albuminuria alone is not neph nor will the presence of transient min and casts following acute fe illness be taken as nephritis. The merular type of nephritis is us	albu- brile glo-

With serologic evidence of hepatitis C infection and the following signs and symptoms due to hepatitis C infection:

Near-constant debilitating symptoms (such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain)

Daily fatigue, malaise, and anorexia, with substantial weight loss (or other indication of malnutrition), and hepatomegaly, or; incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least six weeks during the past 12-month period, but not occurring constantly

Daily fatigue, malaise, and anorexia, with minor weight loss and hepatomegaly, or; incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least four weeks, but less than six weeks, during the past 12-month period ..

Daily fatique, malaise, and anorexia (without weight loss or hepatomegaly), requiring dietary restriction or continuous medication, or; incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least two weeks, but less than four weeks, during the past 12month period ..

lburile ളിറmerular type of nephritis is usually preceded by or associated with severe infectious disease; the onset is sudden, and the course marked by red blood cells, salt retention, and edema; it may clear up entirely or progress to a chronic condition. The nephrosclerotic type, originating in hypertension or arteriosclerosis, develops slowly, with minimum laboratory findings, and is associated with natural progress. Separate ratings are not to be assigned for disability from disease of the heart and any form of nephritis, on account of the close interrelationships of cardiovascular disabilities. If, however, absence of a kidney is the sole renal disability, even if removal was required because of nephritis, the absent kidney

and any hypertension or heart disease

will be separately rated. Also, in the

event that chronic renal disease has

progressed to the point where regular

20

§4.115a

dialysis is required, any coexisting hypertension or heart disease will be separately rated.

 $[41\ {\rm FR}\ 34258,\ {\rm Aug.}\ 13,\ 1976,\ {\rm as}\ {\rm amended}\ {\rm at}\ 59$ FR 2527, Jan. 18, 1994]

§4.115a Ratings of the genitourinary system—dysfunctions.

Diseases of the genitourinary system generally result in disabilities related to renal or voiding dysfunctions, infections, or a combination of these. The following section provides descriptions of various levels of disability in each of these symptom areas. Where diagnostic codes refer the decisionmaker to these specific areas dysfunction, only the predominant area of dysfunction shall be considered for rating purposes. Since the areas of dysfunction described below do not cover all symptoms resulting from genitourinary diseases, specific diagnoses may include a description of symptoms assigned to that diagnosis.

	F
Renal dysfunction:	
Requiring regular dialysis, or precluding more than sedentary activity from one of the following: persistent edema and albuminuria; or, BUN more than 80mg%; or, creatinine more than 8mg%; or, markedly decreased function of kidney or other organ systems, especially cardiovascular	
Persistent edema and albuminuria with BUN 40 to 80mg%; or, creatinine 4 to 8mg%; or, generalized poor health characterized by lethargy, weakness, anorexia, weight loss, or limitation of exertion	
Constant albuminuria with some edema; or, defi- nite decrease in kidney function; or, hyper- tension at least 40 percent disabling under di- agnostic code 7101	
Albumin constant or recurring with hyaline and granular casts or red blood cells; or, transient or slight edema or hypertension at least 10 percent disabling under diagnostic code 7101	
Albumin and casts with history of acute nephritis; or, hypertension non-compensable under diag- nostic code 7101	
Rate particular condition as urine leakage, fre- quency, or obstructed voiding	
Continual Urine Leakage, Post Surgical Urinary Diversion, Urinary Incontinence, or Stress Incontinence:	
Requiring the use of an appliance or the wearing of absorbent materials which must be changed more than 4 times per day	
Requiring the wearing of absorbent materials which must be changed 2 to 4 times per day	
Requiring the wearing of absorbent materials which must be changed less than 2 times per	

	Rat- ing
Urinary frequency:	
Daytime voiding interval less than one hour, or; awakening to void five or more times per night Daytime voiding interval between one and two	40
hours, or; awakening to void three to four times per night	20
Daytime voiding interval between two and three hours, or; awakening to void two times per	40
night Obstructed voiding:	10
Urinary retention requiring intermittent or continuous catheterization	30
Marked obstructive symptomatology (hesitancy, slow or weak stream, decreased force of stream) with any one or combination of the fol-	
lowing:	
 Post void residuals greater than 150 cc. Uroflowmetry; markedly diminished peak flow rate (less than 10 cc/sec). 	
 Recurrent urinary tract infections sec- ondary to obstruction. 	
Stricture disease requiring periodic dilatation every 2 to 3 months	10
Obstructive symptomatology with or without stric- ture disease requiring dilatation 1 to 2 times	
per year	0
Poor renal function: Rate as renal dysfunction.	
Recurrent symptomatic infection requiring drain-	
age/frequent hospitalization (greater than two	
times/year), and/or requiring continuous inten-	
sive management	30
Long-term drug therapy, 1–2 hospitalizations per year and/or requiring intermittent intensive management	10
manayement	10

 $[59~{\rm FR}~2527,~{\rm Jan.}~18,~1994;~59~{\rm FR}~10676,~{\rm Mar.}~7,$

§4.115b Ratings of the genitourinary system—diagnoses.

		Rat- ing
	Note: When evaluating any claim involving loss or loss of use of one or more creative organs, refer to § 3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation. Footnotes in	
	the schedule indicate conditions which potentially establish entitlement	
	to special monthly compensation; however, there are other conditions in this section which under certain cir- cumstances also establish entitlement to special monthly compensation.	
7500	Kidney, removal of one: Minimum evaluation Or rate as renal dysfunction if there is nephritis, infection, or pathology of	30
	the other.	
7501	Kidney, abscess of:	1

100

80

60

30

60 40

20

Department of Veterans Affairs

§	4.	1	1	5	b

		Rat- ing		Rat- ing
7502	Nephritis, chronic:		7523 Testis, atrophy complete:.	
	Rate as renal dysfunction.		Both—201	
7504	Pyelonephritis, chronic:		One—0 1	
	Rate as renal dysfunction or urinary		7524 Testis, removal:.	
	tract infection, whichever is predominant.		Both—30 ¹	
7505	Kidney, tuberculosis of:		One—0 1	
000	Rate in accordance with §§ 4.88b or		Note: In cases of the removal of one	
7507	4.89, whichever is appropriate. Nephrosclerosis, arteriolar: Rate according to predominant symptoms as renal dysfunction, hypertension or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be		testis as the result of a service-in- curred injury or disease, other than an undescended or congenitally un- developed testis, with the absence or nonfunctioning of the other testis un- related to service, an evaluation of 30 percent will be assigned for the serv- ice-connected testicular loss. Testis,	
	elevated to the next higher evalua- tion.		underscended, or congenitally unde-	
7508	Nephrolithiasis:		veloped is not a ratable disability.	
	Rate as hydronephrosis, except for re-		7525 Epididymo-orchitis, chronic only:	
	current stone formation requiring one		Rate as urinary tract infection.	
	or more of the following:		For tubercular infections: Rate in ac-	
	diet therapy drug therapy		cordance with §§ 4.88b or 4.89,	
	3. invasive or non-invasive proce-		whichever is appropriate.	
	dures more than two times/year	30	7527 Prostate gland injuries, infections, hyper- trophy, postoperative residuals:	
7509	Hydronephrosis:		Rate as voiding dysfunction or urinary	
	Severe; Rate as renal dysfunction. Frequent attacks of colic with infection		tract infection, whichever is predomi-	
	(pyonephrosis), kidney function impaired	30	nant.	
	Frequent attacks of colic, requiring catheter		7528 Malignant neoplasms of the genitourinary sys-	
	drainage	20	tem	100
	Only an occasional attack of colic, not in-	40	Note—Following the cessation of sur-	
7510	fected and not requiring catheter drainage Ureterolithiasis:	10	gical, X-ray, antineoplastic chemo- therapy or other therapeutic proce-	
0.0	Rate as hydronephrosis, except for re-		dure, the rating of 100 percent shall	
	current stone formation requiring one		continue with a mandatory VA exam-	
	or more of the following:		ination at the expiration of six	
	1. diet therapy		months. Any change in evaluation	
	drug therapy invasive or non-invasive proce-		based upon that or any subsequent	
	dures more than two times/year	30	examination shall be subject to the provisions of §3.105(e) of this chap-	
7511	Ureter, stricture of:		ter. If there has been no local reoc-	
	Rate as hydronephrosis, except for re-		currence or metastasis, rate on re-	
	current stone formation requiring one or more of the following:		siduals as voiding dysfunction or	
	1. diet therapy		renal dysfunction, whichever is pre-	
	2. drug therapy		dominant.	
	3. invasive or non-invasive proce-		7529 Benign neoplasms of the genitourinary system:	
7540	dures more than two times/year	30	Rate as voiding dysfunction or renal	
	Cystitis, chronic, includes interstitial and all ogies, infectious and non-infectious:		dysfunction, whichever is predomi-	
51101	Rate as voiding dysfunction.		nant.	
	Bladder, calculus in, with symptoms inter-		7530 Chronic renal disease requiring regular dialy-	
ferin	ng with function:		Sis:	
7516	Rate as voiding dysfunction Bladder, fistula of:		Rate as renal dysfunction.	
510	Rate as voiding dysfunction or urinary		7531 Kidney transplant: Following transplant surgery	100
	tract infection, whichever is predomi-		Thereafter: Rate on residuals as renal	100
	_ nant.		dysfunction, minimum rating	30
7547	Postoperative, suprapubic cystotomy	100	3,44 444 , 444 3	
7517	Bladder, injury of: Rate as voiding dysfunction.			
7518	Urethra, stricture of:			
	Rate as voiding dysfunction.			
′519	Urethra, fistual of:			
	Rate as voiding dysfunction.	100		
520	Multiple urethroperineal fistulae	100 30		
J_U	Or rate as voiding dysfunction.	30		
'521	Penis removal of glans	20		
7521		20		

38 CFR Ch. I (7-1-21 Edition)

Rat-ing Note-The 100 percent evaluation shall be assigned as of the date of hospital admission for transplant surgery and shall continue with a mandatory VA examination one year following hospital discharge. Any change in eval-uation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. 7532 Renal tubular disorders (such as renal glycosurias, aminoacidurias, renal tubular acidosis, Fanconi's syndrome, Bartter's syndrome, related disorders of Henle's loop and proximal or distal nephron function, etc.): Minimum rating for symptomatic condi-20 tion .. Or rate as renal dysfunction. 7533 Cystic diseases of the kidneys (polycystic disease, uremic medullary cystic disease, Medullary sponge kidney, and similar conditions): Rate as renal dysfunction. 7534 Atherosclerotic renal disease (renal artery stenosis or atheroembolic renal disease): Rate as renal dysfunction. 7535 Toxic nephropathy (antibotics, radiocontrast agents, nonsteroidal anti-inflammatory agents, heavy metals, and similar agents): Rate as renal dysfunction. 7536 Glomerulonephritis: Rate as renal dysfunction. Interstitial nephritis: 7537 Rate as renal dysfunction. 7538 Papillary necrosis: Rate as renal dysfunction. 7539 Renal amyloid disease: Rate as renal dysfunction. 7540 Disseminated intravascular coagulation with renal cortical necrosis: Rate as renal dysfunction. 7541 Renal involvement in diabetes mellitus, sickle cell anemia, systemic lupus erythematosus, vasculitis, or other systemic disease processes. Rate as renal dysfunction. 7542 Neurogenic bladder: Rate as voiding dysfunction. ¹ Review for entitlement to special monthly compensation under § 3.350 of this chapter.

[59 FR 2527, Jan. 18, 1994; 59 FR 14567, Mar. 29, 1994, as amended at 59 FR 46339, Sept. 8, 1994]

GYNECOLOGICAL CONDITIONS AND DISORDERS OF THE BREAST

§ 4.116 Schedule of ratings—gynecological conditions and disorders of the breast.

	Rating
Note 1: Natural menopause, prima amenorrhea, and pregnancy and chi birth are not disabilities for rating pip poses. Chronic residuals of medical surgical complications of pregnancy middle be disabilities for rating purposes. Note 2: When evaluating any claim invoing loss or loss of use of one or middle creative organs or anatomical loss one or both breasts, refer to §3.350 this chapter to determine whether the veteran may be entitled to specimentally compensation. Footnotes in the schedule indicate conditions which permitted the specimentally establish entitlement to specimentally establish entitlement to specimentally compensation; however, almost any condition in this section might, undicertain circumstances, establish entitiement to special monthly compensation. 7610 Vulva or clitoris, disease or injury of (including vulvovaginitis)	Id- ur- or ay IV- ore of of he iial he oo- iial ost
7611 Vagina, disease or injury of.7612 Cervix, disease or injury of.	
 7613 Uterus, disease, injury, or adhesions of. 7614 Fallopian tube, disease, injury, or adhesio of (including pelvic inflammatory disease (PID)). 7615 Ovary, disease, injury, or adhesions of. General Rating Formula for Disease, Injury, or Ahesions of Female Reproductive Organs (dia 	Ad-
nostic codes 7610 through 7615): Symptoms not controlled by continuo	us
treatment	30
ment	10
Symptoms that do not require continuo treatment	0 lity ne- an ual
7617 Uterus and both ovaries, removal of, co	m-
plete: For three months after removal	1100
Thereafter	
7618 Uterus, removal of, including corpus: For three months after removal Thereafter	1100
7619 Ovary, removal of:	
For three months after removal Thereafter:	1100
Complete removal of both ovaries	
Removal of one with or withon partial removal of the other	
Note: In cases of the removal of one over as the result of a service-connected jury or disease, with the absence or not functioning of a second ovary unrelat to service, an evaluation of 30 perce will be assigned for the service-conected ovarian loss	ary in- on- ed ent
7620 Ovaries, atrophy of both, complete	1 20

	Rating		Ratin
621 Complete or incomplete pelvic organ		(4) Wide local excision (including	
prolapse due to injury, disease, or surgical com-		partial mastectomy,	
plications of pregnancy	10	lumpectomy, tylectomy,	
	10		
Note: Pelvic organ prolapse occurs when		, , , , , , , , , , , , , , , , , , ,	
a pelvic organ such as bladder, urethra, uterus, vagina, small bowel, or rectum		quadrantectomy) means re- moval of a portion of the breast	
drops (prolapse) from its normal place in		tissue	
the abdomen. Conditions associated with		7627 Malignant neoplasms of gynecological sys-	
			10
pelvic organ prolapse include: uterine or		Note: A rating of 100 percent shall con	10
vaginal vault prolapse, cystocele,		Note: A rating of 100 percent shall con-	
urethrocele, rectocele, enterocele, or any		tinue beyond the cessation of any sur-	
combination thereof. Evaluate pelvic		gical, radiation, antineoplastic chemo-	
organ prolapse under DC 7621. Evaluate		therapy or other therapeutic procedures.	
separately any genitourinary, digestive,		Six months after discontinuance of such	
or skin symptoms under the appropriate		treatment, the appropriate disability rat-	
diagnostic code(s) and combine all eval-		ing shall be determined by mandatory	
uations with the 10 percent evaluation		VA examination. Any change in evalua-	
under DC 7621		tion based upon that or any subsequent	
624 Fistula, rectovaginal:		examination shall be subject to the provi-	
Vaginal fecal leakage at least once a day	400	sions of §3.105(e) of this chapter. Rate	
requiring wearing of pad	100	chronic residuals to include scars,	
Vaginal fecal leakage four or more times		lymphedema, disfigurement, and/or other	
per week, but less than daily, requiring		impairment of function under the appro-	
wearing of pad	60	priate diagnostic code(s) within the ap-	
Vaginal fecal leakage one to three times		propriate body system	
per week requiring wearing of pad	30	7628 Benign neoplasms of gynecological system.	
Vaginal fecal leakage less than once a		Rate chronic residuals to include scars,	
week	10	lymphedema, disfigurement, and/or other impair-	
Without leakage	0	ment of function under the appropriate diagnostic	
625 Fistula, urethrovaginal:		code(s) within the appropriate body system	
Multiple urethrovaginal fistulae	100	7629 Endometriosis:	
Requiring the use of an appliance or the		Lesions involving bowel or bladder con-	
wearing of absorbent materials which		firmed by laparoscopy, pelvic pain or	
must be changed more than four times		heavy or irregular bleeding not controlled	
per day	60	by treatment, and bowel or bladder	
Requiring the wearing of absorbent mate-		symptoms	5
rials which must be changed two to four		Pelvic pain or heavy or irregular bleeding	
times per day	40	not controlled by treatment	3
Requiring the wearing of absorbent mate-		Pelvic pain or heavy or irregular bleeding	
rials which must be changed less than		requiring continuous treatment for control	
two times per day	20	Note: Diagnosis of endometriosis must be	
626 Breast, surgery of:		substantiated by laparoscopy.	
Following radical mastectomy:		7630 Malignant neoplasms of the breast	10
Both	180	Note: A rating of 100 percent shall con-	
One	150	tinue beyond the cessation of any sur-	
Following modified radical mastectomy:		gical, radiation, antineoplastic chemo-	
Both	160	therapy or other therapeutic procedure.	
One	140	Six months after discontinuance of such	
Following simple mastectomy or wide local	70	treatment, the appropriate disability rat-	
excision with significant alteration of size		ing shall be determined by mandatory	
or form:		VA examination. Any change in evalua-	
Both	150	tion based upon that or any subsequent	
One	130	examination shall be subject to the provi-	
Following wide local excision without sig-	.30	sions of §3.105(e) of this chapter. Rate	
nificant alteration of size or form:		chronic residuals according to impair-	
Both or one	0	ment of function due to scars,	
Note: For VA purposes:	U	lymphedema, or disfigurement (e.g., limi-	
(1) Radical mastectomy means		tation of arm, shoulder, and wrist motion,	
` ' '		or loss of grip strength, or loss of sensa-	
removal of the entire breast, un-		tion, or residuals from harvesting of mus-	
derlying pectoral muscles, and		cles for reconstructive purposes), and/or	
regional lymph nodes up to the		under diagnostic code 7626	
coracoclavicular ligament		7631 Benign neoplasms of the breast and other	
(2) Modified radical mastectomy		injuries of the breast. Rate chronic residuals ac-	
means removal of the entire		cording to impairment of function due to scars,	
breast and axillary lymph nodes		lymphedema, or disfigurement (e.g., limitation of	
(in continuity with the breast).		arm, shoulder, and wrist motion, or loss of grip	
Pectoral muscles are left intact		strength, or loss of sensation, or residuals from	
(3) Simple (or total) mastectomy		harvesting of muscles for reconstructive pur-	
means removal of all of the		poses), and/or under diagnostic code 7626	
breast tissue, nipple, and a		7632 Female sexual arousal disorder (FSAD)	
small portion of the overlying		1002 I emale sexual alousal disolder (FSAD)	
skin, but lymph nodes and mus-		¹ Review for entitlement to special monthly comp	oncati

(Authority: 38 U.S.C. 1155)

[60 FR 19855, Apr. 21, 1995, as amended at 67 FR 6874, Feb. 14, 2002; 67 FR 37695, May 30, 2002; 83 FR 15071, Apr. 9, 2018]

THE HEMATOLOGIC AND LYMPHATIC Systems

§4.117 Schedule of ratings—hemic and lymphatic systems.

	Rating
7702 Agranulocytosis, acquired: Requiring bone marrow transplant; or infections recurring, on average, at least once every six weeks per 12-month period.	100
riod	100
500/microliter (μl) but less than 1000/μl; or infections recurring, on average, at least once every three months per 12-month period	60
Requiring intermittent myeloid growth factors to maintain ANC greater than 1000/ μl; or infections recurring, on average, at least once per 12-month period but less than once every three months per 12-	
month period	30
Requiring continuous medication (e.g., antibiotics) for control; or requiring intermitent use of a myeloid growth factor to maintain ANC greater than or equal to	10
1500/µl Note: A 100 percent evaluation for bone marrow transplant shall be assigned as of the date of hospital admission and shall continue with a mandatory VA examination six months following hospital discharge. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.	10
7703 Leukemia (except for chronic myelogenous leukemia):	
When there is active disease or during a treatment phase	100
monoclonal B-cell lymphocytosis (MBL), asymptomatic, Rai Stage 0	0
Note (1): A 100 percent evaluation shall continue	

ote (1): A 100 percent evaluation shall continue beyond the cessation of any surgical therapy, radiation therapy, antineoplastic chemotherapy, or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no recurrence, rate on residuals

Note (2): Evaluate symptomatic chronic lymphocytic leu-kemia that is at Rai Stage I, II, III, or IV the same as any other leukemia evaluated under this diagnostic code

Note (3): Evaluate residuals of leukemia or leukemia therapy the appropriate diagnostic code(s) Myeloproliferative Disorders: (Diagnostic Codes 7704, 7718, 7719)

		Rating
7704	Polycythemia vera:	
	Requiring peripheral blood or bone marrow stem-cell transplant or chemotherapy (in- cluding myelosuppressants) for the pur-	
	pose of ameliorating the symptom bur- den	100
	Requiring phlebotomy 6 or more times per 12-month period or molecularly targeted therapy for the purpose of controlling	
	RBC count	60
	Requiring phlebotomy 4–5 times per 12- month period, or if requiring continuous biologic therapy or myelosuppressive agents, to include interferon, to maintain platelets <200,000 or white blood cells	
	(WBC) <12,000	30
	Requiring phlebotomy 3 or fewer times per 12-month period or if requiring biologic therapy or interferon on an intermittent basis as needed to maintain all blood	
	values at reference range levels	10
	Note (1): Rate complications such as hyper	
	gout, stroke, or thrombotic disease separa	telv

Note (2): If the condition undergoes leukemic transformation, evaluate as leukemia under diagnostic code 7703

Note (3): A 100 percent evaluation shall be assigned as of the date of hospital admission for peripheral blood or bone marrow stem cell transplant; or during the period of treatment with chemotherapy (including myelosuppressants). Six months following hospital discharge or, in the case of chemotherapy treatment, six months after completion of treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any reduction in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this

7705 Immune thrombocytopenia:

Requiring chemotherapy for chronic refractory thrombocytopenia; or a platelet	4000
count 30,000 or below despite treatment	100
Requiring immunosuppressive therapy; or	
for a platelet count higher than 30,000	
but not higher than 50,000, with history	
of hospitalization because of severe	
bleeding requiring intravenous immune	
globulin, high-dose parenteral	
corticosteroids, and platelet transfusions	70
Platelet count higher than 30,000 but not	
higher than 50,000, with either immune	
thrombocytopenia or mild mucous mem-	
brane bleeding which requires oral	
corticosteroid therapy or intravenous im-	
mune globulin	30
Platelet count higher than 30,000 but not	
higher than 50,000, not requiring treat-	
ment	10
Platelet count above 50,000 and asymp-	
tomatic; or for immune thrombocytopenia	
in remission	l c

Note (1): Separately evaluate splenectomy under diagnostic code 7706 and combine with an eval-uation under this diagnostic code

Note (2): A 100 percent evaluation shall continue beyond the cessation of chemotherapy. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any reduc-tion in evaluation based upon that or any subse-quent examination shall be subject to the provisions of §3.105(e) of this chapter

7706 Splenectomy ..

		Rating		Rating
	Note: Separately rate complications such		When there is active disease, during treat-	
	temic infections with encapsulated bacteria Note: Separately rate complications such a	1	ment phase, or with indolent and non- contiguous phase of low grade NHL	100
	temic infections with encapsulated bacteria		Note: A 100 percent evaluation shall continu	
7707 7709	Spleen, injury of, healed. Rate for any residuals. Hodgkin's lymphoma: With active disease or during a treatment		yond the cessation of any surgical therag ation therapy, antineoplastic chemother other therapeutic procedures. Two years a continuance of such treatment, the app	apy, or ifter dis- ropriate
	phase	100	disability rating shall be determined by ma VA examination. Any reduction in ev- based upon that or any subsequent exar shall be subject to the provisions of §3.10 this chapter. If there has been no recurren on residuals under the appropriate dia code(s)	aluation nination 05(e) of ce, rate
	discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any reduction in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals under the appropriate diagnostic code(s).		7716 Aplastic anemia: Requiring peripheral blood or bone marrow stem cell transplant; or requiring transfusion of platelets or red cells, on average, at least once every six weeks per 12-month period; or infections recurring, on average, at least once every six weeks per 12-month period	100
	Adenitis, tuberculous, active or inactive: Rate under § 4.88c or 4.89 of this part, whichever is appropriate. Multiple myeloma:		cells, on average, at least once every three months per 12-month period; or in- fections recurring, on average, at least once every three months per 12-month	
	Symptomatic multiple myeloma		period; or using continuous therapy with immunosuppressive agent or newer platelet stimulating factors	60
	, ,	n Soci- national	riod	nt shall mission xamina- ge. Any
	for five years after the diagnosis of symp multiple myeloma, at which time the app disability evaluation shall be determined to datory VA examination. Any reduction in tion based upon that or any subsequent extion shall be subject to the provisi §3.105(e) and §3.344 (a) and (b) of this c	ropriate by man- evalua- xamina- ons of	change in evaluation based upon that or a sequent examination shall be subject to the sions of §3.105(e) of this chapter Note (2): The term "newer platelet stimulat tors" includes medication, factors, or other approved by the United States Food an Administration	e provi- ing fac- agents
7714	Sickle cell anemia: With at least 4 or more painful episodes per 12-month period, occurring in skin, joints, bones, or any major organs, caused by hemolysis and sickling of red blood cells, with anemia, thrombosis, and infarction, with residual symptoms		7717 AL amyloidosis (primary amyloidosis) 7718 Essential thrombocythemia and primary myelofibrosis: Requiring either continuous myelosuppressive therapy or, for six months following hospital admission, pe- ripheral blood or bone marrow stem cell	100
	precluding even light manual labor With 3 painful episodes per 12-month period or with symptoms precluding other	100	transplant, or chemotherapy, or interferon treatment	100
	than light manual labor	60 30	myelosuppressive therapy, or chemotherapy, or interferon treatment to maintain platelet count <500 × 10 °/L	70
	Asymptomatic, established case in remission, but with identifiable organ impairment		Requiring continuous or intermittent myelosuppressive therapy, or chemo- therapy, or interferon treatment to main- tain platelet count of 200,000–400,000,	
	rectly attributable pathological findings, is ratable disability. Cases of symptomatic significant rections and symptomatic significant rections are rectly attributable pathological findings, is		or white blood cell (WBC) count of 4,000-10,000	30
	trait will be forwarded to the Director, Contion Service, for consideration under § 3.32	npensa-	Asymptomatic	0
	of this chapter	- (>) (1)	Note (1): If the condition undergoes leukemi formation, evaluate as leukemia under dia	

		Rating			Rating
sigr peri plar che moi cas con ratii amii upo sub cha 7719 Chronic (chronic mye leukemia): Requi ster	(2): A 100 percent evaluation shall ned as of the date of hospital admis pheral blood or bone marrow stem or try, or during the period of treatme motherapy (including myelosuppressanths following hospital discharge or, or of chemotherapy treatment, six month pletion of treatment, the appropriate or gashall be determined by mandatory nation. Any reduction in evaluation that or any subsequent examination piect to the provisions of §3.105(e) pter myelogenous leukemia (CML) eloid leukemia or chronic granulocytic iring peripheral blood or bone marrow m cell transplant, or continuous elosuppressive or immunosuppressive rapy treatment	be assion for solution for solu		Requiring continuous treatment with Vitamin B ₁₂ injections, Vitamin B ₁₂ sublingual or high-dose oral tablets, or Vitamin B ₁₂ nasal spray or gel	10 nus ane- ussigned equiring urenteral andatory bital dis- therapy. that or bipect to . There- parately us ane- sing pe- a, or re-
	ring intermittent myelosuppressive	100	7723	Acquired hemolytic anemia:	1
ther with inte ent In ap	rapy, or molecularly targeted therapy of tyrosine kinase inhibitors, or refron treatment when not in apparremission	60	7720	Requiring a bone marrow transplant or continuous intravenous or immunosuppressive therapy (e.g., prednisone, Cytoxan, azathioprine, or rituximab) Requiring immunosuppressive medication	100
	ularly targeted therapy with tyrosine ase inhibitors	30		4 or more times per 12-month period Requiring at least 2 but less than 4	60
Note	(1): If the condition undergoes leukem	ic trans-		courses of immunosuppressive therapy	
	nation, evaluate as leukemia under di le 7703	agnostic		per 12-month period	30
	(2): A 100 percent evaluation shall	be as-		suppressive therapy per 12-month period	10
peri plar che mor cas con ratir ami upo	ned as of the date of hospital admisipheral blood or bone marrow stem cent; or during the period of treatmemotherapy (including myelosuppressanths following hospital discharge or, e of chemotherapy treatment, six monimpletion of treatment, the appropriate ong shall be determined by mandatory nation. Any reduction in evaluation in that or any subsequent examination ject to the provisions of §3.105 of this	ell trans- ent with ints). Six in the ths after disability VA ex- based shall be		Asymptomatic	date of for six andatory bital dis- ed upon be sub- chapter y under
	ficiency anemia:		7704	uation under diagnostic code 7723	ı
mor Requi leas 12-t trea Asym by c Note: to b iron	ring intravenous iron infusions 4 or re times per 12-month period	Evaluate s under	7724	Solitary plasmacytoma: Solitary plasmacytoma, when there is active disease or during a treatment phase Note (1): A 100 percent evaluation shall obeyond the cessation of any surgical ther diation therapy, antineoplastic chemothe other therapeutic procedures (in autologous stem cell transplantation). Six after discontinuance of such treatment, the priate disability rating shall be determ mandatory VA examination. Any change uation based upon that or any subsequer ination shall be subject to the provis	rapy, ra- rapy, or ncluding months e appro- ined by in eval- tt exam-
	cid deficiency:			§3.105(e) of this chapter. If there has bee	n no re-
dos Asym	ring continuous treatment with high- e oral supplementationptomatic or requiring treatment only	10		currence, rate residuals under the approp agnostic codes Note (2): Rate a solitary plasmacytoma that	has de-
	dietary modificationbus anemia and Vitamin B ₁₂ defi-	0		veloped into multiple myeloma as symp multiple myeloma	Jonalic
ciency anem For ii due sigr	iia: nitial diagnosis requiring transfusion to severe anemia, or if there are ns or symptoms related to central		7705	Note (3): Rate residuals of plasma cell d (e.g., thrombosis) and adverse effects of treatment (e.g., neuropathy) under the priate diagnostic codes	medical
enc	vous system impairment, such as ephalopathy, myelopathy, or severe ipheral neuropathy, requiring paren-		7725	Myelodysplastic syndromes: Requiring peripheral blood or bone marrow stem cell transplant; or requiring chemo-	
	Il B ₁₂ therapy	100		therapy	

	Rating
Requiring 4 or more blood or platelet trans- fusions per 12-month period; or infec- tions requiring hospitalization 3 or more times per 12-month period	60
Requiring at least 1 but no more than 3 blood or platelet transfusions per 12-month period; infections requiring hospitalization at least 1 but no more than 2 times per 12-month period; or requiring biologic therapy on an ongoing basis or erythropoiesis stimulating agent (ESA) for 12 weeks or less per 12-month period;	
riod	30
Note (1): If the condition progresses to le	ukemia,

Note (1): If the condition progresses to leukemia, evaluate as leukemia under diagnostic code 7703

Note (2): A 100 percent evaluation shall be as-

Note (2): A 100 percent evaluation shall be assigned as of the date of hospital admission for peripheral blood or bone marrow stem cell transplant, or during the period of treatment with chemotherapy, and shall continue with a mandatory VA examination six months following hospital discharge or, in the case of chemotherapy treatment, six months after completion of treatment. Any reduction in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no recurrence, residuals will be rated under the appropriate diagnostic codes

[60 FR 49227, Sept. 22, 1995, as amended at 77 FR 6467, Feb. 8, 2012; 79 FR 2100, Jan. 13, 2014; 83 FR 54254, Oct. 29, 2018; 83 FR 54881, Nov. 1, 2018]

THE SKIN

§ 4.118 Schedule of ratings—skin.

(a) For the purposes of this section, systemic therapy is treatment that is administered through any route (orally, injection, suppository, intranasally) other than the skin, and topical therapy is treatment that is administered through the skin.

(b) Two or more skin conditions may be combined in accordance with §4.25 only if separate areas of skin are involved. If two or more skin conditions involve the same area of skin, then only the highest evaluation shall be used.

	Rat- ing
7800 Burn scar(s) of the head, face, or neck; scar(s) of the head, face, or neck due to other causes; or other disfigurement of the head, face, or neck:	
With visible or palpable tissue loss and ei- ther gross distortion or asymmetry of three or more features or paired sets of features (nose, chin, forehead, eyes (in- cluding eyelids), ears (auricles), cheeks,	
lips), or; with six or more characteristics of disfigurement	80

	Rat- ing
With visible or palpable tissue loss and either gross distortion or asymmetry of two features or paired sets of features (nose, chin, forehead, eyes (including eyelids), ears (auricles), cheeks, lips), or; with four or five characteristics of disfigurement	50
With visible or palpable tissue loss and either gross distortion or asymmetry of one feature or paired set of features (nose, chin, forehead, eyes (including eyelids), ears (auricles), cheeks, lips), or; with two	
or three characteristics of disfigurement	30
With one characteristic of disfigurement Note (1):The 8 characteristics of disfigurement, for purposes of evaluation under §4.118, are: Scar 5 or more inches (13 or more cm.) in	10
length. Scar at least one-quarter inch (0.6 cm.) wide at widest part.	
Surface contour of scar elevated or depressed on palpation.	
Scar adherent to underlying tissue. Skin hypo-or hyper-pigmented in an area exceeding six square inches (39 sq. cm.).	
Skin texture abnormal (irregular, atrophic, shiny, scaly, etc.) in an area exceeding six square inches (39 sq. cm.).	
Underlying soft tissue missing in an area exceeding six square inches (39 sq. cm.).	
Skin indurated and inflexible in an area exceeding six square inches (39 sq. cm.). Note (2): Rate tissue loss of the auricle	
under DC 6207 (loss of auricle) and anatomical loss of the eye under DC 6061 (anatomical loss of both eyes) or DC 6063 (anatomical loss of one eye), as ap-	
propriate. Note (3): Take into consideration unretouched color photographs when	
evaluating under these criteria. Note (4): Separately evaluate disabling ef-	
fects other than disfigurement that are as- sociated with individual scar(s) of the head, face, or neck, such as pain, insta-	
bility, and residuals of associated muscle or nerve injury, under the appropriate di- agnostic code(s) and apply § 4.25 to com- bine the evaluation(s) with the evaluation	
assigned under this diagnostic code. Note (5): The characteristic(s) of disfigure-	
ment may be caused by one scar or by multiple scars; the characteristic(s) required to assign a particular evaluation need not be caused by a single scar in order to assign that evaluation.	
Burn scar(s) or scar(s) due to other causes, f the head, face, or neck, that are associated	
underlying soft tissue damage:	
Area or areas of 144 square inches (929 sq. cm.) or greater	40
inches (929 sq. cm.)	30
Area or areas of at least 12 square inches (77 sq. cm.) but less than 72 square inches (465 sq. cm.)	20
Area or areas of at least 6 square inches (39 sq. cm.) but less than 12 square	

(39 sq. cm.) but less than 12 square inches (77 sq. cm.)

10

7801 not

	Rat-		Rat-
	ing		ing
Note (1): For the purposes of DCs 7801 and 7802, the six (6) zones of the body are defined as each extremity, anterior trunk, and posterior trunk. The midaxillary line divides the anterior trunk from the posterior trunk. Note (2): A separate evaluation may be assigned for each affected zone of the body under this diagnostic code if there are multiple scars, or a single scar, affecting multiple zones of the body. Combine the separate evaluations under §4.25. Alternatively, if a higher evaluation would result from adding the areas affected from		Constant or near-constant systemic therapy including, but not limited to, corticosteroids, phototherapy, retinoids, biologics, photochemotherapy, psoralen with long-wave ultraviolet-A light (PUVA), or other immunosuppressive drugs required over the past 12-month period	60 30
multiple zones of the body, a single eval- uation may also be assigned under this		duration of 6 weeks or more, but not constantly, over the past 12-month period.	
diagnostic code. 7802 Burn scar(s) or scar(s) due to other causes, not of the head, face, or neck, that are not associated with underlying soft tissue damage: Area or areas of 144 square inches (929 sq.		At least one of the following	10
cm.) or greater	10	of exposed areas affected; or Intermittent systemic therapy including, but not limited to, corticosteroids,	
defined as each extremity, anterior trunk, and posterior trunk. The midaxillary line divides the anterior trunk from the posterior trunk.		phototherapy, retinoids, biologics, photochemotherapy, PUVA, or other im- munosuppressive drugs required for a total duration of less than 6 weeks over	
Note (2): A separate evaluation may be as- signed for each affected zone of the body under this diagnostic code if there are		the past 12-month period. No more than topical therapy required over the past 12-month period and at least one of the following	0
multiple scars, or a single scar, affecting multiple zones of the body. Combine the separate evaluations under §4.25. Alternatively, if a higher evaluation would result from adding the areas affected from multiple zones of the body, a single evaluation served to be secured to the second of the s		Characteristic lesions involving less than 5 percent of the entire body affected; or Characteristic lesions involving less than 5 percent of exposed areas affected. Or rate as disfigurement of the head, face, or neck (DC 7800) or scars (DCs 7801,	
uation may also be assigned under this diagnostic code.		7802, 7804, or 7805), depending upon the predominant disability. This rating instruc-	
7804 Scar(s), unstable or painful:.		tion does not apply to DC 7824.	
Five or more scars that are unstable or painful	30	7806 Dermatitis or eczema. Evaluate under the General Rating Formula	
Three or four scars that are unstable or painful	20	for the Skin. 7807 American (New World) leishmaniasis	
One or two scars that are unstable or pain-		(mucocutaneous, espundia):	
ful	10	Rate as disfigurement of the head, face, or neck (DC 7800), scars (DC's 7801, 7802, 7803, 7804, or 7805), or dermatitis (DC 7806), depending upon the predominant	
Note (2): If one or more scars are both un- stable and painful, add 10 percent to the evaluation that is based on the total num-		disability. Note: Evaluate non-cutaneous (visceral) leishmaniasis under DC 6301 (visceral	
ber of unstable or painful scars Note (3): Scars evaluated under diagnostic		leishmaniasis). 7808 Old World leishmaniasis (cutaneous, Oriental	
codes 7800, 7801, 7802, or 7805 may also receive an evaluation under this di- agnostic code, when applicable		sore): Rate as disfigurement of the head, face, or neck (DC 7800), scars (DC's, 7801, 7802,	
7805 Scars, other; and other effects of scars evaluated under diagnostic codes 7800, 7801, 7802, or 7804:		 7803, 7804, or 7805), or dermatitis (DC 7806), depending upon the predominant disability. Note: Evaluate non-cutaneous (visceral) 	
Evaluate any disabling effect(s) not considered in a rating provided under diagnostic codes 7800–04 under an appropriate diagnostic code.		leishmaniasis under DC 6301 (visceral leishmaniasis). 7809 Discoid lupus erythematosus. Evaluate under the General Rating Formula	
General Rating Formula For The Skin For DCs 7806, 7809, 7813–7816, 7820–7822, and 7824:		for the Skin. Note: Do not combine with ratings under DC	
At least one of the following	60	6350. 7811 Tuberculosis luposa (lupus vulgaris), active or	
40 percent of the entire body or more than 40 percent of exposed areas affected; or		inactive: Rate under §§ 4.88c or 4.89, whichever is appropriate.	

	Rat-		Rat-
	ing		ing
7813 Dermatophytosis (ringworm: Of body, tinea corporis; of head, tinea capitis; of feet, tinea pedis; of beard area, tinea barbae; of nails, tinea unguium (onychomycosis); of inguinal area (jock itch), tinea cruris; tinea versicolor). Evaluate under the General Rating Formula for the Skin.		Any extent of involvement of the skin, and no more than topical therapy required over the past 12-month period	0
7815 Bullous disorders (including pemphigus vulgaris, pemphigus foliaceous, bullous pemphigoid, dermatitis herpetiformis, epidermolysis bullosa acquisita, benign chronic familial pemphigus (Hailey-Hailey), and porphyria cutanea		scribed therapy, as documented by medical records. 7818 Malignant skin neoplasms (other than malignant melanoma): Rate as disfigurement of the head, face, or	
tarda). Evaluate under the General Rating Formula		neck (DC 7800), scars (DC's 7801, 7802, 7803, 7804, or 7805), or impairment of	
for the Skin. Note: Rate complications and residuals of		function. Note: If a skin malignancy requires therapy	
mucosal involvement (ocular, oral, gastro- intestinal, respiratory, or genitourinary) separately under the appropriate diag-		that is comparable to that used for systemic malignancies, <i>i.e.</i> , systemic chemotherapy, X-ray therapy more extensive	
nostic code. 7816 Psoriasis. Evaluate under the General Rating Formula		than to the skin, or surgery more exten- sive than wide local excision, a 100-per- cent evaluation will be assigned from the	
for the Skin. Note: Rate complications such as psoriatic arthritis and other clinical manifestations		date of onset of treatment, and will con- tinue, with a mandatory VA examination six months following the completion of	
(e.g., oral mucosa, nails) separately under the appropriate diagnostic code.		such antineoplastic treatment, and any change in evaluation based upon that or	
7817 Erythroderma: Generalized involvement of the skin with		any subsequent examination will be subject to the provisions of §3.105(e) of this	
systemic manifestations (such as fever, weight loss, or hypoproteinemia) AND one of the following	100	chapter. If there has been no local recur- rence or metastasis, evaluation will then be made on residuals. If treatment is con-	
Constant or near-constant systemic therapy such as therapeutic doses of	100	fined to the skin, the provisions for a 100- percent evaluation do not apply.	
corticosteroids, other immunosuppressive drugs, retinoids, PUVA (psoralen with		7819 Benign skin neoplasms: Rate as disfigurement of the head, face, or	
long-wave ultraviolet-A light), UVB (ultra- violet-B light) treatments, biologics, or electron beam therapy required over the		neck (DC 7800), scars (DC's 7801, 7802, 7803, 7804, or 7805), or impairment of function.	
past 12 month period; or No current treatment due to a documented		7820 Infections of the skin not listed elsewhere (including bacterial, fungal, viral, treponemal, and	
history of treatment failure with 2 or more treatment regimens	100	parasitic diseases). Evaluate under the General Rating Formula for the Skin.	
systemic manifestations and one of the following.		7821 Cutaneous manifestations of collagen-vas- cular diseases not listed elsewhere (including	
Constant or near-constant systemic therapy such as therapeutic doses of corticosteroids, other immunosuppressive		scleroderma, calcinosis cutis, subacute cutaneous lupus erythematosus, and dermatomyositis). Evaluate under the General Rating Formula	
drugs, retinoids, PUVA, UVB treatments, biologics, or electron beam therapy re- quired over the past 12-month period; or		for the Skin. 7822 Papulosquamous disorders not listed else-	
No current treatment due to a documented history of treatment failure with 1 treat-	60	where (including lichen planus, large or small plaque parapsoriasis, pityriasis lichenoides et varioliformis acuta (PLEVA), lymphomatoid	
ment regimen Any extent of involvement of the skin, and any of the following therapies required for a total duration of 6 weeks or more, but	80	papulosus, mycosis fungoides, and pityriasis rubra pilaris (PRP)). Evaluate under the General Rating Formula	
not constantly, over the past 12-month period: systemic therapy such as thera- peutic doses of corticosteroids, other im-		for the Skin. 7823 Vitiligo: With exposed areas affected	10
munosuppressive drugs, retinoids, PUVA, UVB treatments, biologics, or electron	00	With no exposed areas affected	0
Any extent of involvement of the skin, and any of the following therapies required for	30	keratoderma). Evaluate under the General Rating Formula	
a total duration of less than 6 weeks over the past 12-month period: systemic ther-		for the Skin. 7825 Chronic urticaria:	
apy such as therapeutic doses of corticosteroids, other immunosuppressive drugs, retinoids, PUVA, UVB treatments,		For the purposes of this diagnostic code, chronic urticaria is defined as continuous urticaria at least twice per week, off treat-	
biologics, or electron beam therapy	10	ment, for a period of six weeks or more.	

	Dat		Det
	Rat- ing		Rat- ing
Chronic refractory urticaria that requires third line treatment for control (e.g., plasmapheresis, immunotherapy, immunosuppressives) due to ineffective-		Deep acne (deep inflamed nodules and pus- filled cysts) affecting less than 40 percent of the face and neck, or deep acne other than on the face and neck	10
ness with first and second line treatments	60	Superficial acne (comedones, papules, pustules) of	
Chronic urticaria that requires second line		any extent	0
treatment (e.g., corticosteroids, sympathomimetics, leukotriene inhibitors, neutrophil inhibitors, thyroid hormone) for		Or rate as disfigurement of the head, face, or neck (DC 7800) or scars (DCs 7801, 7802, 7804, or 7805), depending upon the	
control Chronic urticaria that requires first line treat-	30	predominant disability.	
ment (antihistamines) for control	10	7829 Chloracne:	
7826 Vasculitis, primary cutaneous:		Deep acne (deep inflamed nodules and pus- filled cysts) affecting 40 percent or more	
Persistent documented vasculitis episodes		of the face and neck	30
refractory to continuous immuno-	60	Deep acne (deep inflamed nodules and pus-	
suppressive therapy All of the following	30	filled cysts) affecting the intertriginous	
Recurrent documented vasculitic episodes		areas (the axilla of the arm, the	
occurring four or more times over the past		anogenital region, skin folds of the	00
12-month period; and		breasts, or between digits)	20
Requiring intermittent systemic immuno-	20	Deep acne (deep inflamed nodules and pus- filled cysts) affecting less than 40 percent	
suppressive therapy for control At least one of the following	30 10	of the face and neck; or deep acne affect-	
Recurrent documented vasculitic episodes	10	ing non-intertriginous areas of the body	
occurring one to three times over the past		(other than the face and neck)	10
12-month period, and requiring intermit-		Superficial acne (comedones, papules,	
tent systemic immunosuppressive therapy		pustules) of any extent	0
for control; or Without recurrent documented vasculitic epi-		Or rate as disfigurement of the head, face,	
sodes but requiring continuous systemic		or neck (DC 7800) or scars (DCs 7801, 7802, 7804, or 7805), depending upon the	
medication for control.		predominant disability.	
Or rate as disfigurement of the head, face,		7830 Scarring alopecia:	
or neck (DC 7800) or scars (DCs 7801,		Affecting more than 40 percent of the scalp	20
7802, 7804, or 7805), depending upon the predominant disability.		Affecting 20 to 40 percent of the scalp	10
7827 Erythema multiforme; Toxic epidermal		Affecting less than 20 percent of the scalp	0
necrolysis:		7831 Alopecia areata:	
Recurrent mucosal, palmar, or plantar in-		With loss of all body hair	10
volvement impairing mastication, use of		With loss of hair limited to scalp and face	0
hands, or ambulation occurring four or more times over the past 12-month period		7832 Hyperhidrosis:	
despite ongoing immunosuppressive ther-		Unable to handle paper or tools because of moisture, and unresponsive to therapy	30
ару	60	Able to handle paper or tools after therapy	0
All of the following	30	7833 Malignant melanoma:	
Recurrent mucosal, palmar, or plantar in-		Rate as scars (DC's 7801, 7802, 7803,	
volvement not impairing mastication, use of hands, or ambulation, occurring four or		7804, or 7805), disfigurement of the head,	
more times over the past 12-month pe-		face, or neck (DC 7800), or impairment of	
riod; andrequiring intermittent systemic		function (under the appropriate body system).	
therapy.		Note: If a skin malignancy requires therapy	
At least one of the following One to three episodes of mucosal, palmar,	10	that is comparable to that used for sys-	
or plantar involvement not impairing mas-		temic malignancies, i.e., systemic chemo-	
tication, use of hands, or ambulation, oc-		therapy, X-ray therapy more extensive	
curring over the past 12-month period		than to the skin, or surgery more exten- sive than wide local excision, a 100-per-	
AND requiring intermittent systemic ther-		cent evaluation will be assigned from the	
apy; or Without recurrent episodes, but requiring		date of onset of treatment, and will con-	
continuous systemic medication for con-		tinue, with a mandatory VA examination	
trol.		six months following the completion of	
Or rate as disfigurement of the head, face,		such antineoplastic treatment, and any change in evaluation based upon that or	
or neck (DC 7800) or scars (DCs 7801,		any subsequent examination will be sub-	
7802, 7804, or 7805), depending upon the predominant disability.		ject to the provisions of § 3.105(e). If there	
Note: For the purposes of this DC only, sys-		has been no local recurrence or metas-	
temic therapy may consist of one or more		tasis, evaluation will then be made on re-	
of the following treatment agents:		siduals. If treatment is confined to the skin, the provisions for a 100-percent	
immunosuppressives, antihistamines, or		evaluation do not apply.	
sympathomimetics. 7828 Acne:			
Deep acne (deep inflamed nodules and pus-			
filled cysts) affecting 40 percent or more			
of the face and neck	30		

(Authority: 38 U.S.C. 1155)

[67 FR 49596, July 31, 2002; 67 FR 58448, 58449, Sept. 16, 2002; 73 FR 54710, Oct. 23, 2008; 77 FR 2910, Jan. 20, 2012; 83 FR 32597, July 13, 2018; 83 FR 38663, Aug. 7, 2018]

[67 FR 49596, July 31, 2002; 67 FR 58448, 5	58449,		ing
Sept. 16, 2002; 73 FR 54710, Oct. 23, 2008; 7 2910, Jan. 20, 2012; 83 FR 32597, July 13, 83 FR 38663, Aug. 7, 2018] THE ENDOCRINE SYSTEM	7 FR	Note (2): This evaluation shall continue for six months after initial diagnosis. Thereafter, rate residuals of disease or medical treatment under the most appropriate diagnostic code(s) under the appropriate body system (e.g., eye, digestics and the state of the system of the syste	
THE ENDOCRINE SISTEM		tive, mental disorders). Note (3): If eye involvement, such as	
§4.119 Schedule of ratings—endoc system.	rine	exophthalmos, corneal ulcer, blurred vision, or diplopia, is also present due to thyroid disease, also separately evaluate under the appropriate	
	Rat- ing	diagnostic code(s) in §4.79, Schedule of Ratings—Eye (such as diplopia (DC 6090) or impairment of central visual acuity (DCs 6061–6066)).	
7900 Hyperthyroidism, including, but not limited to, Graves' disease:		7904 Hyperparathyroidism:	
For six months after initial diagnosis	30	For six months from date of discharge following	100
Thereafter, rate residuals of disease or complica- tions of medical treatment within the appro- priate diagnostic code(s) within the appropriate body system. Note (1): If hyperthyroid cardiovascular or cardiac		surgery	100
disease is present, separately evaluate under DC 7008 (hyperthyroid heart disease). Note (2): Separately evaluate eye involvement occurring as a manifestation of Graves' Disease as diplopia (DC 6090); impairment of central visual acuity (DCs 6061–6066); or under the		Hypercalcemia (indicated by at least one of the following: Total Ca greater than 12 mg/dL (3–3.5 mmol/L), lonized Ca greater than 5.6 mg/dL (2–2.5 mmol/L), creatinine clearance less than 60 mL/min, bone mineral density T-score less than 2.5 SD (below mean) at any site or previous fragility fracture)	60
most appropriate DCs in § 4.79. 7901 Thyroid enlargement, toxic:		Note (2): Where surgical intervention is indicated,	"
Note (1): Evaluate symptoms of hyperthyroidism under DC 7900, hyperthyroidism, including, but not limited to, Graves' disease.		this evaluation shall continue until the day of surgery, at which time the provisions pertaining to a 100-percent evaluation shall apply.	
Note (2): If disfigurement of the neck is present due to thyroid disease or enlargement, separately evaluate under DC 7800 (burn scar(s) of the head, face, or neck causes; or other disfigurement of the head, face, or neck).		Note (3): Where surgical intervention is not indi- cated, this evaluation shall continue for six months after pharmacologic treatment begins. After six months, rate on residuals under the appropriate diagnostic code(s) within the appro- priate body system(s) based on a VA examina- tion.	
7902 Thyroid enlargement, nontoxic:		Symptoms such as fatigue, anorexia, nausea, or	
Note (1): Evaluate symptoms due to pressure on adjacent organs (such as the trachea, larynx, or esophagus) under the appropriate diagnostic code(s) within the appropriate body system.		constipation that occur despite surgery; or in in- dividuals who are not candidates for surgery but require continuous medication for control	10
Note (2): If disfigurement of the neck is present		Asymptomatic	0
due to thyroid disease or enlargement, sepa- rately evaluate under DC 7800 (burn scar(s) of the head, face, or neck; scar(s) of the head,		evaluate chronic residuals, such as nephrolithiasis (kidney stones), decreased renal function, fractures, vision problems, and cardio-	
face, or neck due to other causes; or other dis- figurement of the head, face, or neck). 7903 Hypothyroidism:		vascular complications, under the appropriate diagnostic codes.	
Hypothyroidism manifesting as myxedema (cold		7905 Hypoparathyroidism: For three months after initial diagnosis	100
intolerance, muscular weakness, cardiovascular involvement (including, but not limited to hypotension, bradycardia, and pericardial effusion), and mental disturbance (including, but not limited to dementia, slowing of thought and de-		Thereafter, evaluate chronic residuals, such as nephrolithiasis (kidney stones), cataracts, de- creased renal function, and congestive heart failure under the appropriate diagnostic codes.	100
pression))	100	7906 Thyroiditis: With normal thyroid function (euthyroid)	0
Note (1): This evaluation shall continue for six months beyond the date that an examining physician has determined crisis stabilization. Thereafter, the residual effects of hypothyroidism shall be rated under the appropriate diagnostic code(s) within the appropriate body system(s)		Note: Manifesting as hyperthyroidism, evaluate as hyperthyroidism, including, but not limited to, Graves' disease (DC 7900); manifesting as hypothyroidism, evaluate as hypothyroidism (DC 7903).	
(e.g., eye, digestive, and mental disorders).	20	7907 Cushing's syndrome: As active, progressive disease, including areas of	
Hypothyroidism without myxedema	30	osteoporosis, hypertension, and proximal upper and lower extremity muscle wasting that results in inability to rise from squatting position, climb stairs, rise from a deep chair without assist-	400
		ance, or raise arms	100

			_
	Rat- ing		Rat- ing
Proximal upper or lower extremity muscle wasting that results in inability to rise from squatting position, climb stairs, rise from a deep chair without assistance, or raise arms		Requiring more than one daily injection of insulin, restricted diet, and regulation of activities (avoidance of strenuous occupational and recreational activities) with episodes of ketoacidosis or hypoglycemic reactions requiring at least three hospitalizations per year or weekly visits to a diabetic care provider, plus either progressive loss of weight and strength or complications that would be compensable if separately evaluated	
vascular collapse. Note (3): Tuberculous Addison's disease will be evaluated as active or inactive tuberculosis. If inactive, these evaluations are not to be combined with the graduated ratings of 50 percent or 30 percent for non-pulmonary tuberculosis specified under §4.88b. Assign the higher rating. 7912 Polyglandular syndrome (multiple endocrine		7917 Hyperaldosteronism (benign or malignant): Note: Evaluate as malignant or benign neoplasm, as appropriate. 7918 Pheochromocytoma (benign or malignant): Note: Evaluate as malignant or benign neoplasm as appropriate. 7919 C-cell hyperplasia of the thyroid: If antineoplastic therapy is required, evaluate as a	
neoplasia, autoimmune polyglandular syndrome): Evaluate according to major manifestations to include, but not limited to, Type I diabetes mellitus, hyperthyroidism, hypothyroidism, hypoparathyroidism, or Addison's disease. 7913 Diabetes mellitus:		mailignant neoplasm under DC 7914. If a prophylactic thyroidectomy is performed (based upon genetic testing) and antineoplastic therapy is not required, evaluate as hypothyroidism under DC 7903.	

[61 FR 20446, May 7, 1996, as amended at 82 FR 50804, Nov. 2, 2017]

NEUROLOGICAL CONDITIONS AND CONVULSIVE DISORDERS

§ 4.120 Evaluations by comparison.

Disability in this field is ordinarily to be rated in proportion to the impairment of motor, sensory or mental function. Consider especially psychotic manifestations, complete or partial loss of use of one or more extremities. speech disturbances, impairment of vision, disturbances of gait, tremors, visceral manifestations, injury to the skull, etc. In rating disability from the conditions in the preceding sentence refer to the appropriate schedule. In rating peripheral nerve injuries and their residuals, attention should be given to the site and character of the injury, the relative impairment in motor function, trophic changes, or sensory disturbances.

§ 4.121 Identification of epilepsy.

When there is doubt as to the true nature of epileptiform attacks, neurological observation in a hospital adequate to make such a study is necessary. To warrant a rating for epilepsy, the seizures must be witnessed or verified at some time by a physician. As to frequency, competent, consistent lay testimony emphasizing convulsive and immediate post-convulsive characteristics may be accepted. The frequency of seizures should be ascertained under the ordinary conditions of life (while not hospitalized).

§ 4.122 Psychomotor epilepsy.

The term psychomotor epilepsy refers to a condition that is characterized by seizures and not uncommonly by a chronic psychiatric disturbance as well.

(a) Psychomotor seizures consist of episodic alterations in conscious control that may be associated with automatic states, generalized convulsions, random motor movements (chewing, lip smacking, fumbling), hallucinatory phenomena (involving taste, smell, sound, vision), perceptual illusions (deja vu, feelings of loneliness, strangeness, macropsia, micropsia, dreamy states), alterations in thinking (not open to reason), alterations in memory, abnormalities of mood or affect (fear, alarm, terror, anger, dread, well-

being), and autonomic disturbances (sweating, pallor, flushing of the face, visceral phenomena such as nausea, vomiting, defecation, a rising feeling of warmth in the abdomen). Automatic states or automatisms are characterized by episodes of irrational, irrelevant, disjointed, unconventional, asocial, purposeless though seemingly coordinated and purposeful, confused or inappropriate activity of one to several minutes (or, infrequently, hours) duration with subsequent amnesia for the seizure. Examples: A person of high social standing remained seated, muttered angrily, and rubbed the arms of his chair while the National Anthem was being played; an apparently normal person suddenly disrobed in public; a man traded an expensive automobile for an antiquated automobile in poor mechanical condition and after regaining conscious control, discovered that he had signed an agreement to pay an additional sum of money in the trade. The seizure manifestations of psychomotor epilepsy vary from patient to patient and in the same patient from seizure to seizure.

(b) A chronic mental disorder is not uncommon as an interseizure manifestation of psychomotor epilepsy and may include psychiatric disturbances extending from minimal anxiety to severe personality disorder (as distinguished from developmental) or almost complete personality disintegration (psychosis). The manifestations of a chronic mental disorder associated with psychomotor epilepsy, like those of the seizures, are protean in character

§ 4.123 Neuritis, cranial or peripheral.

Neuritis, cranial or peripheral, characterized by loss of reflexes, muscle atrophy, sensory disturbances, and constant pain, at times excruciating, is to be rated on the scale provided for injury of the nerve involved, with a maximum equal to severe, incomplete, paralysis. See nerve involved for diagnostic code number and rating. The maximum rating which may be assigned for neuritis not characterized by organic changes referred to in this section will be that for moderate, or with sciatic nerve involvement, for moderately severe, incomplete paralysis.

ing

100

§4.124

§ 4.124 Neuralgia, cranial or peripheral.

Neuralgia, cranial or peripheral, characterized usually by a dull and intermittent pain, of typical distribution so as to identify the nerve, is to be rated on the same scale, with a maximum equal to moderate incomplete paralysis. See nerve involved for diagnostic code number and rating. Tic douloureux, or trifacial neuralgia, may be rated up to complete paralysis of the affected nerve.

§4.124a Schedule of ratings—neurological conditions and convulsive disorders.

[With the exceptions noted, disability from the following diseases and their residuals may be rated from 10 percent to 100 percent in proportion to the impairment of motor, sensory, or mental function. Consider especially psychotic manifestations, complete or partial loss of use of one or more extremities, speech disturbances, impairment of vision, disturbances of gait, tremors, visceral manifestations, etc., referring to the appropriate bodily system of the schedule. With partial loss of use of one or more extremities from neurological lesions, rate by comparison with the mild, moderate, severe, or complete paralysis of peripheral nerves]

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM

	Rat- ing
8000 Encephalitis, epidemic, chronic:	
As active febrile disease	100
Rate residuals, minimum	10
Brain, new growth of:	400
8002 Malignant	100
for 2 years following cessation of surgical,	
chemotherapeutic or other treatment modality.	
At this point, if the residuals have stabilized,	
the rating will be made on neurological residu-	
als according to symptomatology.	
Minimum rating	30
8003 Benign, minimum	60
Rate residuals, minimum	10
8004 Paralysis agitans:	
Minimum rating	30
8005 Bulbar palsy	100
8007 Brain, vessels, embolism of. 8008 Brain, vessels, thrombosis of.	
8009 Brain, vessels, hemorrhage from:	
Rate the vascular conditions under Codes 8007	
through 8009, for 6 months	100
Rate residuals, thereafter, minimum	10
8010 Myelitis:	
Minimum rating	10
8011 Poliomyelitis, anterior:	
As active febrile disease	100
Rate residuals, minimum	10

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM—Continued

8012 Hematomyelia:

For 6 months

Rate residuals, minimum	10
8013 Syphilis, cerebrospinal.	
8014 Syphilis, meningovascular.	
8015 Tabes dorsalis.	
NOTE: Rate upon the severity of convulsions, pa- ralysis, visual impairment or psychotic involve- ment, etc.	
8017 Amyotrophic lateral sclerosis	100
NOTE: Consider the need for special monthly	100
compensation.	
8018 Multiple sclerosis:	
Minimum rating	30
8019 Meningitis, cerebrospinal, epidemic:	
As active febrile disease	100
Rate residuals, minimum	10
8020 Brain, abscess of:	
As active disease	100
Rate residuals, minimum	10
Spinal cord, new growths of:. 8021 Malignant	100
8021 Malignant	100
for 2 years following cessation of surgical,	
chemotherapeutic or other treatment modality.	
At this point, if the residuals have stabilized,	
the rating will be made on neurological residu-	
als according to symptomatology.	
Minimum rating	30
8022 Benign, minimum rating	60
Rate residuals, minimum	10
8023 Progressive muscular atrophy: Minimum rating	30
8024 Syringomyelia:	00
Minimum rating	30
8025 Myasthenia gravis:	00
Minimum rating	30
NOTE: It is required for the minimum ratings for	
residuals under diagnostic codes 8000-8025,	
that there be ascertainable residuals. Deter-	
minations as to the presence of residuals not	
capable of objective verification, i.e., head-	
aches, dizziness, fatigability, must be approached on the basis of the diagnosis re-	
proached on the basis of the diagnosis re-	
corded; subjective residuals will be accepted	
when consistent with the disease and not more likely attributable to other disease or no	
disease. It is of exceptional importance that	
when ratings in excess of the prescribed min-	
imum ratings are assigned, the diagnostic	
codes utilized as bases of evaluation be cited,	
in addition to the codes identifying the diag-	
noses.	
8045 Residuals of traumatic brain injury (TBI):	
There are three main areas of dysfunction	
that may result from TBI and have pro-	
found effects on functioning: cognitive	
(which is common in varying degrees	
after TBI), emotional/behavioral, and physical. Each of these areas of dysfunc-	
tion may require evaluation	
non may require evaluation	

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM—Continued

ing

Subjective symptoms may be the only residual of TBI or may be associated with cog-nitive impairment or other areas of dysfunction. Evaluate subjective symptoms that are residuals of TBI, whether or not they are part of cognitive impairment, under the subjective symptoms facet in the table titled "Evaluation of Cognitive Impairment and Other Residuals of TBI Not Otherwise Classified." However, separately evaluate any residual with a distinct diagnosis that may be evaluated under another diagnostic code, such as migraine headache or Meniere's disease. even if that diagnosis is based on subjective symptoms, rather than under the "Evaluation of Cognitive Impairment and Other Residuals of TBI Not Otherwise Classified" table.

Evaluate emotional/behavioral dysfunction under §4.130 (Schedule of ratings—mental disorders) when there is a diagnosis of a mental disorder. When there is no diagnosis of a mental disorder, evaluate emotional/behavioral symptoms under the criteria in the table titled "Evaluation of Cognitive Impairment and Other Residuals of TBI Not Otherwise Classified."

Evaluate physical (including neurological) dysfunction based on the following list, under an appropriate diagnostic code: Motor and sensory dysfunction, including pain, of the extremities and face; visual impairment; hearing loss and tinnitus; loss of sense of smell and taste; seizures; gait, coordination, and balance problems; speech and other communication difficulties, including aphasia and related disorders, and dysarthria; neurogenic bladder; neurogenic bowel; cranial nerve dysfunctions; autonomic nerve dysfunctions; and endocrine dysfunctions.

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM—Continued

The preceding list of types of physical dysfunction does not encompass all possible residuals of TBI. For residuals not listed here that are reported on an examination, evaluate under the most appropriate diagnostic code. Evaluate each condition separately, as long as the same signs and symptoms are not used to support more than one evaluation, and combine under § 4.25 the evaluations for each separately rated condition. The evaluation assigned based on the "Evaluation of Cognitive Impairment and Other Residuals of TBI Not Otherwise Classified" table will be considered the evaluation for a single condition for purposes of combining with other disability evaluations.

Consider the need for special monthly compensation for such problems as loss of use of an extremity, certain sensory impairments, erectile dysfunction, the need for aid and attendance (including for protection from hazards or dangers incident to the daily environment due to cognitive impairment), being housebound, etc.

Evaluation of Cognitive Impairment and Subjective Symptoms

The table titled "Evaluation of Cognitive Impairment and Other Residuals of TBI Not Otherwise Classi fied" contains 10 important facets of TBI related to cognitive impairment and subjective symptoms. It provides criteria for levels of impairment for each facet, as appropriate, ranging from 0 to 3, and a 5th level, the highest level of impairment, labeled "total." However, not every facet has every level of severity. The Consciousness facet, for example, does not provide for an impairment level other than "total," since any level of impaired consciousness would be totally disabling. Assign a 100-percent evaluation if "total" is the level of evaluation for one or more facets. If no facet is evaluated as "total," assign the overall percentage evaluation based on the level of the highest facet as follows: 0 = 0 percent; 1 = 10 percent; 2 = 40 percent; and 3 = 70 percent. For example, assign a 70 percent evaluation if 3 is the highest level of evaluation for any facet.

Note (1): There may be an overlap of manifestations of conditions evaluated under the table titled "Evaluation Of Cognitive Impairment And Other Residuals Of TBI Not Otherwise Classified" with manifestations of a comorbid mental or neurologic or other physical disorder that can be separately evaluated under another diagnostic code. In such cases, do not assign more than one evaluation based on the same manifestations. If the manifestations of two or more conditions cannot be clearly separated, assign a single evaluation under whichever set of diagnostic criteria allows the better assessment of overall impaired functioning due to both conditions. However, if the manifestations are clearly separable, assign a separate evaluation for each condition

38 CFR Ch. I (7-1-21 Edition)

ORGANIC DISEASES OF THE CENTRAL NERVOUS

SYSTEM—Continued

§4.124a

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM—Continued

	Rat- ing		Rat- ing
Note (2): Symptoms listed as examples at certain evaluation levels in the table are only examples and are not symptoms that must be present in order to assign a particular evaluation Note (3): "Instrumental activities of daily living" refers to activities other than self-care that are needed for independent living, such as meal preparation, doing housework and other chores, shopping, traveling, doing laundry, being responsible for one's own medications, and using a telephone. These activities are distinguished from "Activities of daily living," which refers to basic self-care and includes bathing or showering, dressing, eating, getting in or out of bed or a chair, and using the toilet. Note (4): The terms "mild," "moderate," and "severe" TBI, which may appear in medical records, refer to a classification of TBI made at, or close to, the time of injury		Purely neurological disabilities, such as hemiplegia, cranial nerve paralysis, etc., due to cerebral arteriosclerosis will be rated under the diagnostic codes dealing with such specific disabilities, with citation of a hyphenated diagnostic code (e.g., 8046–8207). Purely subjective complaints such as headache, dizziness, tinnitus, insomnia and irritability, recognized as symptomatic of a properly diagnosed cerebral arteriosclerosis, will be rated 10 percent and no more under diagnostic code 9305. This 10 percent rating will not be combined with any other rating for a disability due to cerebral or generalized arteriosclerosis. Ratings in excess of 10 percent for cerebral arteriosclerosis under diagnostic code 9305 are not assignable in the absence of a diagnosis of multi-infarct dementia with cerebral arteriosclerosis. NOTE: The ratings under code 8046 apply only when the diagnosis of cerebral arteriosclerosis is substantiated by the entire clinical picture	
rather than to the current level of functioning. This classification does not affect		and not solely on findings of retinal arterio- sclerosis.	

Note tioning. This classification does not affect the rating assigned under diagnostic code 8045..

Note (5): A veteran whose residuals of TBI are rated under a version of §4.124a, di-agnostic code 8045, in effect before Octo-ber 23, 2008 may request review under diagnostic code 8045, irrespective of whether his or her disability has worsened whether his or her disability has worsened since the last review. VA will review that veteran's disability rating to determine whether the veteran may be entitled to a higher disability rating under diagnostic code 8045. A request for review pursuant to this note will be treated as a claim for an increased rating for purposes of deter-mining the effective date of an increased rating awarded as a result of such review; however, in no case will the award be effective before October 23, 2008. For the purposes of determining the effective date of an increased rating awarded as a result of such review, VA will apply 38 CFR 3.114, if applicable..

8046 Cerebral arteriosclerosis:

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED

Facets of cognitive impairment and other residuals of TBI not otherwise classified	Level of im- pair- ment	Criteria
Memory, attention, con- centration, executive functions.	0	No complaints of impairment of memory, attention, concentration, or executive functions. A complaint of mild loss of memory (such as having difficulty following a conversation, recalling recent conversations, remembering names of new acquaintances, or finding words, or often misplacing items), attention, concentration, or executive functions, but without objective evidence on testing. Objective evidence on testing of mild impairment of memory, at-
	3	tention, concentration, or executive functions resulting in mild functional impairment. Objective evidence on testing of moderate impairment of memory, attention, concentration, or executive functions resulting in moderate functional impairment.

§4.124a

Department of Veterans Affairs

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED—Continued

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED—Continued

sequences of choices, and make a reasonable decision, although has little difficulty with simple decisions. 3 Moderately severely impaired judgment. For even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. Total Total Total Severely impaired judgment. For even routine and familiar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. For example, unable to determine appropriate Severely impaired judgment. For even routine and familiar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. For example, unable to determine appropriate	npairment and other esiduals of TBI not	evel of n- air- ent	Criteria	Facets of cognitive impairment and other residuals of TBI not otherwise classified	Level of im- pair- ment	Criteria
Judgment	7	otal	testing of severe im- pairment of memory, attention, concentra- tion, or executive func-	Orientation	3	Social interaction is frequently inappropriate. Social interaction is inappropriate most or all of the time. Always oriented to per-
Mildly impaired judgment. For complex or unfamiliar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. 2 Moderately impaired judgment. For complex or unfamiliar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision, although has little difficulty with simple decisions. 3 Moderately severely impaired judgment. For even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decisions. Total Visual spatial orientation O Total Total		0	vere functional impair- ment.	Chornation		son, time, place, and situation.
and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. 2 Moderately impaired judgment. For complex or unfamiliar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision, although has little difficulty with simple decisions. 3 Moderately severely impaired judgment. For even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. Total Severely impaired judgment. For even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. For example, unable to determine appropriate Total	gment	-	Mildly impaired judg- ment. For complex or unfamiliar decisions,		1	Occasionally disoriented to one of the four as- pects (person, time, place, situation) of ori- entation.
judgment. For complex or unfamiliar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision, although has little difficulty with simple decisions. Moderately severely impaired judgment. For even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. Total Total Total Severely impaired judgment. For even routine and familiar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. For example, unable to determine appropriate Visual spatial orientation 1 Total		2	identify, understand, and weigh the alter- natives, understand the consequences of choices, and make a reasonable decision.		2	Occasionally disoriented to two of the four as- pects (person, time, place, situation) of ori- entation or often dis- oriented to one aspect of orientation.
understand the con- sequences of choices, and make a reason- able decision, although has little difficulty with simple decisions. Moderately severely im- paired judgment. For even routine and familiar decisions, occa- sionally unable to iden- tify, understand, and weigh the alternatives, understand the con- sequences of choices, and make a reason- able decision. Total Total Total Motor activity (with intact motor and sensory sys- tem). 1 1 2 2 3 Total			judgment. For complex or unfamiliar decisions, usually unable to iden- tify, understand, and		3	Often disoriented to two or more of the four aspects (person, time, place, situation) of orientation.
Moderately severely impaired judgment. For even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. Total			understand the con- sequences of choices, and make a reason- able decision, although has little difficulty with		Total	Consistently disoriented to two or more of the four aspects (person, time, place, situation) of orientation.
even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. Total Severely impaired judgment. For even routine and familiar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. For example, unable to determine appropriate		3	Moderately severely impaired judgment. For	motor and sensory sys-	0	Motor activity normal.
Total Severely impaired judgment. For even routine and familiar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. For example, unable to determine appropriate Total			iar decisions, occa- sionally unable to iden- tify, understand, and weigh the alternatives, understand the con- sequences of choices, and make a reason-		1	Motor activity normal most of the time, but mildly slowed at times due to apraxia (inabil- ity to perform pre- viously learned motor activities, despite nor- mal motor function).
tify, understand, and weigh the alternatives, understand the con- sequences of choices, and make a reason- able decision. For ex- ample, unable to de- termine appropriate tify, understand, and Total Total	1	otal	Severely impaired judg- ment. For even routine and familiar decisions,		2	Motor activity mildly de- creased or with mod- erate slowing due to apraxia.
sequences of choices, and make a reason- able decision. For ex- ample, unable to de- termine appropriate Total Total Total Total			tify, understand, and weigh the alternatives,		3	Motor activity moderately decreased due to apraxia.
ample, unable to de- Visual spatial orientation 0 termine appropriate 1			sequences of choices, and make a reason-		Total	Motor activity severely decreased due to apraxia.
weather conditions or judge when to avoid dangerous situations or activities.			ample, unable to de- termine appropriate clothing for current weather conditions or judge when to avoid dangerous situations	Visual spatial orientation		Normal. Mildly impaired. Occasionally gets lost in unfamiliar surroundings, has difficulty reading maps or following directions. Is able to use
Social interaction	ial interaction		Social interaction is routinely appropriate.			assistive devices such as GPS (global positioning system).

38 CFR Ch. I (7-1-21 Edition)

§4.124a

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED—Continued

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED—Continued

Facets of cognitive impairment and other residuals of TBI not otherwise classified	Level of im- pair- ment	Criteria	Facets of cognitive impairment and other residuals of TBI not otherwise classified	Level of im- pair- ment	Criteria
Subjective symptoms	3 Total 0	Moderately impaired. Usually gets lost in unfamiliar surroundings, has difficulty reading maps, following directions, and judging distance. Has difficulty using assistive devices such as GPS (global positioning system). Moderately severely impaired. Gets lost even in familiar surroundings, unable to use assistive devices such as GPS (global positioning system). Severely impaired. May be unable to touch or name own body parts when asked by the examiner, identify the relative position in space of two different objects, or find the way from one room to another in a familiar environment. Subjective symptoms that do not interfere with work; instrumental activities of daily living; or work, family, or other close relationships. Examples are: mild or occasional headaches, mild anxiety. Three or more subjective symptoms that mildly interfere with work; instrumental activities of daily living; or work, family, or other close relationships. Examples of findings that might be seen at this level of impairment are: intermittent dizziness, daily mild to moderate headaches, tinnitus, frequent insomnia, hypersensitivity to sound, hypersensitivity to light.	Neurobehavioral effects	0	Three or more subjective symptoms that moderately interfere with work; instrumental activities of daily living; or work, family, or other close relationships. Examples of findings that might be seen at this level of impairment are: marked fatigability, blurred or double vision, headaches requiring rest periods during most days. One or more neurobehavioral effects that do not interfere with workplace interaction or social interaction. Examples of neurobehavioral effects are: Irritability, impulsivity, unpredictability, lack of motivation, verbal aggression, belligerence, apathy, lack of empathy, lack of empathy, lack of empathy, and impaired awareness of disability. Any of these effects may range from slight to severe, although verbal and physical aggression are likely to have a more serious impact on workplace interaction and social interaction than some of the other effects. One or more neurobehavioral effects that occasionally interfere with workplace interaction, or both but do not preclude them. One or more neurobehavioral effects that frequently interfere with workplace interaction, or both but do not preclude them. One or more neurobehavioral effects that frequently interfere with workplace interaction, social interaction, social interaction, sobid but do not preclude
					them.

Department of Veterans Affairs

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED—Continued

Facets of cognitive impairment and other residuals of TBI not otherwise classified	Level of im- pair- ment	Criteria
	3	One or more neurobehavioral effects that interfere with or preclude workplace interaction, or both on most days or that occasionally require supervision for safety of self or others.
Communication	0	Able to communicate by spoken and written language (expressive communication), and to comprehend spoken and written language.
	1	Comprehension or ex- pression, or both, of either spoken lan- guage or written lan- guage is only occa- sionally impaired. Can communicate complex ideas.
	2	Inability to communicate either by spoken language, written language, or both, more than occasionally but less than half of the time, or to comprehend spoken language.
	3	guage, written lan- guage, or both, more than occasionally but less than half of the time. Can generally communicate complex ideas. Inability to communicate
		either by spoken lan- guage, written lan- guage, or both, at least half of the time but not all of the time, or to comprehend spo- ken language, written language, or both, at least half of the time
	Total	but not all of the time. May rely on gestures or other alternative modes of communication. Able to communicate basic needs. Complete inability to communicate either by spoken language, written language, or both, or to comprehend spoken language, written language, or both. Unable to communicate basic needs.

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED—Continued

Facets of cognitive impairment and other residuals of TBI not otherwise classified	Level of im- pair- ment	Criteria
Consciousness	Total	Persistently altered state of consciousness, such as vegetative state, minimally re- sponsive state, coma.

MISCELLANEOUS DISEASES

	Rat- ing
8100 Migraine:	
With very frequent completely prostrating and prolonged attacks productive of severe economic inadaptability	50
With characteristic prostrating attacks occurring on an average once a month over last several months	30
With characteristic prostrating attacks averaging one in 2 months over last several months	10
With less frequent attacks	0
8103 Tic, convulsive:	00
Severe Moderate	30 10
Mild	0
Note: Depending upon frequency, severity, mus-	U
cle groups involved.	
8104 Paramyoclonus multiplex (convulsive state,	
myoclonic type):	
Rate as tic; convulsive; severe cases	60
8105 Chorea, Sydenham's:	
Pronounced, progressive grave types	100
Severe	80
Moderately severe	50
Moderate	30
Mild	10
NOTE: Consider rheumatic etiology and com-	
plications.	
8106 Chorea, Huntington's.	
Rate as Sydenham's chorea. This, though a fa-	
milial disease, has its onset in late adult life,	
and is considered a ratable disability.	
8107 Athetosis, acquired.	
Rate as chorea.	
8108 Narcolepsy.	
Rate as for epilepsy, petit mal.	
DISEASES OF THE CRANIAL NERVES	

DISEASES OF THE CRANIAL NERVES

	Rat- ing
Disability from lesions of peripheral portions of first, second, third, fourth, sixth, and eighth nerves will be rated under the Organs of Special Sense. The ratings for the cranial nerves are for unilateral involvement; when bilateral, combine but without the bilateral factor. Fifth (trigeminal) cranial nerve 8205 Paralysis of:	
Complete	50
Incomplete, severe	30 10
incomplete, moderate	10

38 CFR Ch. I (7-1-21 Edition)

§4.124a

DISEASES OF THE CRANIAL NERVES—Continued

DISEASES OF THE PERIPHERAL NERVES

	Rat-	Schedule of ratings		ing
	ing	Schedule of ratings	Major	Minor
NOTE: Dependent upon relative degree of sensory manifestation or motor loss. 8305 Neuritis. 8405 Neuralgia. NOTE: Tic douloureux may be rated in accordance with severity, up to complete paralysis. Seventh (facial) cranial nerve		The term "incomplete paralysis," with this and other peripheral nerve injuries, indicates a degree of lost or impaired function substantially less than the type picture for complete paralysis given with each nerve, whether due to varied level of the nerve lesion or to partial regenera-		
3207 Paralysis of: Complete	30 20 10	of the flere lesion of to partial regenera- tion. When the involvement is wholly sensory, the rating should be for the mild, or at most, the moderate degree. The ratings for the peripheral nerves are for unilateral involvement; when bilateral, combine with application of the bilateral factor.		
8407 Neuralgia. Ninth (glossopharyngeal) cranial nerve.		Upper radicular group (fifth and sixth cervicals)		
8209 Paralysis of: Complete Incomplete, severe Incomplete, moderate NOTE: Dependent upon relative loss of ordinary sensation in mucous membrane of the phar-	30 20 10	8510 Paralysis of: Complete; all shoulder and elbow movements lost or severely affected, hand and wrist movements not affected	70	6
ynx, fauces, and tonsils. 8309 Neuritis.		Severe	50	4
8409 Neuralgia.		Moderate Mild	40 20	3 2
Tenth (pneumogastric, vagus) cranial nerve. 8210 Paralysis of: Complete	50 30 10	8610 Neuritis. 8710 Neuralgia. Middle radicular group 8511 Paralysis of: Complete; adduction, abduction and rotation of arm, flexion of elbow, and extension of wrist lost or severely af-		
3310 Neuritis. 3410 Neuralgia.		fectedIncomplete:	70	6
Eleventh (spinal accessory, external branch) cra-		Severe	50	4
nial nerve.		Moderate	40	3
8211 Paralysis of: Complete	30 20 10	Mild	20	2
8311 Neuritis. 8411 Neuralgia. Twelfth (hypoglossal) cranial nerve.		Complete; all intrinsic muscles of hand, and some or all of flexors of wrist and fingers, paralyzed (sub-		
3212 Paralysis of: Complete	50	stantial loss of use of hand)	70	6
Incomplete, severe	30	Severe Moderate	50 40	3
Incomplete, moderate	10	Mild	20	2
8412 Neuralgia.		All radicular groups		
		8513 Paralysis of: Complete Incomplete:	90	8
		Severe	70	6
		Moderate Mild	40 20	30

Department of Veterans Affairs

DISEASES OF THE PERIPHERAL NERVES-Continued

		Rat	ing
	Schedule of ratings		Minor
8613 8713	Neuritis. Neuralgia.		
0/13	Neuraigia.		
The	musculospiral nerve (radial nerve)		
	Paralysis of: omplete; drop of hand and fingers, wrist and fingers perpetually flexed, the thumb adducted falling within the line of the outer border of the index finger; can not extend hand at wrist, extend proximal phalanges of fingers, extend thumb, or make lateral movement of wrist; supination of hand, extension and flexion of elbow weakened, the loss of synergic motion of extensors impairs the hand grip seriously; total paralysis of the triceps occurs only as the greatest		
le.	rarity	70	60
III	Severe	50	40
	Moderate	30	20
	Mild	20	20
8614 8714	Neuritis.		
N	OTE: Lesions involving only "dissocia communis digitorum" and "paralysis I sor communis digitorum," will not e	oelow the	exten-

erate rating under code 8514.

· ·		
The median nerve		
8515 Paralysis of: Complete; the hand inclined to the ulnar side, the index and middle fingers more extended than normally, considerable atrophy of the muscles of the thenar eminence, the thumb in the plane of the hand (ape hand); pronation incomplete and defective, absence of flexion of index finger and feeble flexion of middle finger, cannot make a fist, index and middle fingers remain extended; cannot flex distal phalanx of thumb, defective opposition and abduction of the thumb, at right angles to palm; flexion of wrist weakened; pain with trophic disturbances Incomplete: Severe Moderate Mild Neuritis. 8615 Neuritis.	70 50 30 10	60 40 20 10
The ulnar nerve		
8516 Paralysis of: Complete; the "griffin claw" deformity, due to flexor contraction of ring and little fingers, atrophy very marked in dorsal interspace and thenar and hypothenar eminences; loss of extension of ring and little fingers cannot spread the fingers (or reverse), cannot adduct the thumb; flexor of wrist weakened	60	50
Incomplete:	40	00
Severe Moderate	40 30	30 20

DISEASES OF THE PERIPHERAL NERVES-Continued

Continued		
	Rating	
Schedule of ratings	Major	Minor
Mild 8616 Neuritis. 8716 Neuralgia.	10	10
Musculocutaneous nerve		
8517 Paralysis of:		
Complete; weakness but not loss of flexion of elbow and supination of forearm	30	20
Incomplete:		
Severe	20	20
Moderate Mild	10	10
8617 Neuritis.	0	1
8717 Neuralgia.		
Circumflex nerve		
8518 Paralysis of:		
Complete; abduction of arm is impos- sible, outward rotation is weakened; muscles supplied are deltoid and teres minor	50	40
Incomplete: Severe	30	21
Moderate	10	1
Mild	0	
8618 Neuritis. 8718 Neuralgia.		
Long thoracic nerve		
8519 Paralysis of: Complete; inability to raise arm above shoulder level, winged scapula de-		
formityIncomplete:	30	2
Severe	20	2
Moderate	10	10
Mild	0	(
NOTE: Not to be combined with lost mo der level.	tion above	shoul-
8619 Neuritis. 8719 Neuralgia.		
NOTE: Combined nerve injuries should	be rated	by ref-

NOTE: Combined nerve injuries should be rated by reference to the major involvement, or if sufficient in extent, consider radicular group ratings.

	Rating
Sciatic nerve	
8520 Paralysis of:	
Complete; the foot dangles and drops, no active movement possible of muscles below the knee, flexion of	80
knee weakened or (very rarely) lost Incomplete:	80
Severe, with marked muscular at-	
rophy	60
Moderately severe	40
Moderate	20
Mild	10

38 CFR Ch. I (7-1-21 Edition)

§4.124a

	Rating		Rating
8620 Neuritis.		Incomplete:	
8720 Neuralgia.		Severe	20
		Moderate	10
External popliteal nerve (common		Mild	10
peroneal)		8625 Neuritis.	
8521 Paralysis of:		8725 Neuralgia.	
Complete; foot drop and slight droop		Anterior crural nerve (femoral)	
of first phalanges of all toes, cannot		8526 Paralysis of:	
dorsiflex the foot, extension (dorsal		Complete; paralysis of quadriceps ex-	
flexion) of proximal phalanges of		tensor muscles	40
toes lost; abduction of foot lost,		Incomplete:	
adduction weakened; anesthesia		Severe	30
covers entire dorsum of foot and		Moderate	20
toes	40	Mild	10
Incomplete:		8626 Neuritis.	
Severe	30	8726 Neuralgia.	
Moderate	20	Internal saphenous nerve	
Mild	10	-	
8621 Neuritis.		8527 Paralysis of: Severe to complete	10
8721 Neuralgia.		Mild to moderate	0
Musculocutaneous nerve (superficial		8627 Neuritis.	U
peroneal)		8727 Neuralgia.	
8522 Paralysis of:		-	
Complete; eversion of foot weakened	30	Obturator nerve	
Incomplete:	30	8528 Paralysis of:	
Severe	20	Severe to complete	10
Moderate	10	Mild or moderate	0
Mild	0	8628 Neuritis.	
8622 Neuritis.	O	8728 Neuralgia.	
8722 Neuralgia.		External cutaneous nerve of thigh	
•		8529 Paralysis of:	
Anterior tibial nerve (deep peroneal)		Severe to complete	10
8523 Paralysis of:		Mild or moderate	0
Complete; dorsal flexion of foot lost	30	8629 Neuritis.	
Incomplete:		8729 Neuralgia.	
Severe	20	Ilio-inguinal nerve	
Moderate	10	8530 Paralysis of:	
Mild	0	Severe to complete	10
8623 Neuritis.		Mild or moderate	0
8723 Neuralgia.		8630 Neuritis.	Ū
Internal popliteal nerve (tibial)		8730 Neuralgia.	
8524 Paralysis of:		8540 Soft-tissue sarcoma (of neurogenic	
Complete; plantar flexion lost, frank		origin)	100
adduction of foot impossible, flexion		NOTE: The 100 percent rating will be co	ntinued
and separation of toes abolished; no		for 6 months following the cessation	
muscle in sole can move; in lesions		gical, X-ray, antineoplastic chemothe	
of the nerve high in popliteal fossa,		other therapeutic procedure. At this	point, if
plantar flexion of foot is lost	40	there has been no local recurrence or	metas-
Incomplete:		tases, the rating will be made on resid	uals.
Severe	30		
Moderate	20	THE EPILEPSIES	
Mild	10		
8624 Neuritis.			Rat-
8724 Neuralgia.			ing
Posterior tibial nerve		A thorough study of all material in §§4.121 an	d
8525 Paralysis of:		4.122 of the preface and under the ratings for	
Complete; paralysis of all muscles of		epilepsy is necessary prior to any rating ac	>-
sole of foot, frequently with painful		tion. 8910 Epilepsy, grand mal.	
paralysis of a causalgic nature; toes		Rate under the general rating formula for major	or
cannot be flexed; adduction is weak-		seizures.	
ened; plantar flexion is impaired	30	8911 Epilepsy, petit mal.	1

THE EPILEPSIES—Continued

Rate under the general rating formula for minor seizures.
NOTE (1): A major seizure is characterized by the generalized tonic-clonic convulsion with unconsciousness.
NOTE (2): A minor seizure consists of a brief interruption in consciousness or conscious control associated with staring or rhythmic blinking of the eyes or nodding of the head ("pure" petit mal), or sudden jerking movements of the arms, trunk, or head (myoclonic type) or sudden loss of postural control
(akinetic type).
General Rating Formula for Major and Minor Epileptic Seizures:
Averaging at least 1 major seizure per
month over the last year
Averaging at least 1 major seizure in 3 months over the last year; or more than
10 minor seizures weekly
Averaging at least 1 major seizure in 4 months over the last year; or 9–10 minor seizures per week
At least 1 major seizure in the last 6 months
or 2 in the last year; or averaging at least
5 to 8 minor seizures weekly
At least 1 major seizure in the last 2 years;
or at least 2 minor seizures in the last 6
months
A confirmed diagnosis of epilepsy with a
history of seizures
NOTE (1): When continuous medication is shown
necessary for the control of epilepsy, the min- imum evaluation will be 10 percent. This rating
will not be combined with any other rating for

NOTE (2): In the presence of major and minor seizures, rate the predominating type.

NOTE (3): There will be no distinction between

diurnal and nocturnal major seizures

8912 Epilepsy, Jacksonian and focal motor or sensory.

8913 Epilepsy, diencephalic.

epilepsy.

Rate as minor seizures, except in the presence of major and minor seizures, rate the predominating type.

8914 Epilepsy, psychomotor.

Major seizures:

Psychomotor seizures will be rated as major seizures under the general rating formula when characterized by automatic states and/or generalized convulsions with unconsciousness

Minor seizures:

Psychomotor seizures will be rated as minor seizures under the general rating formula when characterized by brief transient episodes of random motor movements, hallucinations, perceptual illusions, abnormalities of thinking, memory or mood, or autonomic disturbances

Mental Disorders in Epilepsies: A nonpsychotic organic Mental Disorders in Epilepsies: A nonpsychotic organic brain syndrome will be rated separately under the appropriate diagnostic code (e.g., 9304 or 9326). In the absence of a diagnostic ropersychotic organic psychiatric disturbance (psychotic, psychoneurotic or personality disorder) if diagnosed and shown to be secondary to or directly associated with epilepsy will be rated separately. The psychotic or psychroneurotic disorder will be rated under the appropriate diagnostic code. The personality disorder will be rated as a diágnostic code. The personality disorder will be rated as a dementia (e.g., diagnostic code 9304 or 9326). Epilepsy and Unemployability: (1) Rating specialists must bear in mind that the epileptic, although his or her seizures are controlled, may find employment and rehabilitation difficult of attainment due to employer reluctance to the hiring of the

(2) Where a case is encountered with a definite history of unemployment, full and complete development should be un-dertaken to ascertain whether the epilepsy is the determining factor in his or her inability to obtain employment.

(3) The assent of the claimant should first be obtained for

(a) The assent of the challful should his be obtained by permission to conduct this economic and social survey. The purpose of this survey is to secure all the relevant facts and data necessary to permit of a true judgment as to the reason for his or her unemployment and should include information

Rat-

100

80

60

40

20

10

as to:

(a) Education;
(b) Occupations prior and subsequent to service;
(c) Places of employment and reasons for termination;
(d) Wages received;
(e) Number of seizures.
(4) Upon completion of this survey and current examination, the case should have rating board consideration. Where in the judgment of the rating board the veteran's unemployability is due to epilepsy and jurisdiction is not vested in that body by reason of schedular evaluations, the case should be submitted to the Compensation Service or the Director, Pension and Fiduciary Service. and Fiduciary Service.

(Authority: 38 U.S.C. 1155)

[29 FR 6718, May 22, 1964, as amended at 40 FR 42540, Sept. 15, 1975; 41 FR 11302, Mar. 18, 1976; 43 FR 45362, Oct. 2, 1978; 54 FR 4282, Jan. 30, 1989; 54 FR 49755, Dec. 1, 1989; 55 FR 154, Jan. 3, 1990; 56 FR 51653, Oct. 15, 1991; 57 FR 24364, June 9, 1992; 70 FR 75399, Dec. 20, 2005; 73 FR 54705, Sept. 23, 2008; 73 FR 69554, Nov. 19, 2008; 76 FR 78824, Dec. 20, 2011; 79 FR 2100, Jan. 13, 2014]

MENTAL DISORDERS

§ 4.125 Diagnosis of mental disorders.

(a) If the diagnosis of a mental disorder does not conform to DSM-5 or is not supported by the findings on the examination report, the rating agency shall return the report to the examiner to substantiate the diagnosis. Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5), American Psychiatric Association (2013), is incorporated by reference into this section with the approval of the Director of the Federal Register under 5 U.S.C. 552(a) and 1 CFR part 51. To enforce any edition other than that specified in this section, the Department of Veterans Affairs must publish notice of change in the FEDERAL REGISTER and the material must be available to the public. All approved material is available from the American Psychiatric Association, 1000 Wilson Boulevard, Suite 1825, Arlington, VA 22209-3901, 703-907-7300, http://www.dsm5.org. It is also available for inspection at the Office of Regulation Policy and Management, Department of Veterans Affairs, 810 Vermont Avenue NW., Room 1068,

Washington, DC 20420. It is also available for inspection at the National Archives and Records Administration (NARA). For information on the availability of this information at NARA, call 202–741–6030 or go to https://www.archives.gov/federal_register/code_of_federal_regulations/ ibr publications.html.

(b) If the diagnosis of a mental disorder is changed, the rating agency shall determine whether the new diagnosis represents progression of the prior diagnosis, correction of an error in the prior diagnosis, or development of a new and separate condition. If it is not clear from the available records what the change of diagnosis represents, the rating agency shall return the report to the examiner for a determination.

(Authority: 38 U.S.C. 1155)

[61 FR 52700, Oct. 8, 1996, as amended at 79 FR 45099, Aug. 4, 2014]

§ 4.126 Evaluation of disability from mental disorders.

- (a) When evaluating a mental disorder, the rating agency shall consider the frequency, severity, and duration of psychiatric symptoms, the length of remissions, and the veteran's capacity for adjustment during periods of remission. The rating agency shall assign an evaluation based on all the evidence of record that bears on occupational and social impairment rather than solely on the examiner's assessment of the level of disability at the moment of the examination.
- (b) When evaluating the level of disability from a mental disorder, the rating agency will consider the extent of social impairment, but shall not assign an evaluation solely on the basis of social impairment.
- (c) Neurocognitive disorders shall be evaluated under the general rating formula for mental disorders; neurologic deficits or other impairments stemming from the same etiology (e.g., a head injury) shall be evaluated separately and combined with the evaluation for neurocognitive disorders (see §4.25).
- (d) When a single disability has been diagnosed both as a physical condition and as a mental disorder, the rating agency shall evaluate it using a diag-

nostic code which represents the dominant (more disabling) aspect of the condition (see § 4.14).

(Authority: 38 U.S.C. 1155)

[61 FR 52700, Oct. 8, 1996, as amended at 79 FR 45099, Aug. 4, 2014]

§ 4.127 Intellectual disability (intellectual developmental disorder) and personality disorders.

Intellectual disability (intellectual developmental disorder) and personality disorders are not diseases or injuries for compensation purposes, and, except as provided in §3.310(a) of this chapter, disability resulting from them may not be service-connected. However, disability resulting from a mental disorder that is superimposed upon intellectual disability (intellectual developmental disorder) or a personality disorder may be service-connected.

(Authority: 38 U.S.C. 1155)

[79 FR 45100, Aug. 4, 2014]

§ 4.128 Convalescence ratings following extended hospitalization.

If a mental disorder has been assigned a total evaluation due to a continuous period of hospitalization lasting six months or more, the rating agency shall continue the total evaluation indefinitely and schedule a mandatory examination six months after the veteran is discharged or released to nonbed care. A change in evaluation based on that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.

(Authority: 38 U.S.C. 1155)

 $[61~{\rm FR}~52700,\,{\rm Oct.}~8,\,1996]$

§4.129 Mental disorders due to traumatic stress.

When a mental disorder that develops in service as a result of a highly stressful event is severe enough to bring about the veteran's release from active military service, the rating agency shall assign an evaluation of not less than 50 percent and schedule an examination within the six month period

Department of Veterans Affairs

following the veteran's discharge to determine whether a change in evaluation is warranted.

(Authority: 38 U.S.C. 1155) [61 FR 52700, Oct. 8, 1996]

§ 4.130 Schedule of ratings—Mental disorders.

The nomenclature employed in this portion of the rating schedule is based upon the American Psychiatric Association's Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5) (see § 4.125 for availability information). Rating agencies must be thoroughly familiar with this manual to properly implement the directives in § 4.125 through § 4.129 and to apply the general rating formula for mental disorders in § 4.130. The schedule for rating for mental disorders is set forth as follows:

- 9201 Schizophrenia 9202 [Removed] 9203 [Removed] 9204 [Removed] 9205 [Removed] 9208 Delusional disorder 9210 Other specified and unspecified schizophrenia spectrum and other psychotic disorders 9211 Schizoaffective disorder
- 9300 Delirium 9301 Major or mild neurocognitive disorder due to HIV or other infections

- 9304 Major or mild neurocognitive disorder due to traumatic brain injury
- 9305 Major or mild vascular neurocognitive disorder
- 9310 Unspecified neurocognitive disorder
- 9312 Major or mild neurocognitive disorder due to Alzheimer's disease
- 9326 Major or mild neurocognitive disorder due to another medical condition or substance/medication-induced major or mild neurocognitive disorder
- 9327 [Removed]
- 9400 Generalized anxiety disorder
- 9403 Specific phobia; social anxiety disorder (social phobia)
- 9404 Obsessive compulsive disorder
- 9410 Other specified anxiety disorder
- 9411 Posttraumatic stress disorder
- 9412 Panic disorder and/or agoraphobia
- 9413 Unspecified anxiety disorder
- 9416 Dissociative amnesia; dissociative identity disorder
- 9417 Depersonalization/Derealization dis
- 9421 Somatic symptom disorder
- 9422 Other specified somatic symptom and related disorder
- 9423 Unspecified somatic symptom and related disorder
- 9424 Conversion disorder (functional neurological symptom disorder)
- 9425 Illness anxiety disorder
- 9431 Cyclothymic disorder
- 9432 Bipolar disorder
- 9433 Persistent depressive disorder (dysthymia)
- 9434 Major depressive disorder
- 9435 Unspecified depressive disorder
- 9440 Chronic adjustment disorder

GENERAL RATING FORMULA FOR MENTAL DISORDERS

	Rating
Total occupational and social impairment, due to such symptoms as: gross impairment in thought processes or communication; persistent delusions or hallucinations; grossly inappropriate behavior; persistent danger of hurting self or others; intermittent inability to perform activities of daily living (including maintenance of minimal personal hygiene); disorientation to time or place; memory loss for names of close relatives, own occupation, or own name. Occupational and social impairment, with deficiencies in most areas, such as work, school, family relations, judging the processing of the processi	100
ment, thinking, or mood, due to such symptoms as: suicidal ideation; obsessional rituals which interfere with rou- tine activities; speech intermittently illogical, obscure, or irrelevant; near-continuous panic or depression affecting the ability to function independently, appropriately and effectively; impaired impulse control (such as unprovoked irritability with periods of violence); spatial disorientation; neglect of personal appearance and hygiene; difficulty in adapting to stressful circumstances (including work or a worklike setting); inability to establish and maintain effec- tive relationships.	70
Occupational and social impairment with reduced reliability and productivity due to such symptoms as: flattened af- fect; circumstantial, circumlocutory, or stereotyped speech; panic attacks more than once a week; difficulty in un- derstanding complex commands; impairment of short- and long-term memory (e.g., retention of only highly learned material, forgetting to complete tasks); impaired judgment; impaired abstract thinking; disturbances of mo-	70
tivation and mood; difficulty in establishing and maintaining effective work and social relationships. Occupational and social impairment with occasional decrease in work efficiency and intermittent periods of inability to perform occupational tasks (although generally functioning satisfactorily, with routine behavior, self-care, and conversation normal), due to such symptoms as: depressed mood, anxiety, suspiciousness, panic attacks (weekly	50
or less often), chronic sleep impairment, mild memory loss (such as forgetting names, directions, recent events). Occupational and social impairment due to mild or transient symptoms which decrease work efficiency and ability to perform occupational tasks only during periods of significant stress, or symptoms controlled by continuous medi-	30
cation.	10
A mental condition has been formally diagnosed, but symptoms are not severe enough either to interfere with occu- pational and social functioning or to require continuous medication.	0

9520 Anorexia nervosa 9521 Bulimia nervosa

RATING FORMULA FOR EATING DISORDERS

	Rating
Self-induced weight loss to less than 80 percent of expected minimum weight, with incapacitating episodes of at least six weeks total duration per year, and requiring hospitalization more than twice a year for parenteral nutrition or tube feeding.	100
Self-induced weight loss to less than 85 percent of expected minimum weight with incapacitating episodes of six or more weeks total duration per year.	60
Self-induced weight loss to less than 85 percent of expected minimum weight with incapacitating episodes of more than two but less than six weeks total duration per year.	30
Binge eating followed by self-induced vomiting or other measures to prevent weight gain, or resistance to weight gain even when below expected minimum weight, with diagnosis of an eating disorder and incapacitating episodes of up to two weeks total duration per year.	10
Binge eating followed by self-induced vomiting or other measures to prevent weight gain, or resistance to weight gain even when below expected minimum weight, with diagnosis of an eating disorder but without incapacitating episodes.	0

Note 1: An incapacitating episode is a period during which bed rest and treatment by a physician are required.

Note 2: Ratings under diagnostic codes 9201 to 9440 will be evaluated using the General Rating Formula for Mental Disorders. Ratings under diagnostic codes 9520 and 9521 will be evaluated using the General Rating Formula for Eating Disorders.

(Authority: 38 U.S.C. 1155)			Rat-
[79 FR 45100, Aug. 4, 2014]			ing
DENTAL AND ORAL CONDITIONS		Not involving temporomandibular articulation.	
§4.149 [Reserved]		Not replaceable by prosthesis	20 10
84470 01 11 4 4		nostic imaging studies:	
§ 4.150 Schedule of ratings—denta oral conditions.	l and	Severe, with false motion	30
oral conditions.		Moderate, without false motion	10
	Rat-	9904 Mandible, malunion of:	
	ing	Displacement, causing severe anterior or posterior open bite	20
		Displacement, causing moderate anterior or	20
Note (1): For VA compensation purposes, diagnostic		posterior open bite	10
imaging studies include, but are not limited to, con-		Displacement, not causing anterior or pos-	
ventional radiography (X-ray), computed tomography (CT), magnetic resonance imaging (MRI),		terior open bite	0
positron emission tomography (PET), radionuclide		9905 Temporomandibular disorder (TMD):	
bone scanning, or ultrasonography		Interincisal range:	
Note (2): Separately evaluate loss of vocal articula-		0 to 10 millimeters (mm) of max-	
tion, loss of smell, loss of taste, neurological im-		imum unassisted vertical open-	
pairment, respiratory dysfunction, and other impair-		ing.	
ments under the appropriate diagnostic code and combine under § 4.25 for each separately rated		With dietary restrictions to all mechanically altered	
condition 94.25 for each separately rated		foods	50
9900 Maxilla or mandible, chronic osteomyelitis,		Without dietary restrictions	
osteonecrosis or osteoradionecrosis of:		to mechanically altered	
Rate as osteomyelitis, chronic under diag-		foods	40
nostic code 5000		11 to 20 mm of maximum unas-	
9901 Mandible, loss of, complete, between angles	100	sisted vertical opening.	
9902 Mandible, loss of, including ramus, unilaterally		With dietary restrictions to all mechanically altered	
or bilaterally:		foods	40
Loss of one-half or more,.		Without dietary restrictions	
Involving temporomandibular articu- lation.		to mechanically altered	
Not replaceable by prosthesis	70	foods	30
Replaceable by prosthesis	50	21 to 29 mm of maximum unas-	
Not involving temporomandibular	"	sisted vertical opening.	
articulation.		With dietary restrictions to full liquid and pureed	
Not replaceable by prosthesis	40	foods	40
Replaceable by prosthesis	30	With dietary restrictions to	
Loss of less than one-half,		soft and semi-solid	
Involving temporomandibular articu-		foods	30
lation.	70	Without dietary restrictions	
Not replaceable by prosthesis Replaceable by prosthesis	70 50	to mechanically altered	20
neplaceable by prostnesis	1 50	foods	20

Department of Veterans Affairs

Pt. 4, App. A

	Rat- ing		Rat- ing
30 to 34 mm of maximum unas-		Where the loss of masticatory surface can	
sisted vertical opening. With dietary restrictions to full liquid and pureed		be restored by suitable prosthesis NOTE—These ratings apply only to bone loss through trauma or disease such as	0
foods	30	osteomyelitis, and not to the loss of the alveolar process as a result of periodontal	
foodsWithout dietary restrictions	20	disease, since such loss is not considered disabling. 9914 Maxilla, loss of more than half:	
to mechanically altered foods	10	Not replaceable by prosthesis	100 50
Lateral excursion range of motion: 0 to 4 mm	10	9915 Maxilla, loss of half or less: Loss of 25 to 50 percent:	50
Note (1): Ratings for limited interincisal movement shall not be combined with ratings for limited lat- eral excursion.		Not replaceable by prosthesis Replaceable by prosthesis	40 30
Note (2): For VA compensation purposes, the normal maximum unassisted range of vertical jaw opening		Loss of less than 25 percent: Not replaceable by prosthesis	20
is from 35 to 50 mm. Note (3): For VA compensation purposes, mechanically altered foods are defined as altered by blend-		Replaceable by prosthesis	0
ing, chopping, grinding or mashing so that they are		Nonunion, With false motion	30
easy to chew and swallow. There are four levels of mechanically altered foods: full liquid, puree, soft, and semisolid foods. To warrant elevation based		Without false motion	10
on mechanically altered foods, the use of texture- modified diets must be recorded or verified by a physician.		With displacement, causing severe anterior or posterior open bite With displacement, causing mod-	30
9908 Condyloid process, loss of, one or both sides 9909 Coronoid process, loss of:	30	erate anterior or posterior open bite	10
Bilateral	20 10	With displacement, causing mild anterior or posterior open bite Note: For VA compensation purposes, the severity of	0
9911 Hard palate, loss of: Loss of half or more, not replaceable by prosthesis	30	maxillary nonunion is dependent upon the degree of abnormal mobility of maxilla fragments following	
Loss of less than half, not replaceable by prosthesis	20	treatment (i.e., presence or absence of false motion), and maxillary nonunion must be confirmed by	
Loss of half or more, replaceable by pros- thesis Loss of less than half, replaceable by pros-	10	diagnostic imaging studies. 9917 Neoplasm, hard and soft tissue, benign: Rate as loss of supporting structures (bone	
thesis	0	or teeth) and/or functional impairment due to scarring.	
body of maxilla or mandible without loss of continuity:		9918 Neoplasm, hard and soft tissue, malignant Note: A rating of 100 percent shall continue beyond	100
Where the lost masticatory surface cannot be restored by suitable prosthesis:		the cessation of any surgical, radiation, antineoplastic chemotherapy or other therapeutic	
Loss of all teeth Loss of all upper teeth	40 30	procedure. Six months after discontinuance of	
Loss of all lower teeth	30	such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any	
missingAll upper and lower anterior teeth	20	subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. If there has	
missing All upper anterior teeth missing	20 10	been no local recurrence or metastasis, rate on re- siduals such as loss of supporting structures (bone	
All lower anterior teeth missing All upper and lower teeth on one side missing	10	or teeth) and/or functional impairment due to scarring.	

 $[59~{\rm FR}~2530,\,{\rm Jan.}~18,\,1994,\,{\rm as}~{\rm amended}~{\rm at}~82~{\rm FR}~36083,\,{\rm Aug.}~3,\,2017]$

APPENDIX A TO PART 4—TABLE OF AMENDMENTS AND EFFECTIVE DATES SINCE 1946

Sec.	Diagnostic code No.	
4.71a4.71a	5001 5002 5003 5009	Evaluation February 1, 1962. Evaluation March 11, 1969; criterion February 7, 2021. Evaluation March 1, 1963; title, criteria, note February 7, 2021. Added July 6, 1950; title February 7, 2021. Title, evaluation, note February 7, 2021. Title, criteria February 7, 2021.

4, App. A		30 CFR CII. 1 (7-1-21 Edilloli
Sec.	Diagnostic code No.	
	5011	Title, criteria February 7, 2021.
	5012	Criterion March 10, 1976; title, note February 7, 2021.
	5013	Title February 7, 2021.
	5014	Title February 7, 2021.
	5015	Title February 7, 2021.
	5018 5020	Removed February 7, 2021. Removed November 30, 2020.
	5022	Removed February 7, 2021.
	5023	Title February 7, 2021.
	5024	Criterion March 1, 1963; title, criteria February 7, 2021.
	5025 5051	Added May 7, 1996. Added September 22, 1978; note February 7, 2021.
	5052	Added September 22, 1978; note February 7, 2021.
	5053	Added September 22, 1978; note February 7, 2021.
	5054	Added September 22, 1978; title, criterion, and note February 7, 2021.
	5055	Added September 22, 1978; title, criterion, and note February 7, 2021.
	5056 5100–5103	Added September 22, 1978; note February 7, 2021. Removed March 10, 1976.
	5104	Criterion March 10, 1976.
	5105	Criterion March 10, 1976.
	5120	Title, criterion February 7, 2021.
	5160	Title, criterion, note February 7, 2021.
	5164 5166	Evaluation June 9, 1952. Criterion September 22, 1978.
	5170	Title February 7, 2021.
	5172	Added July 6, 1950.
	5173	Added June 9, 1952.
	5174	Added September 9, 1975; removed September 22, 1978.
	5201 5202	Criterion February 7, 2021. Criterion February 7, 2021.
	5211	Criterion September 22, 1978.
	5212	Criterion September 22, 1978.
	5214	Criterion September 22, 1978.
	5216	Preceding paragraph criterion September 22, 1978.
	5217 5218	Criterion August 26, 2002. Criterion August 26, 2002.
	5219	Criterion September 22, 1978; criterion August 26, 2002.
	5220	Preceding paragraph criterion September 22, 1978; criterion August 26, 2002.
	5223	Criterion August 26, 2002.
	5224 5225	Criterion August 26, 2002. Criterion August 26, 2002.
	5226	Criterion August 26, 2002.
	5227	Criterion September 22, 1978; criterion August 26, 2002.
	5228	Added August 26, 2002.
	5229	Added August 26, 2002.
	5230 5235	Added August 26, 2002. Replaces 5285–5295 September 26, 2003.
	5236	Replaces 5285–5295 September 26, 2003.
	5237	Replaces 5285–5295 September 26, 2003.
	5238	Replaces 5285–5295 September 26, 2003.
	5239 5240	Replaces 5285–5295 September 26, 2003. Replaces 5285–5295 September 26, 2003.
	5240	Replaces 5285–5295 September 26, 2003.
	5242	Replaces 5285–5295 September 26, 2003; Title February 7, 2021.
	5243	Replaces 5285–5295 September 26, 2003; Criterion September 26, 2003; Ti February 7, 2021.
	5244	Added February 7, 2021.
	5255 5257	Criterion July 6, 1950; criterion February 7, 2021. Evaluation July 6, 1950; criterion and note February 7, 2021.
	5262	Criterion February 7, 2021.
	5264	Added September 9, 1975; removed September 22, 1978.
	5269	Added February 7, 2021.
	5271	Criterion February 7, 2021.
	5275 5293	Criterion March 10, 1976; criterion September 22, 1978. Criterion March 10, 1976; criterion September 23, 2002; revised and moved
	5294	5235–5243 September 26, 2003. Evaluation March 10, 1976; revised and moved to 5235–5243 September 2 2003.
	5295	Evaluation March 10, 1976; revised and moved to 5235–5243 September 2 2003.
	5296	Criterion March 10, 1976.
	5297	Criterion August 23, 1948; criterion February 1, 1962.
	5298	Added August 23, 1948.

Sec.	Diagnostic code No.	
4.73		Introduction Note criterion July 3, 1997; second Note added February 7, 2021.
	5317	Criterion September 22, 1978.
	5324	Added February 1, 1962.
	5325	Criterion July 3, 1997.
	5327	Added March 10, 1976; criterion October 15, 1991; criterion July 3, 1997.
	5328	Added NOTE March 10, 1976.
	5329	Added NOTE July 3, 1997.
	5330 5331	Added February 7, 2021. Added February 7, 2021.
l.77		Revised May 13, 2018.
.78		Revised May 13, 2018.
1.79		Introduction criterion May 13, 2018; Revised General Rating Formula for Dis
		eases of the Eye NOTE revised May 13, 2018.
	6000	Criterion May 13, 2018.
	6001	Criterion May 13, 2018.
	6002	Criterion May 13, 2018.
	6006	Title May 13, 2018. Criterion May 13, 2018.
	6007	Criterion May 13, 2018.
	6008 6009	Criterion May 13, 2018.
	6011	Criterion May 13, 2018. Evaluation May 13, 2018.
	6012	Evaluation May 13, 2018.
	6013	Evaluation May 13, 2018.
	6014	Title May 13, 2018.
	6015	Title May 13, 2018.
	6017	Evaluation May 13, 2018.
	6018	Evaluation May 13, 2018.
	6019	Evaluation.
	6026	Evaluation May 13, 2018.
	6027	Evaluation May 13, 2018.
	6034	Evaluation May 13, 2018.
	6035 6036	Evaluation May 13, 2018.
	6040	Evaluation May 13, 2018. Added May 13, 2018.
	6042	Added May 13, 2018.
	6046	Added May 13, 2018.
	6091	Evaluation May 13, 2018.
l.84a		Table V criterion July 1, 1994.
	6010	Criterion March 11, 1969.
	6019	Criterion September 22, 1978.
	6029	NOTE August 23, 1948; criterion September 22, 1978.
	6035	Added September 9, 1975.
	6050-6062	Removed March 10, 1976.
	6061	Added March 10, 1976.
	6062	Added March 10, 1976.
	6063–6079 6064	Criterion September 22, 1978. Criterion March 10, 1976.
	6071	Criterion March 10, 1976.
	6076	Evaluation August 23, 1948.
	6080	Criterion September 22, 1978.
	6081	Criterion March 10, 1976.
	6090	Criterion September 22, 1978; criterion September 12, 1988.
1.84b	6260	Added October 1, 1961; criterion October 1, 1961; evaluation March 10, 1976
		removed December 18, 1987; re-designated § 4.87a December 18, 1987.
l.87		Tables VI and VII replaced by new Tables VI, VIA, and VII December 18, 198
		6200-6260 revised and re-designated § 4.87 June 10, 1999.
l.87a	6200–6260	Moved to §4.87 June 10, 1999.
	6275–6276	Moved from § 4.87b June 10, 1999.
	6277–6297	March 23, 1956 removed, December 17, 1987; Table II revised Table V March
		10, 1976; Table II revised to Table VII September 22, 1978; text from §4.84
	0000	Schedule of ratings-ear re-designated from §4.87 December 17, 1987.
	6286	Removed December 17, 1987.
	6291 6297	Criterion March 10, 1976; removed December 17, 1987. Criterion March 10, 1976; removed December 17, 1987.
I.87b		Removed June 10, 1999.
l.88a		March 11, 1969; re-designated § 4.88b November 29, 1994; § 4.88a added
r.00a		read "Chronic fatigue syndrome"; criterion November 29, 1994.
4.88b		Added March 11, 1969; re-designated §4.88c November 29, 1994; §4.88a re
		designated to §4.88b November 29, 1994; General Rating Formula for Infec
		tious Diseases added August 11, 2019.
	6300	Criterion August 30, 1996; title, criterion, and note August 11, 2019.
	6301	Criterion, note August 11, 2019.
	6302	Criterion September 22, 1978; criterion August 30, 1996; criterion, note August
	0002	Ontenon deplember 22, 1970, chienon August 30, 1930, chienon, note August

Sec.	Diagnostic code No.	
	6304 6305	Evaluation August 30, 1996; criterion, note August 11, 2019. Criterion March 1, 1989; evaluation August 30, 1996; title, criterion, note August 11, 2019.
	6306	Evaluation August 30, 1996; criterion, note August 11, 2019.
	6307	Criterion May 13, 2018; criterion, note August 11, 2019.
	6308 6309	Criterion August 30, 1996; criterion, note August 11, 2019.
	6310	Added March 1, 1963; criterion March 1, 1989; criterion August 30, 1996; cr terion, note August 11, 2019. Criterion, note August 11, 2019.
	6311	Criterion, note August 11, 2019.
	6312	Added August 11, 2019.
	6314	Evaluation March 1, 1989; evaluation August 30, 1996.
	6315 6316	Criterion August 30, 1996. Evaluation March 1, 1989; evaluation August 30, 1996; criterion, note August 1: 2019.
	6317	Criterion August 30, 1996; title, criterion, note August 11, 2019.
	6318	Added March 1, 1989; criterion August 30, 1996; criterion, note August 11, 2019
	6319	Added August 30, 1996; criterion, note August 11, 2019.
	6320	Added August 30, 1996; criterion, note August 11, 2019.
	6325 6326	Added August 11, 2019. Added August 11, 2019.
	6329	Added August 11, 2019.
	6330	Added August 11, 2019.
	6331	Added August 11, 2019.
	6333 6334	Added August 11, 2019. Added August 11, 2019.
	6335	Added August 11, 2019. Added August 11, 2019.
	6350	Evaluation March 1, 1963; evaluation March 10, 1976; evaluation August 3 1996.
	6351	Added March 1, 1989; evaluation March 24, 1992; criterion August 30, 1996; criterion, note August 11, 2019.
	6352 6353	Added March 1, 1989; removed March 24, 1992.
	6354	Added March 1, 1989; removed March 24, 1992. Added November 29, 1994; criterion August 30, 1996; title, criterion, note Augu 11, 2019.
4.88c		Re-designated from § 4.88b November 29, 1994.
1.89		Ratings for nonpulmonary TB December 1, 1949; criterion March 11, 1969.
1.97	6502 6504	Criterion October 7, 1996. Criterion October 7, 1996.
	6510-6514	Criterion October 7, 1996.
	6515	Criterion March 11, 1969.
	6516	Criterion October 7, 1996.
	6517 6518	Removed October 7, 1996. Criterion October 7, 1996.
	6519	Criterion October 7, 1996.
	6520	Criterion October 7, 1996.
	6521	Added October 7, 1996.
	6522	Added October 7, 1996.
	6523 6524	Added October 7, 1996. Added October 7, 1996.
	6600	Evaluation September 9, 1975; criterion October 7, 1996.
	6601	Criterion October 7, 1996.
	6602	
		Criterion September 9, 1975; criterion October 7, 1996.
	6603	Added September 9, 1975; criterion October 7, 1996.
	6603 6604	Added September 9, 1975; criterion October 7, 1996. Added October 7, 1996.
	6603	Added September 9, 1975; criterion October 7, 1996.
	6603 6604 6701 6702 6703	Added September 9, 1975; criterion October 7, 1996. Added October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996.
	6603 6604 6701 6702 6703 6704	Added September 9, 1975; criterion October 7, 1996. Added October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Subparagraph (1) following December 1, 1949; criterion March 11, 1969; criterio September 22, 1978.
	6603 6604 6701 6702 6703 6704	Added September 9, 1975; criterion October 7, 1996. Added October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Subparagraph (1) following December 1, 1949; criterion March 11, 1969; criterion September 22, 1978. Removed March 11, 1969.
	6603 6604 6701 6702 6703 6704	Added September 9, 1975; criterion October 7, 1996. Added October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Subparagraph (1) following December 1, 1949; criterion March 11, 1969; criterion September 22, 1978. Removed March 11, 1969. Added March 11, 1969; removed September 22, 1978. Criterion July 6, 1950; criterion September 22, 1978. Second note following December 1, 1949; criterion March 11, 1969; evaluation
	6603 6604 6701 6702 6703 6704 6705 6707–6710 6721 6724	Added September 9, 1975; criterion October 7, 1996. Added October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Subparagraph (1) following December 1, 1949; criterion March 11, 1969; criterion September 22, 1978. Removed March 11, 1969. Added March 11, 1969; removed September 22, 1978. Criterion July 6, 1950; criterion September 22, 1978. Second note following December 1, 1949; criterion March 11, 1969; evaluation October 7, 1996.
	6603 6604 6701 6702 6703 6704 6705 6707–6710 6721 6724 6725–6728	Added September 9, 1975; criterion October 7, 1996. Added October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Subparagraph (1) following December 1, 1949; criterion March 11, 1969; criterion September 22, 1978. Removed March 11, 1969; removed September 22, 1978. Criterion July 6, 1950; criterion September 22, 1978. Second note following December 1, 1949; criterion March 11, 1969; evaluation October 7, 1996. Added March 11, 1969; removed September 22, 1978.
	6603 6604 6701 6702 6703 6704 6705 6707–6710 6721 6724 6725–6728 6730	Added September 9, 1975; criterion October 7, 1996. Added October 7, 1996. Evaluation October 7, 1996. Subparagraph (1) following December 1, 1949; criterion March 11, 1969; criterion September 22, 1978. Removed March 11, 1969. Added March 11, 1969; removed September 22, 1978. Criterion July 6, 1950; criterion September 22, 1978. Second note following December 1, 1949; criterion March 11, 1969; evaluation October 7, 1996. Added March 11, 1969; removed September 22, 1978. Added September 22, 1978; criterion October 7, 1996.
	6603 6604 6701 6702 6703 6704 6705 6707-6710 6721 6724 6725-6728 6730 6731	Added September 9, 1975; criterion October 7, 1996. Added October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Subparagraph (1) following December 1, 1949; criterion March 11, 1969; criterion September 22, 1978. Removed March 11, 1969. Added March 11, 1969; removed September 22, 1978. Criterion July 6, 1950; criterion September 22, 1978. Second note following December 1, 1949; criterion March 11, 1969; evaluation October 7, 1996. Added March 11, 1969; removed September 22, 1978. Added September 22, 1978; criterion October 7, 1996. Evaluation September 22, 1978; criterion October 7, 1996.
	6603 6604 6701 6702 6703 6704 6705 6707–6710 6721 6724 6725–6728 6730	Added September 9, 1975; criterion October 7, 1996. Added October 7, 1996. Evaluation October 7, 1996. Subparagraph (1) following December 1, 1949; criterion March 11, 1969; criterion September 22, 1978. Removed March 11, 1969. Added March 11, 1969; removed September 22, 1978. Criterion July 6, 1950; criterion September 22, 1978. Second note following December 1, 1949; criterion March 11, 1969; evaluation October 7, 1996. Added March 11, 1969; removed September 22, 1978. Added September 22, 1978; criterion October 7, 1996.
	6603 6604 6701 6702 6703 6704 6705 6707–6710 6721 6724 6725–6728 6730 6731 6732	Added September 9, 1975; criterion October 7, 1996. Added October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Evaluation October 7, 1996. Subparagraph (1) following December 1, 1949; criterion March 11, 1969; criterion September 22, 1978. Removed March 11, 1969; removed September 22, 1978. Criterion July 6, 1950; criterion September 22, 1978. Criterion July 6, 1950; criterion September 22, 1978. Second note following December 1, 1949; criterion March 11, 1969; evaluation October 7, 1996. Added March 11, 1969; removed September 22, 1978. Added September 22, 1978; criterion October 7, 1996. Evaluation September 22, 1978; criterion October 7, 1996. Criterion March 11, 1969.

Sec.	Diagnostic code No.	
	6814	Criterion March 10, 1976; removed October 7, 1996.
	6815	Removed October 7, 1996.
	6816 6817	Removed October 7, 1996. Evaluation October 7, 1996.
	6818	Removed October 7, 1996.
	6819	Criterion March 10, 1976; criterion October 7, 1996.
	6821	Evaluation August 23, 1948.
l.104	6822–6847 7000	Added October 7, 1996. Evaluation July 6, 1950; evaluation September 22, 1978; evaluation January 12.
1.104	7000	1998. Evaluation January 12, 1998.
	7001	Evaluation January 12, 1998.
	7003	Evaluation January 12, 1998.
	7004	Criterion September 22, 1978; evaluation January 12, 1998.
	7005	Evaluation September 9, 1975; evaluation September 22, 1978; evaluation January 12, 1998.
	7006	Evaluation January 12, 1998.
	7007	Evaluation September 22, 1978; evaluation January 12, 1998.
	7008 7010	Evaluation January 12, 1998; criterion December 10, 2017. Evaluation January 12, 1998.
	7011	Evaluation January 12, 1998.
	7013	Removed January 12, 1998.
	7014	Removed January 12, 1998.
	7015	Evaluation September 9, 1975; criterion January 12, 1998.
	7016 7017	Added September 9, 1975; evaluation January 12, 1998. Added September 22, 1978; evaluation January 12, 1998.
	7017	Added January 12, 1998.
	7019	Added January 12, 1998.
	7020	Added January 12, 1998.
	7100 7101	Evaluation July 6, 1950.
		Criterion September 1, 1960; criterion September 9, 1975; criterion January 12 1998.
	7110 7111	Evaluation September 9, 1975; evaluation January 12, 1998.
	7112	Criterion September 9, 1975; evaluation January 12, 1998. Evaluation January 12, 1998.
	7113	Evaluation January 12, 1998.
	7114	Added June 9, 1952; evaluation January 12, 1998.
	7115	Added June 9, 1952; evaluation January 12, 1998.
	7116 7117	Added June 9, 1952; evaluation March 10, 1976; removed January 12, 1998. Added June 9, 1952; evaluation January 12, 1998.
	7118	Criterion January 12, 1998.
	7119	Evaluation January 12, 1998.
	7120	Note following July 6, 1950; evaluation January 12, 1998.
	7121 7122	Criterion July 6, 1950; evaluation March 10, 1976; evaluation January 12, 1998. Last sentence of Note following July 6, 1950; evaluation January 12, 1998; cri-
	7123	terion August 13, 1998. Added October 15, 1991; criterion January 12, 1998.
4.114	7304	Introduction paragraph revised March 10, 1976. Evaluation November 1, 1962.
	7305	Evaluation November 1, 1962.
	7308	Evaluation April 8, 1959.
	7311	Criterion July 2, 2001.
	7312	Evaluation March 10, 1976; evaluation July 2, 2001.
	7313 7319	Evaluation March 10, 1976; removed July 2, 2001. Evaluation November 1, 1962.
	7319	Evaluation July 6, 1950; criterion March 10, 1976.
	7328	Evaluation November 1, 1962.
	7329	Evaluation November 1, 1962.
	7330	Evaluation November 1, 1962.
	7331 7332	Criterion March 11, 1969. Evaluation November 1, 1962.
	7334	Evaluation July 6, 1950; evaluation November 1, 1962.
	7339	Criterion March 10, 1976.
	7341	Removed March 10, 1976.
	7343	Criterion March 10, 1976; criterion July 2, 2001.
	7344 7345	Criterion July 2, 2001. Evaluation August 23, 1948; evaluation February 17, 1955; evaluation July 2
	7346	2001. Evaluation February 1, 1962.
	7347	Added September 9, 1975.
	7348	Added March 10, 1976.
	7351 7354	Added July 2, 2001. Added July 2, 2001.

Sec.	Diagnostic code No.	
4.115a		Re-designated and revised as §4.115b; new §4.115a "Ratings of the genito- urinary system-dysfunctions" added February 17, 1994.
4.115b	7500	Note July 6, 1950; evaluation February 17, 1994, criterion September 8, 1994.
	7501	Evaluation February 17, 1994.
	7502	Evaluation February 17, 1994.
	7503	Removed February 17, 1994.
	7504 7505	Criterion February 17, 1994. Criterion March 11, 1969; evaluation February 17, 1994.
	7507	Criterion February 17, 1994.
	7508	Evaluation February 17, 1994.
	7509	Criterion February 17, 1994.
	7510	Evaluation February 17, 1994.
	7511 7512	Evaluation February 17, 1994.
	7512 7513	Evaluation February 17, 1994. Removed February 17, 1994.
	7514	Criterion March 11, 1969; removed February 17, 1994.
	7515	Criterion February 17, 1994.
	7516	Criterion February 17, 1994.
	7517	Criterion February 17, 1994.
	7518	Evaluation February 17, 1994.
	7519 7520	Evaluation March 10, 1976; evaluation February 17, 1994.
	7520 7521	Criterion February 17, 1994. Criterion February 17, 1994.
	7522	Criterion September 8, 1994.
	7523	Criterion September 8, 1994.
	7524	Note July 6, 1950; evaluation February 17, 1994; evaluation September 8, 1994.
	7525	Criterion March 11, 1969; evaluation February 17, 1994.
	7526	Removed February 17, 1994.
	7527 7528	Criterion February 17, 1994. Criterion March 10, 1976; criterion February 17, 1994.
	7529	Criterion February 17, 1994.
	7530	Added September 9, 1975; evaluation February 17, 1994.
	7531	Added September 9, 1975; criterion February 17, 1994.
	7532–7542	Added February 17, 1994.
4.116		§4.116 removed and §4.116a re-designated §4.116 "Schedule of ratings-gyne cological conditions and disorders of the breasts" May 22, 1995.
	7610	Criterion May 22, 1995; title May 13, 2018.
	7611 7612	Criterion May 22, 1995. Criterion May 22, 1995.
	7613	Criterion May 22, 1995.
	7614	Criterion May 22, 1995.
	7615	Criterion May 22, 1995; note May 13, 2018.
	7617	Criterion May 22, 1995.
	7618	Criterion May 22, 1995.
	7619 7620	Criterion May 22, 1995; note May 13, 2018.
	7621	Criterion May 22, 1995. Criterion May 22, 1995; evaluation May 13, 2018.
	7622	Removed May 13, 2018.
	7623	Removed May 13, 2018.
	7624	Criterion August 9, 1976; evaluation May 22, 1995.
	7625	Criterion August 9, 1976; evaluation May 22, 1995.
	7626 7627	Criterion May 22, 1995; criterion March 18, 2002. Criterion March 10, 1976; criterion May 22, 1995; title, note May 13, 2018.
	7628	Added May 22, 1995; title, criterion May 13, 2018.
	7629	Added May 22, 1995.
	7630	Added May 13, 2018.
	7631	Added May 13, 2018.
	7632	Added May 13, 2018.
1.117	7700	Removed December 9, 2018.
	7701 7702	Removed October 23, 1995.
	7702	Evaluation October 23, 1995; title December 9, 2018; evaluation December 9, 2018. Evaluation August 23, 1948; criterion October 23, 1995; evaluation December 9
	7703	2018; criterion December 9, 2018.
	7704	Evaluation October 23, 1995; evaluation December 9, 2018.
	7705	Evaluation October 23, 1995; title December 9, 2018; evaluation December 9 2018; criterion December 9, 2018.
	7706	Evaluation October 23, 1995; note December 9, 2018; criterion October 23 1995.
	7707	Criterion October 23, 1995.
	7709	Evaluation March 10, 1976; criterion October 23, 1995; title December 9, 2018
		criterion December 9, 2018.
	7710	Criterion October 23, 1995; criterion December 9, 2018.

Sec.	Diagnostic code No.	
	7711	Criterion October 23, 1995.
	7712	Added December 9, 2018.
	7713	Removed October 23, 1995.
	7714	Added September 9, 1975; criterion October 23, 1995; criterion December 9, 2018.
	7715	Added October 26, 1990; criterion December 9, 2018.
	7716	Added October 23, 1995; evaluation December 9, 2018; criterion December 9,
	7717	2018. Added March 9, 2012.
	7718	Added December 9, 2018.
	7719	Added December 9, 2018.
	7720	Added December 9, 2018.
	7721 7722	Added December 9, 2018. Added December 9, 2018.
	7723	Added December 9, 2018.
	7724	Added December 9, 2018.
	7725	Added December 9, 2018.
4.118	7800 7801	Evaluation August 30, 2002; criterion October 23, 2008.
	7001	Criterion July 6, 1950; criterion August 30, 2002; criterion October 23, 2008; title, note 1, note 2 August 13, 2018.
	7802	Criterion September 22, 1978; criterion August 30, 2002; criterion October 23,
		2008; title, note 1, note 2 August 13, 2018.
	7803	Criterion August 30, 2002; removed October 23, 2008.
	7804	Criterion July 6, 1950; criterion September 22, 1978; criterion and evaluation October 23, 2008.
	7805	Criterion October 23, 2008; title August 13, 2018.
		General Rating Formula for DCs 7806, 7809, 7813-7816, 7820-7822, and 7824
	7000	added August 13, 2018.
	7806	Criterion September 9, 1975; evaluation August 30, 2002; criterion August 13, 2018.
	7807	Criterion August 30, 2002.
	7808	Criterion August 30, 2002.
	7809	Criterion August 30, 2002; title, criterion August 13, 2018.
	7810 7811	Removed August 30, 2002. Criterion March 11, 1969; evaluation August 30, 2002.
	7812	Removed August 30, 2002.
	7813	Criterion August 30, 2002; title, criterion August 13, 2018.
	7814	Removed August 30, 2002.
	7815 7816	Evaluation August 30, 2002; criterion, note August 13, 2018. Evaluation August 30, 2002; criterion, note August 13, 2018.
	7817	Evaluation August 30, 2002; title, criterion, note August 13, 2018.
	7818	Criterion August 30, 2002.
	7819	Criterion August 30, 2002.
	7820 7821	Added August 30, 2002; criterion August 13, 2018. Added August 30, 2002; title, criterion August 13, 2018.
	7822	Added August 30, 2002; title, criterion August 13, 2018.
	7823	Added August 30, 2002; criterion August 13, 2018.
	7824	Added August 30, 2002; criterion August 13, 2018.
	7825	Added August 30, 2002; title, criterion August 13, 2018.
	7826 7827	Added August 30, 2002; criterion August 13, 2018. Added August 30, 2002; criterion August 13, 2018.
	7828	Added August 30, 2002; criterion August 13, 2018.
	7829	Added August 30, 2002; criterion August 13, 2018.
	7830	Added August 30, 2002; criterion August 13, 2018.
	7831 7832	Added August 30, 2002; criterion August 13, 2018. Added August 30, 2002; criterion August 13, 2018.
	7833	Added August 30, 2002; criterion August 13, 2018.
4.119	7900	Criterion August 13, 1981; evaluation June 9, 1996; title December 10, 2017;
		evaluation December 10, 2017; criterion December 10, 2017; note December
	7901	10, 2017.
	7901	Criterion August 13, 1981; evaluation June 9, 1996; title December 10, 2017; evaluation December 10, 2017; criterion December 10, 2017; note December
		10, 2017.
	7902	Evaluation August 13, 1981; criterion June 9, 1996; title December 10, 2017;
		evaluation December 10, 2017; criterion December 10, 2017; note December
	7903	10, 2017. Criterion August 13, 1981; evaluation June 9, 1996; evaluation December 10,
	7300	2017; criterion December 10, 2017; note December 10, 2017.
	7904	Criterion August 13, 1981; evaluation June 9, 1996; evaluation December 10,
	7005	2017; criterion December 10, 2017; note <i>December 10, 2017</i> .
	7905	Evaluation; August 13, 1981; evaluation June 9, 1996; evaluation December 10, 2017; criterion December 10, 2017.
	7906	Added December 10, 2017.
		-, -

Sec.	Diagnostic code No.	
	7907	Evaluation; August 13, 1981; evaluation June 9, 1996; criterion December 10, 2017; note December 10, 2017.
	7908	Criterion August 13, 1981; criterion June 9, 1996; criterion December 10, 2017.
	7909	Evaluation August 13, 1981; criterion June 9, 1996; evaluation June 9, 1996; criterion December 10, 2017; evaluation December 10, 2017; note December 10,
	7910	2017. Removed June 9, 1996.
	7911	Evaluation March 11, 1969; evaluation August 13, 1981; criterion June 9, 1996;
		title December 10, 2017; note December 10, 2017.
	7912 7913	Title December 10, 2017; criterion December 10, 2017.
	7913	Criterion September 9, 1975; criterion August 13, 1981; criterion June 6, 1996; evaluation June 9, 1996; criterion December 10, 2017; note <i>December 10</i> , 2017.
	7914	Criterion March 10, 1976; criterion August 13, 1981; criterion June 9, 1996.
	7915 7916	Criterion June 9, 1996; criterion December 10, 2017.
	7916 7917	Added June 9, 1996; note <i>December 10, 2017</i> . Added June 9, 1996; note <i>December 10, 2017</i> .
	7918	Added June 9, 1996; note <i>December 10, 2017.</i>
	7919	Added June 9, 1996; evaluation June 9, 1996; criterion December 10, 2017; note December 10, 2017.
4.124a	8002 8021	Criterion September 22, 1978. Criterion September 22, 1978; criterion October 1, 1961; criterion March 10,
	0021	1976; criterion March 1, 1989.
	8045	Criterion and evaluation October 23, 2008.
	8046	Added October 1, 1961; criterion March 10, 1976; criterion March 1, 1989.
	8100 8540	Evaluation June 9, 1953. Added October 15, 1991.
	8910	Added October 1, 1961.
	8911	Added October 1, 1961; evaluation September 9, 1975.
	8912 8913	Added October 1, 1961. Added October 1, 1961.
	8914	Added October 1, 1961; criterion September 9, 1975; criterion March 10, 1976.
	8910-8914	Evaluations September 9, 1975.
4.125—4.132		All Diagnostic Codes under Mental Disorders October 1, 1961; except as to eval-
4.130		uation for Diagnostic Codes 9500 through 9511 September 9, 1975. Re-designated from § 4.132 November 7, 1996.
	9200	Removed February 3, 1988.
	9201	Criterion February 3, 1988; Title August 4, 2014.
	9202 9203	Criterion February 3, 1988; removed August 4, 2014. Criterion February 3, 1988; removed August 4, 2014.
	9204	Criterion February 3, 1988; removed August 4, 2014.
	9205	Criterion February 3, 1988; criterion November 7, 1996; Removed August 4, 2014.
	9206 9207	Criterion February 3, 1988; removed November 7, 1996. Criterion February 3, 1988; removed November 7, 1996.
	9208	Criterion February 3, 1988; removed November 7, 1996.
	9209	Criterion March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9210 9211	Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996; Title August 4, 2014. Added November 7, 1996.
	9300	Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996.
	9301	Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996; Title August 4, 2014.
	9302 9303	Criterion March 10, 1976; criterion February 3, 1988; removed November 7, 1996. Criterion March 10, 1976; criterion February 3, 1988; removed November 7,
	9304	1996. Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996;
	9305	Title August 4, 2014. Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996;
	9306	Title August 4, 2014. Criterion March 10, 1976; criterion February 3, 1988; removed November 7,
	9307	1996. Criterion March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9308	Criterion March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9309	Criterion March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9310	Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996; Title August 4, 2014.
	9311	Criterion March 10, 1976; criterion February 3, 1988; removed November 7,

Sec.	Diagnostic code No.	
	9312	Added March 10, 1976; criterion February 3, 1988; criterion November 7, 1996;
	9313	Title August 4, 2014.
	9314	Added March 10, 1976; removed February 3, 1988. Added March 10, 1976; removed February 3, 1988.
	9315	Added March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9316–9321	Added March 10, 1976; removed February 3, 1988.
	9322 9323	Added March 10, 1976; criterion February 3, 1988; removed November 7, 1996. Added March 10, 1976; removed February 3, 1988.
	9324	Added March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9325	Added March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9326	Added March 10, 1976; removed February 3, 1988; added November 7, 1996; Title August 4, 2014.
	9327 9400–9411	Added November 7, 1996; removed August 4, 2014. Evaluations February 3, 1988.
	9400	Criterion March 10, 1976; criterion February 3, 1988.
	9401	Criterion March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9402	Criterion March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9403 9410	Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996; Title August 4, 2014.
	9411	Added March 10, 1976; criterion February 3, 1988; Title August 4, 2014. Added February 3, 1988.
	9412	Added November 7, 1996.
	9413	Added November 7, 1996; Title August 4, 2014.
	9416 9417	Added November 7, 1996; Title August 4, 2014. Added November 7, 1996; Title August 4, 2014.
	9421	Added November 7, 1996; Title August 4, 2014.
	9422	Added November 7, 1996; Title August 4, 2014.
	9423 9424	Added November 7, 1996; Title August 4, 2014. Added November 7, 1996; Title August 4, 2014.
	9425	Added November 7, 1996; Title August 4, 2014.
	9431	Added November 7, 1996.
	9432 9433	Added November 7, 1996. Added November 7, 1996; Title August 4, 2014.
	9434	Added November 7, 1996, Title Adgust 4, 2014.
	9435	Added November 7, 1996; Title August 4, 2014.
	9440 9500	Added November 7, 1996. Criterion March 10, 1976; criterion February 3, 1988.
	9501	Criterion March 10, 1976; Criterion February 3, 1988.
	9502	Criterion March 10, 1976; criterion February 3, 1988.
	9503	Removed March 10, 1976.
	9504 9505	Criterion September 9, 1975; removed March 10, 1976. Added March 10, 1976; criterion February 3, 1988.
	9506	Added March 10, 1976; criterion February 3, 1988.
	9507	Added March 10, 1976; criterion February 3, 1988.
	9508 9509	Added March 10, 1976; criterion February 3, 1988.
	9510	Added March 10, 1976; criterion February 3, 1988. Added March 10, 1976; criterion February 3, 1988.
	9511	Added March 10, 1976; criterion February 3, 1988.
	9520 9521	Added November 7, 1996. Added November 7, 1996.
4.132	9521	Re-designated as § 4.130 November 7, 1996.
4.150	9900	Criterion September 22, 1978; criterion February 17, 1994; title September 10, 2017.
	9901 9902	Criterion February 17, 1994. Criterion February 17, 1994; evaluation September 10, 2017; title September 10, 2017.
	9903	2017. Criterion February 17, 1994; evaluation September 10, 2017; title September 10, 2017.
	9904	Criterion September 10, 2017.
	9905	Criterion September 22, 1978; evaluation February 17, 1994; evaluation September 10, 2017; title September 10, 2017.
	9906 9907	Removed September 10, 2017. Removed September 10, 2017.
	9910	Removed September 10, 2017. Removed February 17, 1994.
	9911	Criterion and title September 10, 2017.
	9912	Removed September 10, 2017.
	9913 9914	Criterion February 17, 1994. Added February 17, 1994.
		Criterion February 17, 1994. Added February 17, 1994. Added February 17, 1994. Added February 17, 1994; criterion September 10, 2017.

Sec.	Diagnostic code No.	
	9918	Added September 10, 2017.

[72 FR 12983, Mar. 20, 2007; 72 FR 16728, Apr. 5, 2007, as amended at 73 FR 54708, 54711, Sept. 23, 2008; 73 FR 69554, Nov. 19, 2008; 77 FR 6467, Feb. 8, 2012; 79 FR 45101, Aug. 4, 2014; 80 FR 42042, July 16, 2015; 82 FR 36084, Aug. 3, 2017; 82 FR 50806, Nov. 2, 2017; 83 FR 15072, Apr. 9, 2018; 83 FR 15323, Apr. 10, 2018; 83 FR 32600, July 13, 2018; 83 FR 54257, Oct. 29, 2018; 84 FR 28233, June 18, 2019; 85 FR 76464, Nov. 30, 2020; 86 FR 8143, Feb. 4, 2021]

APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES

Diagnostic Code No.		
THE MUSCULOSKELETAL SYSTEM Acute, Subacute, or Chronic Diseases		
5000	Osteomyelitis, acute, subacute, or chronic.	
5001	Bones and Joints, tuberculosis.	
5002	Multi-joint arthritis (except post-traumatic and gout), 2 or more joints, as an active process.	
003	Degenerative arthritis, other than post-traumatic.	
6004	Arthritis, gonorrheal.	
005	Arthritis, pneumococcic.	
6006 6007	Arthritis, typhoid.	
5008	Arthritis, syphilitic. Arthritis, streptococcic.	
009	Other specified forms of arthropathy (excluding gout).	
010	Post-traumatic arthritis.	
011		
012	Bones, neoplasm, malignant, primary or secondary.	
013	Osteoporosis, residuals of.	
014		
015	Bones, neoplasm, benign.	
016	Osteitis deformans.	
017	Gout.	
018	[Removed]	
019	Bursitis.	
020	[Removed]	
021	Myositis.	
022		
023		
024	Tenosynovitis, tendinitis, tendinosis or tendinopathy.	
025	Fibromyalgia.	
	Prosthetic Implants	
5051	Shoulder replacement (prosthesis).	
5052	Elbow replacement (prosthesis).	
053	Wrist replacement (prosthesis).	
6054	Hip, resurfacing or replacement (prosthesis).	
5055	Knee, resurfacing or replacement (prosthesis).	
5056	Ankle replacement (prosthesis).	
	Combination of Disabilities	
i104	Anatomical loss of one hand and loss of use of one foot.	
105	Anatomical loss of one foot and loss of use of one hand.	
106	Anatomical loss of both hands.	
107	Anatomical loss of both feet.	
108	Anatomical loss of one hand and one foot.	
5109	Loss of use of both hands.	
5110	Loss of use of both feet.	
5111	Loss of use of one hand and one foot.	
	Amputations: Upper Extremity	
Arm amputation of:		
5120	Complete amputation, upper extremity.	
5121	Above insertion of deltoid.	
5122	Below insertion of deltoid.	
Forearm amputation of:		
oroam amputation or.		
5123	Above insertion of pronator teres.	

Diagnostic Code No.	
5125	Hand, loss of use of.
5125	
	Multiple Finger Amputations
5126	Five digits of one hand.
Four digits of one hand:	
5127	Thumb, index, long and ring.
5128 5129	Thumb, index, long and little. Thumb, index, ring and little.
5130	Thumb, long, ring and little.
5131	Index, long, ring and little.
Three digits of one hand:	
5132	Thumb, index and long.
5133	Thumb, index and ring.
5134	Thumb, index and little.
5135	Thumb, long and ring.
5136	Thumb, long and little.
5137	Thumb, ring and little.
5138	Index, long and ring.
5139 5140	Index, long and little. Index, ring and little.
5140	Long, ring and little.
-	Long, ring and ittle.
Two digits of one hand:	I Thomas and today
5142	Thumb and index.
5143	Thumb and long.
5144 5145	Thumb and ring.
	Thumb and little.
5146 5147	Index and long. Index and ring.
5148	Index and little.
5149	Long and ring.
5150	Long and little.
5151	Ring and little.
0: 1 "	
Single finger:	1. Though
5152	Thumb.
5153 5154	Index finger.
5155	Long finger. Ring finger.
5156	Little finger.
	Amputations: Lower Extremity
Thigh amputation of:	
5160	Complete amputation, lower extremity.
5161 5162	Upper third. Middle or lower thirds.
	Mildle of lower tillus.
Leg amputation of: 5163	With defeative atumn
5164	With defective stump.
5165	Not improvable by prosthesis controlled by natural knee action. At a lower level, permitting prosthesis.
5166	Forefoot, proximal to metatarsal bones.
5167	Foot, loss of use of.
5170	Toes, all, amputation of, without metatarsal loss or transmetatarsal, amputation of, with up to
0170	half of metatarsal loss.
5171	Toe, great.
5172	Toes, other than great, with removal of metatarsal head.
5173	Toes, three or more, without metatarsal involvement.
	Shoulder and Arm
5200	Scapulohumeral articulation, ankylosis.
5201	Arm, limitation of motion.
5202	Humerus, other impairment.
5203	Clavicle or scapula, impairment.
	Elbow and Forearm
5205	Elbow, ankylosis.
5206	Forearm, limitation of flexion.
5207	Forearm, limitation of extension.

38 CFR Ch. I (7-1-21 Edition)

Diagnostic Octobal	
Diagnostic Code No.	
5208	Forearm, flexion limited.
5209	Elbow, other impairment.
5210	Radius and ulna, nonunion.
5211 5212	Ulna, impairment. Radius, impairment.
5213	Supination and pronation, impairment.
	Wrist
5214 5215	Wrist, ankylosis. Wrist, limitation of motion.
	Limitation of Motion
Multiple Digits: Unfavorable Anky	
5216 5217	Five digits of one hand. Four digits of one hand.
5218	Three digits of one hand.
5219	Two digits of one hand.
Multiple Digits: Favorable Anky-	
losis:	
5220	Five digits of one hand.
5221 5222	Four digits of one hand. Three digits of one hand.
5223	Two digits of one hand.
-	
Ankylosis of Individual Digits: 5224	Thumb.
5225	Index finger.
5226	Long finger.
5227	Ring or little finger.
Limitation of Motion of Individual	Digits:
5228	Thumb.
5229	Index or long finger.
5230	Ring or little finger.
	Spine
5235	Vertebral fracture or dislocation.
5236	Sacroiliac injury and weakness.
5237	Lumbosacral or cervical strain.
5238 5239	Spinal stenosis. Spondylolisthesis or segmental instability.
5240	Ankylosing spondylitis.
5241	Spinal fusion.
5242	Degenerative arthritis, degenerative disc disease other than intervertebral disc syndrome (also
	see either DC 5003 or 5010).
5243 5244	Intervertebral disc syndrome. Traumatic paralysis, complete.
<u></u>	
	Hip and Thigh
5250	Hip, ankylosis.
5251	Thigh, limitation of extension.
5252	Thigh, limitation of flexion.
5253	Thigh, impairment.
5254 5255	Hip, flail joint. Femur, impairment.
	Knee and Leg
5256	Knee, ankylosis.
5257	Knee, other impairment.
5258	Cartilage, semilunar, dislocated.
5259 5260	Cartilage, semilunar, removal. Leg, limitation of flexion.
5261	Leg, limitation of flexion. Leg, limitation of extension.
5262	Tibia and fibula, impairment.
5263	Genu recurvatum.
	Ankle
5270	Ankle, ankylosis.

Diagnostic Code No.		
5271	Ankle, limited motion.	
5272	Subastragalar or tarsal joint, ankylosis.	
5273	Os calcis or astragalus, malunion.	
5274	Astragalectomy.	
	Shortening of the Lower Extremity	
	<u>`</u>	
5275	Bones, of the lower extremity	
	The Foot	
5269	Plantar fasciitis.	
5276	Flatfoot, acquired.	
5277	Weak foot, bilateral.	
5278 5279	Claw foot (pes cavus), acquired. Metatarsalqia, anterior (Morton's disease).	
5280	Hallux valgus.	
5281	Hallux rigidus.	
5282	Hammer toe.	
5283	Tarsal or metatarsal bones.	
5284	Foot injuries, other.	
	The Skull	
5296	Loss of part of.	
	The Ribs	
5297	Removal of.	
	The Coccyx	
5298	Removal of.	
	MUSCLE IN HIDIES	
	MUSCLE INJURIES Shoulder Girdle and Arm	
5301	Group I Function: Upward rotation of scapula.	
5302	Group II Function: Depression of arm.	
5303	Group III Function: Elevation and abduction of arm.	
5304	Group IV Function: Stabilization of shoulder.	
5305 5306	Group V Function: Elbow supination. Group VI Function: Extension of elbow.	
	<u>'</u>	
	Forearm and Hand	
5307	Group VII Function: Flexion of wrist and fingers.	
5308	Group VIII Function: Extension of wrist, fingers, thumb.	
5309	Group IX Function: Forearm muscles.	
	Foot and Leg	
5310	Group X Function: Movement of forefoot and toes.	
5311	Group XI Function: Propulsion of foot.	
5312	Group XII Function: Dorsiflexion.	
	Pelvic Girdle and Thigh	
5313	Group XIII Function: Extension of hip and flexion of knee.	
5314	Group XIV Function: Extension of knee.	
5315	Group XV Function: Adduction of hip.	
5316	Group XVII Function: Flexion of hip.	
5317 5318	Group XVII Function: Extension of hip. Group XVIII Function: Outward rotation of thigh.	
	Torso and Neck	
5319	Group XIX Function: Abdominal wall and lower thorax.	
5320 5321	Group XX Function: Postural support of body. Group XXI Function: Respiration.	
5322	Group XXII Function: Respiration. Group XXII Function: Rotary and forward movements, head.	
5323	Group XXIII Function: Movements of head.	
	Miscellaneous	
5324	Diaphragm, rupture.	

38 CFR Ch. I (7-1-21 Edition)

Muscle injury, facial muscles.
Muscle hernia.
Muscle, neoplasm of, malignant.
Muscle, neoplasm of, benign.
Sarcoma, soft tissue.
Rhabdomyolysis, residuals of.
Compartment syndrome.
THE EYE Diseases of the Eye
Choroidopathy, including uveitis, iritis, cyclitis, or choroiditis.
Keratopathy.
Scleritis.
Iritis.
Cyclitis Choroiditis.
Choroiditis Retinopathy or maculopathy not otherwise specified.
Intraocular hemorrhage.
Detachment of retina.
Unhealed eye injury.
Tuberculosis of eye.
Retinal scars, atrophy, or irregularities.
Angle-closure glaucoma.
Open-angle glaucoma. Malignant popularms of the over orbit, and adnova (evaluding skin)
 Malignant neoplasms of the eye, orbit, and adnexa (excluding skin). Benign neoplasms of the eye, orbit, and adnexa (excluding skin).
Nystagmus, central.
Conjunctivitis, trachomatous, chronic.
Conjunctivitis, other, chronic.
Ptosis unilateral or bilateral.
Ectropion.
Entropion.
Lagophthalmos. Evebrows, loss,
Eyebrows, loss. Eyelashes, loss.
Disorders of the lacrimal apparatus (epiphora, dacrocystitis, etc.).
Optic neuropathy.
Cataract.
Cataract, senile, and others.
Aphakia.
Accommodation, paralysis Dacryocystitis.
Eyelids, loss of portion.
Lens, crystalline, dislocation.
Pterygium.
Keratoconus.
Status post corneal transplant.
Diabetic retinopathy.
Retinal dystrophy (including retinitis pigmentosa, wet or dry macular degeneration, early-onse
macular degeneration, rod and/or cone dystrophy) Post-chiasmal disorders.
Impairment of Central Visual Acuity
Anatomical loss both eyes.
Blindness, both eyes, only light perception.
Other eye 5/200 (1.5/60).
Other eye 10/200 (3/60).
Other eye 15/200 (4.5/60).
Other eye 20/200 (6/60).
Other eye 20/100 (6/30).
Other eye 20/70 (6/21).
Other eye 20/50 (6/15). Other eye 20/40 (6/12).
Outer eye 20/40 (0/12).
perception:
Other eye 5/200 (1.5/60).
Other eye 5/200 (1.5/60). Other eye 10/200 (3/60).
Other eye 5/200 (1.5/60).

Diagnostic Code No.	
6069	Other eye 20/70 (6/21).
6069	Other eye 20/50 (6/15).
6070	Other eye 20/40 (6/12).
Vision in 1 eye 5/200 (1.5/60):	
6071	Other eye 5/200 (1.5/60).
6072	Other eye 10/200 (3/60).
6072	Other eye 15/200 (4.5/60).
6072	Other eye 20/200 (6/60).
6073	Other eye 20/100 (6/30).
6073	Other eye 20/70 (6/21).
6073	Other eye 20/50 (6/15).
6074	Other eye 20/40 (6/12).
Vision in 1 eye 10/200 (3/60):	
6075	Other eye 10/200 (3/60).
6075	Other eye 15/200 (4.5/60).
6075	Other eye 20/200 (6/60).
6076	Other eye 20/100 (6/30).
6076	Other eye 20/70 (6/21).
6076 6077	Other eye 20/50 (6/15). Other eye 20/40 (6/12).
0077	Other eye 20/40 (0/12).
Vision in 1 eye 15/200 (4.5/60):	
6075	Other eye 15/200 (4.5/60).
6075	Other eye 20/200 (6/60).
6076	Other eye 20/100 (6/30).
6076	Other eye 20/70 (6/21).
6076	Other eye 20/50 (6/15). Other eye 20/40 (6/12).
6077	Other eye 20/40 (6/12).
Vision in 1 eye 20/200 (6/60):	
6075	Other eye 20/200 (6/60).
6076	Other eye 20/100 (6/30).
6076	Other eye 20/70 (6/21).
6076	Other eye 20/50 (6/15).
6077	Other eye 20/40 (6/12).
Vision in 1 eye 20/100 (6/30):	
6078	Other eve 20/100 (6/30).
6078 6078	Other eye 20/100 (6/30). Other eye 20/70 (6/21).
6078	Other eye 20/70 (6/21).
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15).
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12).
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21).
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15).
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21).
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12).
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/50 (6/15).
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12).
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/50 (6/15).
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/50 (6/15).
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/50 (6/15). Other eye 20/50 (6/15). Other eye 20/40 (6/12).
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Field vision, impairment.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Field vision, impairment. Scotoma.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Field vision, impairment. Scotoma.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Field vision, impairment. Scotoma. Diplopia. Symblepharon.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Field vision, impairment. Scotoma. Diplopia. Symblepharon. Diplopia, limited muscle function.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Field vision, impairment. Scotoma. Diplopia. Symblepharon.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Field vision, impairment. Scotoma. Diplopia. Symblepharon. Diplopia, limited muscle function. THE EAR
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Field vision, impairment. Scotoma. Diplopia. Symblepharon. Diplopia, limited muscle function. THE EAR Chronic suppurative otitis media.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Field vision, impairment. Scotoma. Diplopia. Symblepharon. Diplopia, limited muscle function. THE EAR Chronic suppurative otitis media. Chronic nonsuppurative otitis media.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Field vision, impairment. Scotoma. Diplopia. Symblepharon. Diplopia, limited muscle function. THE EAR Chronic suppurative otitis media.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Field vision, impairment. Scotoma. Diplopia. Symblepharon. Diplopia, limited muscle function. THE EAR Chronic suppurative otitis media. Otosclerosis.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Field vision, impairment. Scotoma. Diplopia. Symblepharon. Diplopia, limited muscle function. THE EAR Chronic suppurative otitis media. Chronic nonsuppurative otitis media. Otosclerosis. Peripheral vestibular disorders.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Field vision, impairment. Scotoma. Diplopia. Symblepharon. Diplopia, limited muscle function. THE EAR Chronic suppurative otitis media. Chronic nonsuppurative otitis media. Otosclerosis. Peripheral vestibular disorders. Meniere's syndrome.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Field vision, impairment. Scotoma. Diplopia. Symblepharon. Diplopia, limited muscle function. THE EAR Chronic suppurative otitis media. Chronic nonsuppurative otitis media. Otosclerosis. Peripheral vestibular disorders. Meniere's syndrome. Loss of auricle. Malignant neoplasm. Benign neoplasm.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Field vision, impairment. Scotoma. Field vision, impairment. Scotoma. Diplopia. Symblepharon. Diplopia, limited muscle function. THE EAR Chronic suppurative otitis media. Chronic nonsuppurative otitis media. Otosclerosis. Peripheral vestibular disorders. Meniere's syndrome. Loss of auricle. Malignant neoplasm. Benign neoplasm. Benign neoplasm. Chronic otitis externa.
6078	Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/70 (6/21). Other eye 20/50 (6/15). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Other eye 20/40 (6/12). Field vision, impairment. Scotoma. Diplopia. Symblepharon. Diplopia, limited muscle function. THE EAR Chronic suppurative otitis media. Chronic nonsuppurative otitis media. Otosclerosis. Peripheral vestibular disorders. Meniere's syndrome. Loss of auricle. Malignant neoplasm. Benign neoplasm.

Diagnostic Code No.	
	OTHER SENSE ORGANS
275	Smell, complete loss.
276	
INFECTIO	DUS DISEASES, IMMUNE DISORDERS AND NUTRITIONAL DEFICIENCIES
300	·
301	
302	
304	Malaria.
305	
3306	
307	1 3 3
308	
3309 3310	
311	
312	
313	
314	Beriberi.
315	
316	
3317	
3318	
6319 6320	
3325	
326	
6329	
330	
3331	
333	Nontyphoid salmonella infections.
334	
335	
3350	
6351 6354	
0304	Chronic Fatigue Syndrome (CFS).
	THE RESPIRATORY SYSTEM Nose and Throat
2500	Ourhorn march desiration
6502 6504	
5510	
5511	
5512	
513	
514	
515	Laryngitis, tuberculous.
516	
5518	
5519	
5520	
3521 3522	1 , ,
5523	
5524	
70L-1	
	Trachea and Bronchi
600	Bronchitis, chronic.
6601	
6602	
6603 6604	
0004	<u> </u>
	Lungs and Pleura Tuberculosis
Ratings for Pulmonary Tubercu	ılosis (Chronic) Entitled on August 19, 1968: Active, far advanced.
6702 6703 6704	Active, minimal.

Diagnostic Code No.		
6722	Inactive, moderately advanced.	
6723	Inactive, minimal.	
6724	Inactive, advancement unspecified.	
	<u> </u>	
	sis Initially Evaluated After August 19, 1968:	
6730	Chronic, active.	
6731	Chronic, inactive.	
6732	Pleurisy, active or inactive.	
	Nontuberculous Diseases	
6817	Pulmonary Vascular Disease.	
6819	Neoplasms, malignant.	
6820	Neoplasms, benign.	
	Troopidanie, zanigni	
	Bacterial Infections of the Lung	
6822	Actinomycocio	
6823	Actinomycosis.	
6824	Nocardiosis. Chronic lung abscess.	
0024	Official lung abscess.	
	Interstitial Lung Disease	
	Fibuseis of lune diffuse intenstities	
6825	Fibrosis of lung, diffuse interstitial.	
6826	Desquamative interstitial pneumonitis.	
6827 6828	Pulmonary alveolar proteinosis.	
6829	Eosinophilic granuloma.	
6830	Drug-induced, pneumonitis & fibrosis.	
	Radiation-induced, pneumonitis & fibrosis.	
6831 6832	Hypersensitivity pneumonitis. Pneumoconiosis.	
6833	Asbestosis.	
	Asuesiosis.	
	Mycotic Lung Disease	
6834	Histoplasmosis.	
6835	Coccidioidomycosis.	
6836	Blastomycosis.	
6837	Cryptococcosis.	
6838	Aspergillosis.	
6839	Mucormycosis.	
	Postulation Lawre Diseases	
	Restrictive Lung Disease	
6840	Diaphragm paralysis or paresis.	
6841	Spinal cord injury with respiratory insufficiency.	
6842	Kyphoscoliosis, pectus excavatum/carinatum.	
6843	Traumatic chest wall defect.	
6844	Post-surgical residual.	
6845	Pleural effusion or fibrosis.	
6846 6847	Sarcoidosis. Sleep Apnea Syndromes.	
0047	Gleep Aprilea Syndionies.	
	THE CARDIOVASCULAR SYSTEM Diseases of the Heart	
7000	Valuulas kaast slisaasa	
7000	Valvular heart disease.	
7001	Endocarditis. Pericarditis.	
7002	Pericardial adhesions.	
7004		
7005	Syphilitic heart disease. Arteriosclerotic heart disease.	
7006	Myocardial infarction.	
7007	Hypertensive heart disease.	
7008	Hyperthyroid heart disease.	
7010	Supraventricular arrhythmias.	
7011	Ventricular arrhythmias.	
7015	Atrioventricular block.	
7016	Heart valve replacement.	
7017	Coronary bypass surgery.	
7018	Implantable cardiac pacemakers.	
7019	Cardiac transplantation.	
7020	Cardiomyopathy.	

Diagnostic Code No.		
	Diseases of the Arteries and Veins	
7101	Hypertensive vascular disease.	
7110	Aortic aneurysm.	
7111	Aneurysm, large artery.	
7112	Aneurysm, small artery. Arteriovenous fistula, traumatic.	
7114	Arteriosclerosis obliterans.	
7115	Thrombo-angiitis obliterans (Buerger's Disease).	
7117	Raynaud's syndrome.	
7118	Angioneurotic edema.	
7119 7120	Erythromelalgia. Varicose veins.	
7121	Post-phlebitic syndrome.	
7122	Cold injury residuals.	
7123	Soft tissue sarcoma.	
	THE DIGESTIVE SYSTEM	
7200	Mouth, injuries.	
7201	Lips, injuries.	
7202	Tongue, loss.	
7203	Esophagus, stricture.	
7204 7205	Esophagus, spasm. Esophagus, diverticulum.	
7301	Peritoneum, adhesions.	
7304	Ulcer, gastric.	
7305	Ulcer, duodenal.	
7306	Ulcer, marginal.	
7307	Gastritis, hypertrophic.	
7308 7309	Postgastrectomy syndromes. Stomach, stenosis.	
7310	Stomach, injury of, residuals.	
7311	Liver, injury of, residuals.	
7312	Liver, cirrhosis.	
7314	Cholecystitis, chronic.	
7315 7316	Cholelithiasis, chronic.	
7317	Cholangitis, chronic. Gall bladder, injury.	
7318	Gall bladder, removal.	
7319	Colon, irritable syndrome.	
7321	Amebiasis.	
7322	Dysentery, bacillary.	
7323 7324	Colitis, ulcerative.	
7325	Distomiasis, intestinal or hepatic. Enteritis, chronic.	
7326	Enterocolitis, chronic.	
7327	Diverticulitis.	
7328	Intestine, small, resection.	
7329	Intestine, large, resection.	
7330 7331	Intestine, fistula. Peritonitis.	
7332	Rectum & anus, impairment.	
7333	Rectum & anus, stricture.	
7334	Rectum, prolapse.	
7335	Ano, fistula in.	
7336 7337	Hemorrhoids. Pruritus ani.	
7338	Hernia, inguinal.	
7339	Hernia, ventral, postoperative.	
7340	Hernia, femoral.	
7342	Visceroptosis.	
7343	Neoplasms, malignant. Neoplasms, benign.	
7344 7345	Liver disease, chronic, without cirrhosis.	
7346	Hernia, hiatal.	
7347	Pancreatitis.	
7348	Vagotomy.	
7351	Liver transplant.	
7354	Hepatitis C.	
	THE GENITOURINARY SYSTEM	
7500	500 Kidney, removal.	

7501	Kidney, abscess.
7502	Nephritis, chronic.
7504	Pyelonephritis, chronic.
7505	Kidney, tuberculosis.
7507	Nephrosclerosis, arteriolar.
7508	Nephrolithiasis.
7509	Hydronephrosis.
7510	Ureterolithiasis.
7511	Ureter, stricture.
7512 7515	Cystitis, chronic. Bladder, calculus.
7516	Bladder, fistula.
7517	Bladder, injury.
7518	Urethra, stricture.
7519	Urethra, fistula.
7520	Penis, removal of half or more.
7521	Penis, removal of glans.
7522	Penis, deformity, with loss of erectile power.
7523	Testis, atrophy, complete.
7524	Testis, removal.
7525	Epididymo-orchitis, chronic only.
7527 7528	Prostate gland.
7528 7529	Malignant neoplasms. Benign neoplasms.
7530	Renal disease, chronic.
7531	Kidney transplant.
7532	Renal tubular disorders.
7533	Kidneys, cystic diseases.
7534	Atherosclerotic renal disease.
7535	Toxic nephropathy.
7536	Glomerulonephritis.
7537	Interstitial nephritis.
7538	Papillary necrosis.
7539	Renal amyloid disease.
7540	Disseminated intravascular coagulation. Renal involvement in systemic diseases.
7541 7542	Neurogenic bladder.
Gir	IECOLOGICAL CONDITIONS AND DISORDERS OF THE BREAST
7610	Vulva or clitoris, disease or injury of (including vulvovaginitis).
7611	Vagina, disease or injury.
7612	Cervix, disease or injury.
7613 7614	Uterus, disease or injury. Fallopian tube, disease or injury.
7615	Ovary, disease or injury.
7617	Uterus and both ovaries, removal.
7617 7618	Uterus and both ovaries, removal. Uterus, removal.
7617 7618 7619	Uterus and both ovaries, removal.
7617 7618 7619	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal.
7617	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy.
7617 7618	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal.
7617 7618	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal.
7617	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Breast, surgery.
7617	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system.
7617 7618	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system.
7617 7618 7619 7620 7624 7625 7626 7627 7628	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system. Endometriosis.
7617 7618 7619 7620 7621 7624 7625 7626 7627 7628 7629 7630	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system. Endometriosis. Malignant neoplasms of the breast.
7617 7618 7619 7620 7621 7624 7625 7626 7627 7628 7629 7630	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complicitions of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system. Endometriosis. Malignant neoplasms of the breast. Benign neoplasms of the breast and other injuries of the breast.
7617 7618	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system. Endometriosis. Malignant neoplasms of the breast. Benign neoplasms of the breast. Benign neoplasms of the breast and other injuries of the breast. Female sexual arousal disorder (FSAD).
7617 7618	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complicitions of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system. Endometriosis. Malignant neoplasms of the breast. Benign neoplasms of the breast and other injuries of the breast.
7617 7618 7619 7620 7621 7624 7625 7626 7627 7628 7629 7630 7631	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system. Endometriosis. Malignant neoplasms of the breast. Benign neoplasms of the breast. Benign neoplasms of the breast and other injuries of the breast. Female sexual arousal disorder (FSAD).
7617 7618 7619 7619 7620 7621 7624 7625 7626 7627 7628 7629 7630 7631 7632	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system. Endometriosis. Malignant neoplasms of the breast. Benign neoplasms of the breast and other injuries of the breast. Female sexual arousal disorder (FSAD). The Hematologic and Lymphatic Systems
7617 7618 7619 7620 7621 7624 7625 7626 7627 7628 7629 7630 7631 7632	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system. Endometricsis. Malignant neoplasms of the breast. Benign neoplasms of the breast. Benign neoplasms of the breast and other injuries of the breast. Female sexual arousal disorder (FSAD). The Hematologic and Lymphatic Systems [Removed] Agranulocytosis, acquired. Leukemia.
7617 7618 7619 7620 7621 7624 7625 7626 7627 7628 7629 7630 7631 7700 7700 7702 7703 7704	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system. Endometriosis. Malignant neoplasms of the breast. Benign neoplasms of the breast. Benign neoplasms of the breast and other injuries of the breast. Female sexual arousal disorder (FSAD). The Hematologic and Lymphatic Systems [Removed] Agranulocytosis, acquired. Leukemia. Polycythemia vera.
7617 7618 7619 7620 7621 7624 7625 7626 7627 7628 7630 7631 7763 77700 77702 77704 77704	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system. Endometriosis. Malignant neoplasms of the breast. Benign neoplasms of the breast. Benign neoplasms of the breast. Benign neoplasms of the breast and other injuries of the breast. Female sexual arousal disorder (FSAD). The Hematologic and Lymphatic Systems [Removed] Agranulocytosis, acquired. Leukemia. Polycythemia vera. Immune thrombocytopenia.
7617 7618 7619 7620 7621 7624 7625 7626 7627 7628 7629 7630 7631 7632 7700 7700 7702 7702 7703 7704 7704 7706	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system. Endometricsis. Malignant neoplasms of the breast. Benign neoplasms of the breast and other injuries of the breast. Female sexual arousal disorder (FSAD). The Hematologic and Lymphatic Systems [Removed] Agranulocytosis, acquired. Leukemia. Polycythemia vera. Immune thrombocytopenia. Splenectomy.
7617 7618 7619 7620 7621 7624 7625 7626 7627 7628 7629 7630 7631 7700 7702 7703 7704 7705 7706 7707	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of the breast. Benign neoplasms of the breast. Benign neoplasms of the breast. Benign neoplasms of the breast and other injuries of the breast. Female sexual arousal disorder (FSAD). The Hematologic and Lymphatic Systems [Removed] Agranulocytosis, acquired. Leukemia. Polycythemia vera. Immune thrombocytopenia. Splenectomy. Spleen, injury of, healed.
7617 7618 7619 7620 7621 7624 7625 7626 7627 7628 7630 7631 7632 7700 7700 7702 7704 7705 7706 7707	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system. Endometriosis. Malignant neoplasms of the breast. Benign neoplasms of the breast. Benign neoplasms of the breast. Benign neoplasms of the breast. Female sexual arousal disorder (FSAD). The Hematologic and Lymphatic Systems [Removed] Agranulocytosis, acquired. Leukemia. Polycythemia vera. Immune thrombocytopenia. Splenectomy. Spleen, injury of, healed. Hodgkin's lymphoma.
7617 7618	Uterus and both ovaries, removal. Uterus, removal. Ovary, removal. Ovary, removal. Ovaries, atrophy of both. Complete or incomplete pelvic organ prolapse due to injury or disease or surgical compliations of pregnancy. Fistula, rectovaginal. Fistula, urethrovaginal. Fistula, urethrovaginal. Breast, surgery. Malignant neoplasms of gynecological system. Benign neoplasms of gynecological system. Endometriosis. Malignant neoplasms of the breast. Benign neoplasms of the breast and other injuries of the breast. Female sexual arousal disorder (FSAD). The Hematologic and Lymphatic Systems [Removed] Agranulocytosis, acquired. Leukemia. Polycythemia vera. Immune thrombocytopenia. Splenectomy. Spleen, injury of, healed.

Diagnostic Code No.	
7714	. Sickle cell anemia.
7715	. Non-Hodgkin's lymphoma.
7716	
7717	
7718	
7719	- ,g (, (, (
	kemia).
7720	
7721	1
7722 7723	
7724	
7725	
	THE SKIN
	I TE SKIN
7800	
	or other disfigurement of the head, face, or neck.
7801	
7000	with underlying soft tissue damage.
7802	
7804	ated with underlying soft tissue damage.
7805	
7000	or 7804.
7806	
7807	
7808	
7809	
7811	
7813	. Dermatophytosis.
7815	. Bullous disorders.
7816	
7817	
7818	
7819	
7820	
7821	
7822	
7823 7824	
7825	
7826	
7827	
7828	
7829	
7830	
7831	3 1
7832	
7833	
	THE ENDOCRINE SYSTEM
7900	. Hyperthyroidism, including, but not limited to, Graves' disease.
7901	
7902	
7903	. Hypothyroidism.
7904	
7905	
7906	
7907	
7908 7909	
7911	
7912 7913	
7914	
7915	
7916	
7917	
7918	
7919	
	The same New Comments and Comme

	Diagnostic Code No.	
NEUROLOGICAL CONDITIONS AND CONVULSIVE DISORDERS Organic Diseases of the Central Nervous System		
3000		Encephalitis, epidemic, chronic.
		Brain, New Growth of
		Malignant.
		Benign.
		Paralysis agitans. Bulbar palsy.
		Brain, vessels, embolism.
8008		Brain, vessels, thrombosis.
		Brain, vessels, hemorrhage.
		Myelitis.
		Poliomyelitis, anterior. Hematomyelia.
		Syphilis, cerebrospinal.
		Syphilis, meningovascular.
		Tabes dorsalis.
		Amyotrophic lateral sclerosis. Multiple sclerosis.
		Meningitis, cerebrospinal, epidemic.
3020		Brain, abscess.
		Spinal Cord, New Growths
		Malignant.
		Benign.
		Progressive muscular atrophy. Syringomyelia.
		Myasthenia gravis.
3045		Residuals of traumatic brain injury (TBI).
3046		Cerebral arteriosclerosis.
		Miscellaneous Diseases
3100		Migraine
		Tic, convulsive.
		Paramyoclonus multiplex. Chorea, Sydenham's.
		Chorea, Huntington's.
		Athetosis, acquired.
3108		Narcolepsy.
		The Cranial Nerves
		Fifth (trigeminal), paralysis.
		Seventh (facial), paralysis. Ninth (glossopharyngeal), paralysis.
		Tenth (glossopharyngear), paralysis.
3211		Eleventh (spinal accessory, external branch), paralysis.
		Twelfth (hypoglossal), paralysis.
		Neuritis, fifth cranial nerve. Neuritis, seventh cranial nerve.
		Neuritis, ninth cranial nerve.
3310		Neuritis, tenth cranial nerve.
		Neuritis, eleventh cranial nerve.
		Neuritis, twelfth cranial nerve. Neuralgia, fifth cranial nerve.
		Neuralgia, iitir cranial nerve. Neuralgia, seventh cranial nerve.
		Neuralgia, ninth cranial nerve.
3410		Neuralgia, tenth cranial nerve.
		Neuralgia, eleventh cranial nerve. Neuralgia, twelfth cranial nerve.
		Troutagia, tronai orania norro.
		Peripheral Nerves
8412		Peripheral Nerves Upper radicular group, paralysis.
3412 3510 3511		Upper radicular group, paralysis. Middle radicular group, paralysis.
3510 3511 3512		Upper radicular group, paralysis. Middle radicular group, paralysis. Lower radicular group, paralysis.
3510 3511 3512 3513		Upper radicular group, paralysis. Middle radicular group, paralysis. Lower radicular group, paralysis. All radicular groups, paralysis.
3510 3511 3512 3513 3514		Upper radicular group, paralysis. Middle radicular group, paralysis. Lower radicular group, paralysis.

Diagnostic Code No.	
517	Musculocutaneous nerve, paralysis.
518	
519	
520	
521	
522 523	
524	
525	
526	
527	
528	
529 530	
540	
610	
611	
612	
613	
614	
615 616	
617	
618	
619	Neuritis, long thoracic nerve.
620	
621	
622	
8623 8624	
625	
626	
627	Neuritis, internal saphenous nerve.
628	
629	1,
630	
710 711	
712	
713	
714	
3715	
716	
717	
718 719	
720	
721	
722	
3723	Neuralgia, anterior tibial nerve (deep peroneal).
1724	
725	
726 727	
728	
729	
730	
	The Epilepsies
010	· ·
910 911	Grand mal. Petit mal.
912	
913	
914	
	Mental Disorders
0201	
208	
9210	
211	Schizoaffective Disorder.
9300 9301	

Diagnostic Code No.	
9304	Major or mild neurocognitive disorder due to traumatic brain injury.
9305	Major or mild vascular neurocognitive disorder.
9310	Unspecified neurocognitive disorder.
9312	Major or mild neurocognitive disorder due to Alzheimer's disease.
9326	Major or mild neurocognitive disorder due to another medical condition or substance/medical tion-induced major or mild neurocognitive disorder.
9400	
9403	Specific phobia; social anxiety disorder (social phobia).
9404	
9410	
9411	
9412	Panic disorder and/or agoraphobia.
9413	
9416	
9417	
9421	
9422	
9423	
9424	
9425	
9431	
9432	
9433	Production and the second seco
9434	
9435	
9440	
9520	
9521	
3321	
	DENTAL AND ORAL CONDITIONS
9900	,
9901	
9902	
9903	1
9904	
9905	
9908	
9909	
9911	
9913	
9914	
9915	
9916	
9917	
9918	Neoplasm, hard and soft tissue, malignant.

 $[72\ FR\ 12990,\ Mar.\ 20,\ 2007,\ as\ amended\ at\ 73\ FR\ 54708,\ 54711,\ Sept.\ 23,\ 2008;\ 74\ FR\ 18467,\ Apr.\ 23,\ 2009;\ 77\ FR\ 6467,\ Feb.\ 8,\ 2012;\ 79\ FR\ 45102,\ Aug.\ 4,\ 2014;\ 82\ FR\ 36085,\ Aug.\ 3,\ 2017;\ 82\ FR\ 50807,\ Nov.\ 2,\ 2017;\ 83\ FR\ 15073,\ Apr.\ 9,\ 2018;\ 83\ FR\ 15323,\ Apr.\ 10,\ 2018;\ 83\ FR\ 32600,\ July\ 13,\ 2018;\ 83\ FR\ 54258,\ Oct.\ 29,\ 2018;\ 84\ FR\ 28234,\ June\ 18,\ 2019;\ 85\ FR\ 76466,\ Nov.\ 30,\ 2020;\ 86\ FR\ 8143,\ Feb.\ 4,\ 2021]$

APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES

	Diagnostic code No.
Abscess:	
Brain	8020
Kidney	7501
Lung	6824
Acne	7828
Acromegaly	7908
Actinomycosis	6822
Addison's disease	7911
Agranulocytosis, acquired	7702
AL amyloidosis	7717
Alopecia areata	7831
Amebiasis	7321
Amputation:	
· Arm:	
Complete amputation, upper extremity	5120

		Diagnos code No
Al	pove insertion of deltoid	Ę
Be	elow insertion of deltoid	ŧ
Digits, five	of one hand	
	of one hand:	
TI	numb, index, long and ring	
Th	numb, index, long and little	
	numb, index, ring and little	ŧ
Th	numb, long, ring and little	
In	dex, long, ring and little	
Digits, thre	e of one hand:.	
TI	numb, index and long	
	numb, index and ring	ŧ
	numb, index and little	
Th	numb, long and ring	
	numb, long and little	
Th	numb, ring and little	
In	dex, long and ring	
	dex, long and little	į
	dex, ring and little	
	ong, ring and little	į
	of one hand:	`
	numb and index	
	numb and long	į
	numb and ring	į
	numb and little	į
	dex and long	į
	dex and ring	į
	dex and little	į
	ong and ring	į
		į
	ong and little	ì
	ing and little	•
Single fing		,
	numb	į
	dex finger	-
	ong finger	
	ing finger	
	ttle finger	
Forearm:		
	pove insertion of pronator teres	
	elow insertion of pronator teres	
Leg:		_
	ith defective stump	Ę
	ot improvable by prosthesis controlled by natural knee action	Ę
	lower level, permitting prosthesis	
Fo	prefoot, proximal to metatarsal bones	
	pes, all, amputation of, without metatarsal loss or transmetatarsal, amputation of, with up to	
10		Ę
	half of metatarsal loss	Ę
	half of metatarsal loss	,
To		į
To To	pe, great	
To To	pe, greatpe, other than great, with removal metatarsal head	Ę
To To To Thigh:	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement	Ę
To To To Thigh: Co	pe, greatpe, other than great, with removal metatarsal head	į
To To Thigh: Cl U _l	pe, great pe, other than great, with removal metatarsal head pes, three or more, without metatarsal involvement pemplete amputation, lower extremity	; ;
To To Thigh: Co U M	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Des, three or more, with three or more, without metatarsal involvement Des, three or more, without	; ;
To To To Thigh: Co Uj M yotrophic lateral	pe, great pe, other than great, with removal metatarsal head pes, three or more, without metatarsal involvement pomplete amputation, lower extremity	; ;
To To Thigh: Ci Ul yotrophic lateral atomical loss of:	pe, great pe, other than great, with removal metatarsal head pes, three or more, without metatarsal involvement per third per third iddle or lower thirds sclerosis	
To To Thigh: Ch Ul Wyotrophic lateral atomical loss of: Both eyes	pe, great pe, other than great, with removal metatarsal head pes, three or more, without metatarsal involvement per third iddle or lower thirds sclerosis	; ;
To To Thigh: Cr U M Nyotrophic lateral atomical loss of: Both eyes e eye, with visua	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Des, three or more, without metatarsal head Des, three or more, without metatarsal involvement Des, three or more, with	
To To Thigh: C U _I Myotrophic lateral atomical loss of: Both eyes e eye, with visua	pe, great pe, other than great, with removal metatarsal head pes, three or more, without metatarsal involvement per third iddle or lower thirds sclerosis I acuity of other eye: 200 (1.5/60)	
To T	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Despit third Diddle or lower thirds Sclerosis I acuity of other eye: 200 (1.5/60) J/200 (3/60); 15/200 (4.5/60); 20/200 (6/60)	
To Tromble To Tromble Thigh: Cromble Thigh: Cromble Thigh: Notice Thigh Thi	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Des, three or more, without metatarsal involvement Dest, three or more destination of the second did not be second or more destination of the second did not be second or more destination of the second did not be second or more destination of the second did not be second din the second did not be second did not be second did not be secon	
To Tr Thigh: C. Uly yotrophic lateral atomical loss of: Both eyes e eye, with visua 5/ 10 20	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Des, three or more, without metatarsal involvement Dest three or more involvement Dest three or more involvement Dest three or more involvement Dest three or more, without metatarsal head Dest three or more, without metatarsal involvement Dest three or more, without metatarsal head Dest three or more or more, without metatarsal head Dest three or more or more, without metatarsal head Dest three or more or m	
To Trigh: Thigh: Cright Conditions of: Both eyes are eye, with visual formula in the conditions of: Both eyes are eye, with visual formula in the conditions of: Both eyes are eye, with visual formula in the conditions of th	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Dest three or more, without metatarsal involvement Dest third Dest third Diddle or lower thirds Dest third Diddle or lower	
To T	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Des, three or more, without metatarsal involvem	
To Trigh: Thigh: Co Ul Myotrophic lateral atomical loss of: Both eyes e eye, with visua 5/ 10/ 20/ Both feet . Both feet . Both hand done hand	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Despit three or more, without metatarsal involvement Despit three or more, without metatarsal head Despit three or more or m	
To Trigh: Thigh: Co Up Myotrophic lateral atomical loss of: Both eyes e eye, with visua 65/10/20 Both feet Both feet Both hands One hand	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Dest three or more, without metatarsal head Dest three or more, without metatarsal involvement Dest three or more, without metatarsal involveme	
Thigh: Thigh: Critical Control	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Despit three or more, without metatarsal involvement Despit three or more, without metatarsal head Despit three or more or m	
To Trigh: Thigh: Ci Ul Mayotrophic lateral atomical loss of: Both eyes 6 e eye, with visua 5/ 10 20 Both feet 1. Both head 0. One hand 0. One hand emia:	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Des, three or more oremore or more or	
Thigh: Thigh: C. C. Ui, Mayotrophic lateral atomical loss of: Both eyes e eye, with visua 5/10 20 20 20 20 20 20 20 20 20 20 20 20 20	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Dest three or more, without metatarsal involvem	
Thigh: Thigh: C. C. Ui, Mayotrophic lateral atomical loss of: Both eyes e eye, with visua 5/10 20 20 20 20 20 20 20 20 20 20 20 20 20	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Des, three or more oremore or more or	
To Trigh: Thigh: Ci Up Myotrophic lateral atomical loss of: Both eyes 6 e eye, with visua 5/ 10 20 Both feat 6. Both head 6. One hand One foot a One hand emia: Acquired h Folic acid of Iron deficie	De, great De, other than great, with removal metatarsal head Des, three or more, without metatarsal involvement Dest three or more, without metatarsal involvem	

	Diagnostic code No.
Aortic	711
Large artery	711
Small artery	711
Angioneurotic edema	711
nkylosis:	507
Ankle	527
igits, individual: Thumb	522
Index finger	522
Long finger	522
Ring or little finger	522
Elbow	520
Hand	
avorable:	
Five digits of one hand	522
Four digits of one hand	522 522
Two digits of one hand	522
Unfavorable:	322
Five digits of one hand	521
Four digits of one hand	52
Three digits of one hand	52
Two digits of one hand	52°
lip	52
Knee	52
Scapulohumeral articulation	520
Subastragalar or tarsal joint	52
Wrist	52
Ankylosing spondylitisAphakia	524 602
Aphonia, organic	65
Aplastic anemia	77
Arrhythmia:	• • •
Supraventricular	701
Ventricular	701
Arteriosclerosis obliterans	711
Arteriosclerotic heart disease	700
Arteriovenous fistula	71
Arthritis:	500
Degenerative, other than post-traumatic	500
Other specified forms (excluding gout)	500
Pneumococcic	50
Post-traumatic	50
Multi-joint (except post-traumatic and gout)	50
Streptococcic	50
Syphilitic	50
Typhoid	50
Arthropathy	50
Asbestosis	68
Aspergillosis	68: 66:
Astragalectomy	52
Atherosclerotic renal disease	75
Athetosis	81
Atrioventricular block	70
Avitaminosis	63
Bartonellosis	63
	63
Bladder:	75
Bladder: Calculus in	
Bladder: Calculus in Fistula in	75
Sladder: Calculus in Fistula in Injury of	75 75
Bladder: Calculus in Fistula in Injury of Neurogenic	75 75 75
Bladder: Calculus in Fistula in Injury of Neurogenic Blastomycosis	75 75 75
Bladder: Calculus in Fistula in Injury of Neurogenic Blastomycosis Blastomycosis Blastomycosis	75 75 75 68
Bladder: Calculus in Fistula in Injury of Neurogenic Blastomycosis	75 75 75 68
Bladder: Calculus in Fistula in Injury of Neurogenic Blastomycosis Blindness: see also Vision and Anatomical Loss Both eyes, only light perception On eye, only light perception and other eye: 5/200 (1.5/60)	75 75- 75- 683 600
Fistula in Injury of Neurogenic Slatent See also Vision and Anatomical Loss Blastomycosis Blindness: see also Vision and Anatomical Loss Both eyes, only light perception One eye, only light perception and other eye: 5/2/00 (1.5/60) 10/200 (3/60); 15/200 (4.5/60); 20/200 (6/60)	75: 75: 754 683 606 606
Bladder: Calculus in Fistula in Injury of Neurogenic Blastomycosis Blindness: see also Vision and Anatomical Loss Both eyes, only light perception One eye, only light perception and other eye: 5/200 (1.5/60)	75 75 75 68 60 60

		Diagnost code No
	Neoplasm, benign	5
	Neoplasm, malignant, primary or secondary	5
	Shortening of the lower extremity	5
Brain:		
	Abscess	8
	irgery	7
	ctasis	6
	S	6
	is	6
	s disease	7
	isorders	7
	15010615	5
	pacter jejuni infection	6
	Pacemakers, implantable	7
	Transplantation	7
ardiomy	ropathy	7
-cell hy	perplasia, thyroid	7
ataract:		
	Senile and others	6
	Traumatic	6
	arteriosclerosis	8
	strain	5
	sease or injury	7
norea:	I I realization in	_
	Huntington's	3
	Sydenham's	-
	e	7
	titis, chronic	7
	asis, chronic	7
	dsis, citionic	6
	Fatigue Syndrome (CFS)	6
	ung abscess	6
	obstructive pulmonary disease	ě
	domycosis	6
	ry residuals	7
	cerative	7
ompartr	ment syndrome	5
	e or incomplete pelvic organ prolapse due to injury or disease or surgical complications of pregnancy, ng uterine or vaginal vault prolapse, cystocele, urethrocele, rectocele, enterocele, or combination vitis:	7
•	Trachomatous	6
	Other	6
oronary	bypass surgery	7
oxiella l	purnetii infection (Q Fever)	6
	ccosis	6
	s syndrome	7
	is manifestations of collagen-vascular diseases not listed elsewhere	7
	shuania	-
	chronic	7
	stitisession illness	
	ESSIOIT IIITIESS	-
ecompr	c or oozoma	,
ecompr ermatiti	s or eczema	-
ecompr ermatiti ermator esquam	phytosis	
ecomprermatitiermator ermator esquamator abetes:	phytosis	6
ecomprermatitiermator esquamabetes:	ohytosis ative interstitial pneumonitis	7
ecomprermatitiermatopesquam abetes:	ohytosis	·
ecomprermatitiermatopesquam abetes:	ohytosis	6 7
ecomprermatitiermatopesquamabetes:	ohytosis	6 7
ecompr ermatiti ermator esquam abetes:	ohytosis Lative interstitial pneumonitis Insipidus Mellitus Paralysis or paresis Rupture	6 7 7
ecomprermatitiem at the compression of the compress	ohytosis lative interstitial pneumonitis Insipidus Mellitus m: Paralysis or paresis	
ecomprermatitiermatopesquamabetes: aphrag plopia plopia, sease:	ohytosis Lative interstitial pneumonitis Linsipidus Mellitus m: Paralysis or paresis Rupture	6 5 6
ecompr ermatiti ermator esquam iabetes: iaphrag iplopia iplopia, isease:	ohytosis lative interstitial pneumonitis Insipidus Mellitus m: Paralysis or paresis Rupture Ilimited muscle function, eye Addison's Buerger's	6 7 7 8 8 8
ecompriermatitii ermatopi esquamiabetes: iaphrag iplopia iplopia, isease:	ohytosis lative interstitial pneumonitis Insipidus Mellitus m: Paralysis or paresis Rupture Ilimited muscle function, eye Addison's	7 6 7 7 6 5 6 6
ecompriermatitii ermatopesquamiabetes: iaphrag iplopia iplopia, isease:	ohytosis lative interstitial pneumonitis Insipidus Mellitus m: Paralysis or paresis Rupture Ilimited muscle function, eye Addison's Buerger's	6 7 7 6 5 6 6
ecompr ermatiti ermatop esquam iabetes: iaphrag iplopia iplopia, isease:	ohytosis lative interstitial pneumonitis Insipidus Mellitus m: Paralysis or paresis Rupture Ilimited muscle function, eye Addison's Buerger's Chronic obstructive pulmonary disease	6 7 7 6 5 6 6

	Diagnosti code No
Parasitic	6
Disfigurement of, head, face or neck	78
Dislocated:	,,
Cartilage, semilunar	5
	6
Lens, crystalline	7:
isseminated intravascular coagulation	
istomiasis, intestinal or hepatic	7:
iverticulitis	7:
ysentery, bacillary	7:
ctropion	6
mbolism, brain	8
mphysema, pulmonary	6
ncephalitis, epidemic, chronic	8
ndocarditis	7
ndometriosis	7
nteritis, chronic	7
nterocolitis, chronic	7:
ntropion	6
osinophilic granuloma of lung	6
pididymo-orchitis	7:
pilepsies:	
Diencephalic	8
Grand mal	8
Jacksonian and focal motor or sensory	8
Petit mal	8
Psychomotor	8
	6
piphora	
rythema multiforme	7
rythroderma	7
rythromelalgia	7
sophagus:	
Diverticulum	7
Spasm	7
Stricture	7
allopian tube	7
emale sexual arousal disorder (FSAD)	7
critate sexual disorder (1 6/15)	
ever:	6
ever: Relapsing	
ever: Relapsing Rheumatic	6
ever: Relapsing	6 6
ever: Relapsing	6 6
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia	6 6 5
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano	6 6 5
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano	6 6 5 7
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal	6 6 5 7
ever: Relapsing Rheumatic brosis of lung, diffuse interstitial bromyalgia stula in ano sistula: Rectovaginal Urethrovaginal	6 5 7 7 7
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal	6 5 7 7 7 5
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial bromyalgia istula in ano stula: Rectovaginal Urethrovaginal atfoot, acquired astritis, hypertrophic	6 6 5 7 7 7 5 7
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired asstritis, hypertrophic ienu recurvatum	6 6 5 7 7 7 5 7
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia sistula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired asstritis, hypertrophic eneu recurvatum	6 6 5 7 7 7 5 7
ever: Relapsing Rheumatic brosis of lung, diffuse interstitial bromyalgia sistula in ano situla: Rectovaginal Urethrovaginal atfoot, acquired astritis, hypertrophic enu recurvatum laucoma: Congestive or inflammatory	6 6 5 7 7 7 5 7 5
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia sistula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired asstritis, hypertrophic eneu recurvatum	6 6 5 7 7 7 5 5
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired astritis, hypertrophic enu recurvatum laucoma: Congestive or inflammatory Simple, primary, noncongestive	66 55 7 7 7 5 7 5 6 6
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia sistula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired asstritis, hypertrophic eneu recurvatum laucomae: Congestive or inflammatory Simple, primary, noncongestive	6 6 5 7 7 7 5 7 5 6 6
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired iastritis, hypertrophic ienu recurvatum ilaucoma: Congestive or inflammatory Simple, primary, noncongestive ilout	66 55 7 7 7 55 7 5 6 6 6 6 7
Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibrowyalgia istula: Rectovaginal Urethrovaginal latfoot, acquired Gastritis, hypertrophic sienu recurvatum Silaucoma: Congestive or inflammatory Simple, primary, noncongestive Graves' disease	6 6 5 7 7 7 5 5 6 6 6 7 5
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired iastritis, hypertrophic ienur recurvatum islaucoma: Congestive or inflammatory Simple, primary, noncongestive islomerulonephritis iout irraves' disease lallux:	66 57 77 55 75 66 67 75
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired iastritis, hypertrophic ienu recurvatum ilaucoma: Congestive or inflammatory Simple, primary, noncongestive iout iraves' disease lallux: Rigidus	666557775566667755575557555575555555555
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula: Rectovaginal Urethrovaginal latfoot, acquired iastritis, hypertrophic enu recurvatum llaucoma: Congestive or inflammatory Simple, primary, noncongestive lout iraves' disease allux: Rigidus Valgus	66 55 77 77 75 55 66 77 55 55
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia sistula in ano sistula: Rectovaginal Urethrovaginal latfoot, acquired asstritis, hypertrophic enur recurvatum slaucoma: Congestive or inflammatory Simple, primary, noncongestive slout raves' disease allux: Rigidus Valgus ammer toe	66655777556666775555555555
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired asstritis, hypertrophic ienu recurvatum ilaucoma: Congestive or inflammatory Simple, primary, noncongestive ilomerulonephritis iout raves' disease allux: Rigidus Valgus armer toe eart valve replacement	66655777556666775555555555
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired iastritis, hypertrophic enu recurvatum ilaucoma: Congestive or inflammatory Simple, primary, noncongestive ilomerulonephritis iout rraves' disease allux: Rigidus Valgus ammer toe eart valve replacement ematologic:	66655777755666775577557755775577
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired asstritis, hypertrophic ienu recurvatum ilaucoma: Congestive or inflammatory Simple, primary, noncongestive ilomerulonephritis iout raves' disease allux: Rigidus Valgus armer toe eart valve replacement	66655777755666775577557755775577
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired iastritis, hypertrophic enu recurvatum ilaucoma: Congestive or inflammatory Simple, primary, noncongestive ilomerulonephritis iout rraves' disease allux: Rigidus Valgus ammer toe eart valve replacement ematologic:	666557775566677755777
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal alatfoot, acquired asstritis, hypertrophic een urecurvatum ilaucoma: Congestive or inflammatory Simple, primary, noncongestive ilomerulonephritis iout raves' disease allux: Rigidus Valgus ammer toe eart valve replacement ematologic: Essential thrombocytopenia	66655777556667557755777577777777777777
ever: Relapsing Rheumatic brosis of lung, diffuse interstitial bromyalgia sistula in ano stula: Rectovaginal Urethrovaginal atfoot, acquired astritis, hypertrophic enu recurvatum laucoma: Congestive or inflammatory Simple, primary, noncongestive lomerulonephritis out raves' disease allux: Rigidus Valgus ammer toe eart valve replacement ematologic: Essential thrombocythemia and primary myelofibrosis Immune thrombocytopenia Multiple myeloma	66655777755566677755777777777777777777
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia sistula in ano sistula: Rectovaginal Urethrovaginal latfoot, acquired aastritis, hypertrophic eren recurvatum slaucoma: Congestive or inflammatory Simple, primary, noncongestive slomerulonephritis out raves' disease allux: Rigidus Valgus ammer toe eart valve replacement ematologic: Essential thrombocythemia and primary myelofibrosis Immune thrombocytopenia Multiple myeloma Myelodysplastic syndromes	66655777555666777557777777777777777777
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired iastritis, hypertrophic ienu recurvatum ilaucoma: Congestive or inflammatory Simple, primary, noncongestive ilomerulonephritis iout irraves' disease lallux: Rigidus Valgus lammer toe leart valve replacement lematologic: Essential thrombocythemia and primary myelofibrosis Immune thrombocytopenia Multiple myeloma Myelodysplastic syndromes Solitary plasmacytoma	66655777556667777577777777777777777777
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired iastritis, hypertrophic ienu recurvatum islaucoma: Congestive or inflammatory Simple, primary, noncongestive islomerulonephritis iout irraves' disease lallux: Rigidus Valgus lammer toe leart valve replacement lematologic: Essential thrombocythemia and primary myelofibrosis Immune thrombocythemia Multiple myeloma Myelodysplastic syndromes Solitary plasmacytoma lematomyelia Memorylasmacytoma	66655777556666775577777777777777777777
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired iastritis, hypertrophic ienur recurvatum islaucoma: Congestive or inflammatory Simple, primary, noncongestive islomerulonephritis iout rraves' disease lallux: Rigidus Valgus lammer toe leart valve replacement lematologic: Essential thrombocythemia and primary myelofibrosis Immune thrombocytopenia Multiple myeloma Myelodysplastic syndromes Solitary plasmacytoma lematomyelia lematomyelia lemorrhage:	666557775556667775577777788
Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibrowyalgia istula in ano istula: Rectovaginal Urethrovaginal Definition Congestive or inflammatory Simple, primary, noncongestive Glomerulonephritis Cout Cout	66 55 77 55 77 55 57 77 77 77 77 77 78 8
Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibrowyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired iastritis, hypertrophic ieanu recurvatum Slaucoma: Congestive or inflammatory Simple, primary, noncongestive Slomerulonephritis iout Rigidus Valgus Idammer toe Ideard valve replacement Idematologic: Essential thrombocythemia and primary myelofibrosis Immune thrombocytopenia Multiple myeloma Myelodysplastic syndromes Solitary plasmacytoma Idematomyelia Idemorrhage: Idemorrhage: Idemorrhage: Idemorrhage: Idemorr	6 6 6 5 7 7 7 7 5 5 5 5 5 7 7 7 7 7 8 8
Relapsing Rheumatic ibrorsis of lung, diffuse interstitial ibromyalgia istula: Rectovaginal Urethrovaginal latfoot, acquired asstritis, hypertrophic senu recurvatum slaucoma: Congestive or inflammatory Simple, primary, noncongestive slout araves' disease tallux: Rigidus Valgus dammer toe leart valve replacement tematologic: Essential thrombocythemia and primary myelofibrosis Immune thrombocytopenia Multiple myeloma Myelodysplastic syndromes Solitary plasmacytoma tematorhage: Brain Intra-ocular	6 6 6 6 5 5 7 7 7 7 7 7 7 7 8 8 8 8 8
Relapsing	6 6 6 5 7 7 7 7 5 5 6 6 7 7 7 7 7 7 7 7
ever: Relapsing Rheumatic ibrosis of lung, diffuse interstitial ibromyalgia istula in ano istula: Rectovaginal Urethrovaginal latfoot, acquired iastritis, hypertrophic ienu recurvatum ilaucoma: Congestive or inflammatory Simple, primary, noncongestive ilomerulonephritis iout riaves' disease lallux: Rigidus Valgus lammer toe leart valve replacement lematologic: Essential thrombocythemia and primary myelofibrosis Immune thrombocytopenia Multiple myeloma Myelodysplastic syndromes Solitary plasmacytoma lematomyalia lemorrhage: Brain Intra-ocular	666557777555666777557777777777888

	Diagnost code No
Femoral	7
Hiatal	7
Inguinal	7
Muscle	5
Ventral	7
terotopic ossification	5
Flail joint	
toplasmosis	
-Related Illness	
lgkin's disease	
gkin's lymphoma	
Ironephrosis	
peraldosteronism	
perhidrosis	
perinfection syndrome or disseminated strongyloidiasis	
perparathyroidism	
perpituitarism	
persensitivity	
pertensive:	
Heart disease	
Vascular disease	
perthyroid heart disease	
perthyroidism	
poparathyroidism	
pothyroidism	
pairment of:	
Humerus	
Clavicle or scapula	
Elbow	
Thigh	
Femur	
Knee, other	
Field vision	
Tibia and fibula	
Rectum & anus	
Ulna	
plantable cardiac pacemakers	
ections of the skin	
ury:	
Bladder	
Breast	
Eye, unhealed	
Foot	
Gall bladder	
Lips	
Liver, residuals	
Mouth	
iscle:	
Facial	
Group I Function: Upward rotation of scapula	
Group II Function: Depression of arm	
Group III Function: Depression of arm	
Group IV Function: Stabilization of shoulder	
Group V Function: Elbow supination	
Group VII Function: Extension of elbow	
Group VIII Function: Flexion of wrist and fingers	
Group VIII Function: Extension of wrist, fingers, thumb	
Group IX Function: Forearm muscles	
Group X Function: Movement of forefoot and toes	
Group XI Function: Propulsion of foot	
Group XII Function: Dorsiflexion	
Group XIII Function: Extension of hip and flexion of knee	
Group XIV Function: Extension of knee	
Group XV Function: Adduction of hip	
Group XVI Function: Flexion of hip	
Group XVII Function: Extension of hip	
Group XVII Function: Extension of hip	
Group XVIII Function: Outward rotation of thigh	
Group XVIII Function: Outward rotation of thigh	
Group XVIII Function: Outward rotation of thigh	!
Group XVIII Function: Outward rotation of thigh	!

	Diagnostic code No.
Pharynx	65
Sacroiliac	52
Spinal cord	68
Stomach, residuals of	73
ritis	60
nterstitial nephritis	75
ntervertebral disc syndrome	52
ntestine, fistula of	73
rritable colon syndrome	73
eratinization, diseases of	78
eratitis	60
eratoconus	60
idney:	
Abscess	7:
Cystic diseases	7
Removal	7
Transplant	7
Tuberculosis	7
/phoscoliosis, pectus excavatum / carinatum	6
gophthalmos	6 6
ryngectomy	б
ryngitis:	_
Tuberculous	6
Chronic	6
ırynx, stenosis of	6
ishmaniasis:	
American (New World)	7
Old World	7
prosy (Hansen's Disease)	6
ukemia:	
Chronic myelogenous leukemia (CML) (chronic myeloid leukemia or chronic granulocytic leukemia)	7
Leukemia	7
nitation of extension:	•
Forearm	5
Leq	5
_ ~	5
Radius	
Supination and pronation	5
Thigh	5
mitation of extension and flexion:	_
Forearm	5
mitation of flexion:	
Forearm	5
Leg	5
Thigh	5
mitation of motion:	
Ankle	5
Arm	5
Index or long finger	5
Ring or little finger	5
Temporomandibular	9
Thumb	5
Wrist, limitation of motion	5
/er:	
Disease, chronic, without cirrhosis	7
·	
Transplant	7
Cirrhosis	7
oss of:	_
Auricle	6
Condyloid process	9
Coronoid process	9
Eyebrows	ϵ
Eyelashes	6
Eyelids	6
Palate, hard	9
andible:	
Including ramus, unilaterally or bilaterally	g
axilla:	
More than half	^
MOLE MAIL DAIL	9
	9
Less than half	6
Less than halfose, part of, or scars	
Less than half	5
Less than halfose, part of, or scars	

	Diagnosti code No
peth, loss of	9:
ongue, loss of whole or part	7:
oss of use of:	
Both feet	5
Both hands	5
Foot	5
Hand	5
One hand and one foot	5
mbosacral strain	5
pus:	
Erythematosus	ε
Erythematosus, discoid	7
ne disease	6
nphatic filariasis, to include elephantiasis	6
ılaria	6
lignant melanoma	7
dunion:	_
Mandible	9
Os calcis or astragalus	5
xilla, malunion or nonunion	9
xilla or mandible, chronic osteomyelitis, osteonecrosis, or osteoradionecrosis of	9
lioidosis	6
niere's syndrome	6
ningitis, cerebrospinal, epidemic	8
ntal disorders:	
Anorexia nervosa	9
Bipolar disorder	9
Bulimia nervosa	9
Chronic adjustment disorder	9
Conversion disorder (functional neurological symptom disorder).	Ş
Cyclothymic disorder	9
Delirium	9
Delusional disorder	9
Depersonalization/derealization disorder	S
Dissociative amnesia; dissociative identity disorder	9
Generalized anxiety disorder	S
Illness anxiety disorder	9
Major depressive disorder	9
Major or mild neurocognitive disorder due to Alzheimer's disease	9
Major or mild neurocognitive disorder due to another medical condition or substance/medication-in-	
duced major or mild neurocognitive disorder	9
Major or mild neurocognitive disorder due to HIV or other infections	9
Major or mild neurocognitive disorder due to traumatic brain injury	9
Major or mild vascular neurocognitive disorder	9
Obsessive compulsive disorder	9
Other specified and unspecified schizophrenia spectrum and other psychotic disorders	9
Other specified anxiety disorder	9
Other specified somatic symptom and related disorder	9
Panic disorder and/or agoraphobia	Ş
Persistent depressive disorder (dysthymia)	9
Posttraumatic stress disorder	9
Schizoaffective disorder	9
Schizophrenia	9
Somatic symptom disorder	(
Specific phobia; social anxiety disorder (social phobia)	9
Unspecified somatic symptom and related disorder	9
Unspecified anxiety disorder	(
Unspecified depressive disorder	(
Unspecified neurocognitive disorder	(
tatarsalgia	
graine	8
rton's disease	į
cormycosis	6
ıltiple sclerosis	8
asthenia gravis	8
relitis	8
rocardial infarction	7
rositis	5
rcolepsy	8
eoplasms:	
opiasms: Benign:	

		Diagnost code No
	Ear	6
	Endocrine	7
	Genitourinary	7
	Gynecological	7 9
	Hard and soft tissue	5
	Respiratory	6
	Skin	7
Maligna		
	Breast	7
	Digestive system	7
	Ear	6
	Endocrine	7
	Genitourinary	7
	Hard and soft tissue	g
	Muscle	5
	Respiratory	6
	Skin	7
phritis, chronic		7
		7
	arteriolar	7
uralgia:		
Cranial		_
	Fifth (trigeminal)	8
	Seventh (facial)	8
	Ninth (glossopharyngeal)	8
	Tenth (pneumogastric, vagus)	8
	Twelfth (hypoglossal)	8
Periphe	al Nerves	_
	Upper radicular group	8
	Middle radicular group	8
	Lower radicular group	8
	All radicular groups	8
	Musculospiral (radial)	8
	Median	8
	Ulnar	8
	Musculocutaneous	8
	Circumflex Long thoracic	3
	Sciatic	8
	External popliteal (common peroneal)	8
	Musculocutaneous (superficial peroneal)	8
	Anterior tibial (deep peroneal)	
	Internal popliteal (tibial)	8
	Posterior tibial	8
	Anterior crural (femoral)	8
	Internal saphenous	8
	Obturator	8
	External cutaneous nerve of thigh	8
	Ilio-inguinal	8
uritis:		
Cranial		,
	Fifth (trigeminal)	8
	Seventh (facial)	8
	Tenth (pneumogastric, vagus)	8
	Eleventh (spinal accessory, external branch)	8
	Twelfth (hypoglossal)	8
Optic	() /	6
	al Nerves	
	Upper radicular group	8
	Middle radicular group	8
	Lower radicular group	8
	All radicular groups	8
	Musculospiral (radial)	8
	Median	8
	Ulnar	8
	Musculocutaneous	8
	Circumflex	8
	Long thoracic	

		Diagnost code No
E	external popliteal (common peroneal)	8
	fusculocutaneous (superficial peroneal)	8
	Interior tibial (deep peroneal)	8
	nternal popliteal (tibial)	8
	osterior tibial	8
	Interior crural (femoral)	8
	nternal saphenous	8
	ixternal cutaneous nerve of thigh	8
	io-inguinal	8
	r	7
w growths:		
Benign		
E	ones	5
	rain	8
	ye, orbit, and adnexa	6
	pinal cord	8
Malignant		
	ones	
	train	{
	ye, orbit, and adnexa	6
	pinal cord	
	phoma	
	/cobacterium infection	
	ella infection	Č
nunion:		
	confirmed by diagnostic imaging studies	
	d ulna	
stagmus, centra		(
teitis deformans		
teomalacia, resi	duals of	
teoporosis, resid	luals of	
tis media:		
	rative	(
	/e	(
	L. al.	(
	both	
ary:	r injury	
	i lijury	-
	isorders	
ralysis:		
	dation	
	traumatic	
ralysis, nerve:		
Cranial ne		
	ifth (trigeminal)	
	eventh (facial)	
	linth (glossopharyngeal)	
	enth (pneumogastric, vagus)	
	cleventh (spinal accessory, external branch)	
ا Periphera	welfth (hypoglossal)	
	Ipper radicular group	
	fiddle radicular group	
	ower radicular group	
	Il radicular groups	
	Musculospiral (radial)	
	Median	
	llnar	
	Musculocutaneous	
	Sircumflex	
	ong thoracic	
	ciatic	8
	xternal popliteal (common peroneal)	1

		Diagnost code No
	Anterior tibial nerve (deep peroneal)	8
	Internal popliteal (tibial)	8
	Posterior tibial nerve	8
	Anterior crural nerve (femoral)	8
	Internal saphenous	8
	Obturator	8
	External cutaneous nerve of thighllio-inguinal	8
aramyoclo	onus multiplex	8
	sease	6
		6
nis		
De	eformity, with loss of erectile power	7
	emoval of glans	7
Re	emoval of half or more	7
ricardial	adhesions	7
		7
	vestibular disorders	ϵ
	, adhesions	7
		7
	(Claw foot) acquired	5
	ocytoma	7
	99-	6
	Dilitis	5
	sion or fibrosis	6
	lar syndrome	7
	niosis	6
	s & fibrosis:	
	ug-induced	6
	adiation-induced	8
	s, anterioria vera	7
	lar syndrome	7
	nal disorders	6
	ctomy syndromes	7
	tic syndrome	7
	al residual	6
	e muscular atrophy	8
	and	7
osthetic ii		
An	ıkle replacement	5
Ell	bow replacement	5
Hip	p, resurfacing or replacement	5
Kn	nee, resurfacing or replacement	5
Sh	noulder replacement	5
W	rist replacement	5
		7
		6
		6
Imonary:		,
	veolar proteinosis	6
	iscular disease	7
	itis, chronic	· · · · · · · · · · · · · · · · · · ·
	syndrome	
ctum:	sylutolite	•
	ectum & anus, stricture	7
	olapse	-
moval:	,	
	artilage, semilunar	5
0-	DCCYX	5
	all bladder	7
	dney	7
	enis glans	7
	enis half or more	7
	bs	5
Rit		7
Ril Te	estis	,
Ril Te Ov	vary	7
Rit Te Ov Ute	varyerus	7
Rit Te Ov Ute Ute	vary	, 7 7
Rit Te Ov Uto Uto enal:	varyerus	7 7

	Diagnosti code No
Involvement in systemic diseases	7:
Tubular disorders	7
esection of intestine:	
Large	7:
Small	7:
etina detachment of	6
etinal dystrophy (including retinitis pigmentosa, wet or dry macular degeneration, early-onset macular degen-	6
eration, rod and/or cone dystrophy)etinopathy, diabetic	6
etinopathy, diabetic	6
habdomyolysis, residuals of	5
hinitis:	0.
Allergic or vasomotor	6
Bacterial	6
Granulomatous	6
ickettsial, ehrlichia, and anaplasma Infections	6
arcoidosis	6
carring alopecia	78
cars:	
Burn scar(s) of the head, face, or neck; scar(s) of the head, face, or neck due to other causes; or	
other disfigurement of the head, face, or neck	78
Burn scar(s) or scar(s) due to other causes, not of the head, face, or neck that are associated with un-	_
derlying soft tissue damage	78
Burn scar(s) or scar(s) due to other causes, not of the head, face, or neck that are not associated with	-
underlying soft tissue damage	7
Retina	6
Scars, other; and other effects of scars evaluated under diagnostic codes 7800, 7801, 7802, or 7804 Unstable or painful	7 7
chistosomiasis	6
nigella infections	6
inusitis:	· ·
Ethmoid	6
Frontal	6
Maxillary	6
Pansinusitis	6
Sphenoid	6
leep Apnea Syndrome	6
oft tissue sarcoma:	
Muscle, fat, or fibrous connected	5
Neurogenic origin	8
Vascular origin	7
pine:	
Degenerative arthritis, degenerative disc disease other than intervertebral disc syndrome	5
pinal fusion	5
pinal stenosis	5
pleen, injury of, healed	7
plenectomy	7
condylolisthesis or segmental instability, spine	5
tomach, stenosis of	7
ymblepharon	6
/ndromes:	0
Chronic Fatigue Syndrome (CFS)	6
Cushing's	7
Meniere's	6 7
Raynaud's	6
/philis	6
	U
unhilis:	8
	8
Cerebrospinal	7
Cerebrospinal	
Cerebrospinal	
Cerebrospinal Meningovascular yphilitic heart disease yringomyelia yringomyelia	8
Cerebrospinal Meningovascular // / //philitic heart disease // / //ringomyelia // / // sbes dorsalis // // /	8 8
Cerebrospinal Meningovascular yphilitic heart disease yringomyelia abes dorsalis arsal or metatarsal bones enosynovitis, tendinitis, tendiniosis or tendinopathy	8 8 5
Cerebrospinal Meningovascular yphilitic heart disease yringomyelia abes dorsalis arsal or metatarsal bones anosynovitis, tendinitis, tendinosis or tendinopathy sestis:	8 8 5 5
Cerebrospinal Meningovascular yphilitic heart disease yringomyelia abes dorsalis arsal or metatarsal bones enosynovitis, tendinitis, tendinosis or tendinopathy estis: Atrophy, complete	8 8 5 5
Meningovascular yphilitic heart disease yphilitic heart disease yringomyelia abes dorsalis arsal or metatarsal bones enosynovitis, tendinitis, tendinosis or tendinopathy estis: Atrophy, complete Removal	8 8 5 5 7 7
Cerebrospinal Meningovascular	8 8 5 5 7 7 7
Cerebrospinal Meningovascular Meningovascular Syphilitic heart disease Syringomyelia Abes dorsalis A	8 8 5 5 7 7
Cerebrospinal Meningovascular	8 8 5 5 7 7 7

	Diagnostic code No.
Thyroiditis	79
ic, convulsive	8
innitus, recurrent	62
oxic nephropathy	75
raumatic brain injury residuals	80
raumatic chest wall defect	6
uberculosis:	
Adenitis	7
Bones and joints	50
Eye	6
Kidney	7:
Luposa (lupus vulgaris)	7
Miliary Pleurisy, active or inactive	6
Pulmonary:	0
Active, far advanced	6
Active, moderately advanced	6
Active, minimal	6
Active, advancement unspecified	6
Active, chronic	6
Inactive, chronic	6
Inactive, far advanced	6
Inactive, moderately advanced	6
Inactive, minimal	6
Inactive, advancement unspecified	6
Fuberculosis luposa (lupus vulgaris)	7
Tympanic membrane	6
Jicer:	
Duodenal	7
Gastric	7
Marginal	7:
Jreter, stricture of	7.
Ureterolithiasis	7
Urethra.	_
Fistula	7
Stricture	7
Urticaria, chronic.	7
Uterus: And both ovaries, removal	70
Disease or injury	70
Prolapse	7
Removal	7
Jveitis	6
Vagina, disease or injury	7
Vagotomy	7
Valvular heart disease	7
Varicose veins	7
/asculitis, primary cutaneous	7
Vertebral fracture or dislocation	5
Vibriosis (Cholera, Non-cholera)	6
Visceral Leishmaniasis	6
Visceroptosis	7
Vision: see also Blindness and Loss of	
One eye 5/200 (1.5/60), with visual acuity of other eye:	
5/200 (1.5/60)	6
10/200 (3/60); 15/200 (4.5/60); 20/200 (6/60)	6
20/100 (6/30); 20/70 (6/21); 20/50 (6/15)	6
20/40 (6/12)	6
One eye 10/200 (3/60), with visual acuity of other eye:	_
10/200 (3/60); 15/200 (4.5/60); 20/200 (6/60)	6
20/100 (6/30); 20/70 (6/21); 20/50 (6/15)	6
20/40 (6/12)	6
15/200 (4.5/60) or 20/200 (6/60)	6
20/100 (6/30); 20/70 (6/21); 20/50 (6/15)	6
20/40 (6/12)	6
20/40 (6/12)	"
20/200 (6/60)	_
	6
20/100 (6/30); 20/70 (6/21); 20/50 (6/15)	6
One eve 20/100 (6/30), with visual acuity of other eve: and other eve:	
20/100 (6/30); 20/70 (6/21); 20/50 (6/15)	6

Pt. 5

	Diagnostic code No.
One eye 20/70 (6/21), with visual acuity of other eye:	
20/70 (6/21) or 20/50 (6/15)	6078
20/40 (6/12)	6079
One eye 20/50 (6/15), with visual acuity of other eye:	
20/50 (6/15)	6078
20/40 (6/12)	6079
Each eye 20/40 (6/12)	6079
Vitiligo	7823
Vulva or clitoris, disease or injury of	7610
Weak foot	5277
West Nile virus infection	6335

[72 FR 13003, Mar. 20, 2007, as amended at 73 FR 54708, 54712, Sept. 23, 2008; 73 FR 69554, Nov. 19, 2008; 74 FR 18467, Apr. 23, 2009; 77 FR 6467, Feb. 8, 2012; 79 FR 45103, Aug. 4, 2014; 82 FR 36085, Aug. 3, 2017; 82 FR 50807, Nov. 2, 2017; 83 FR 15073, Apr. 9, 2018; 83 FR 15323, Apr. 10, 2018; 83 FR 32601, July 13, 2018; 83 FR 5259, Oct. 29, 2018; 84 FR 28234, June 18, 2019; 85 FR 76467, Nov. 30, 2020; 85 FR 85523, Dec. 29, 2020; 86 FR 8143, Feb. 4, 2021]

PART 5—ADMINISTRATIVE PROCE-DURES:GUIDANCE DOCUMENTS

Sec.

- 5.0 Purpose.
- 5.10 Definitions relating to guidance documents.
- 5.15 Procedures for issuing guidance documents.
- 5.20 Procedures for petition for the withdrawal or modification of a guidance document
- 5.25 Guidance website.

AUTHORITY: 38 U.S.C. 501; E.O. 13891, 84 FR 55235.

EFFECTIVE DATE NOTE: At 86 FR 30184, June 7, 2021, the authority citation for part 5 was revised, effective July 7, 2021. For the convenience of the user, the revised text is set forth as follows:

AUTHORITY: 38 U.S.C. 501; E.O. 12866, 58 FR 51735, 3 CFR, 1993 Comp., p. 638.

Source: 85 FR 72570, Nov. 13, 2020, unless otherwise noted.

§ 5.0 Purpose.

Sections 5.0 through 5.25 provide VA's processes and procedures for issuing and managing guidance documents in accordance with Executive Order 13891.

EFFECTIVE DATE NOTE: At 86 FR 30184, June 7, 2021, §5.0 was revised, effective July 7, 2021. For the convenience of the user, the revised text is set forth as follows:

§ 5.0 Purpose.

This part provides the Department of Veterans Affairs' (VA's) processes and procedures for issuing and managing guidance documents.

§5.10 Definitions relating to guidance documents.

The following definitions apply to $\S 5.0$ through 5.25.

Guidance document means an agency statement of general applicability (i.e., it applies to more than just one person, event, or transaction), that is intended to have a future effect on the behavior or actions of regulated parties (to include non-VA actors), and that sets forth a policy on a statutory, regulatory, or technical issue, or an interpretation of a statute or regulation. A guidance document does not include the following:

- (1) Rules promulgated pursuant to notice and comment under section 553 of title 5, United States Code, or similar statutory provisions;
- (2) Rules exempt from rulemaking requirements under section 553(a) of title 5, United States Code;
- (3) Rules of agency organization, procedure, or practice;
- (4) Decisions of agency adjudications under section 554 of title 5, United States Code, or similar statutory provisions;
- (5) Internal guidance directed to the issuing agency or other agencies that is not intended to have substantial future effect on the behavior of regulated parties; or
- (6) Internal executive branch legal advice or legal opinions addressed to executive branch officials.

Significant guidance document means a guidance document that the Administrator of the Office of Information and