

§ 526.363

21 CFR Ch. I (4–1–25 Edition)

§ 526.363 Cephapirin benzathine.

(a) *Specifications.* Each single-dose, 10-milliliter syringe contains 300 milligrams cephapirin activity (as cephapirin benzathine).

(b) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.115 of this chapter.

(d) *Conditions of use in dry cows*—(1) *Amount.* Infuse the contents of one syringe (300 mg cephapirin activity) into each quarter following last milking, but no later than 30 days before calving.

(2) *Indications for use.* For the treatment of mastitis caused by susceptible strains of *Streptococcus agalactiae* and *Staphylococcus aureus*, including penicillin-resistant strains.

(3) *Limitations.* For use in dry cows only. Not to be used within 30 days of calving. Milk from treated cows must not be used for food during the first 72 hours after calving. Animals infused with this product must not be slaughtered for food until 42 days after the latest infusion. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 37174, Aug. 22, 1978, as amended at 53 FR 27851, July 25, 1988; 73 FR 12262, Mar. 7, 2008; 75 FR 10168, Mar. 5, 2010; 76 FR 17338, Mar. 29, 2011; 86 FR 13186, Mar. 8, 2021; 88 FR 16549, Mar. 20, 2023]

§ 526.365 Cephapirin sodium.

(a) *Specifications.* Each single-dose, 10-milliliter syringe contains 200 milligrams (mg) cephapirin sodium activity.

(b) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.115 of this chapter.

(d) *Conditions of use in lactating cows*—(1) *Amount.* Infuse the contents of one syringe (200 mg cephapirin activity) into each infected quarter immediately after the quarter has been completely milked out. Do not milk out for 12 hours. Repeat once only in 12 hours.

(2) *Indications for use.* For the treatment of mastitis in lactating cows caused by susceptible strains of *Streptococcus agalactiae* and *Staphylococcus aureus* including strains resistant to penicillin.

(3) *Limitations.* Milk that has been taken from animals during treatment

and for 96 hours after the last treatment must not be used for food. Treated animals must not be slaughtered for food until 4 days after the last treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 57455, Dec. 10, 1975, as amended at 53 FR 27852, July 25, 1988. Redesignated at 63 FR 8349, Feb. 19, 1998; 65 FR 20733, Apr. 18, 2000; 73 FR 3181, Jan. 17, 2008; 75 FR 10168, Mar. 5, 2010; 86 FR 13186, Mar. 8, 2021; 88 FR 16549, Mar. 20, 2023]

§ 526.464 Cloxacillin benzathine.

(a) *Specifications.* Each single-dose, 7.5- or 10-milliliter syringe contains cloxacillin benzathine equivalent to 500 milligrams (mg) cloxacillin.

(b) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.165 of this chapter.

(d) *Conditions of use in dry cows*—(1) *Amount.* Infuse the contents of one syringe (equivalent to 500 mg cloxacillin) into each quarter immediately after last milking, but no later than 30 days before calving.

(2) *Indications for use.* For the treatment of mastitis caused by *Staphylococcus aureus* and *Streptococcus agalactiae* including penicillin resistant strains in dairy cows during the dry period.

(3) *Limitations.* Animals infused with this product must not be slaughtered for food until 30 days after the latest infusion. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[86 FR 13186, Mar. 8, 2021]

§ 526.465 Cloxacillin sodium.

(a) *Specifications.* Each single-dose, 10-milliliter syringe contains cloxacillin sodium equivalent to 200 milligrams (mg) cloxacillin.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.165 of this chapter.

(d) *Conditions of use in lactating cows*—(1) *Amount.* Infuse the contents of one syringe (equivalent to 200 mg cloxacillin) into each infected quarter. Treatment should be repeated at 12-hour intervals for a total of 3 doses.