

DEPARTMENT OF THE INTERIOR

Fish and Wildlife Service

50 CFR Part 17

Endangered and Threatened Wildlife and Plants; 12-Month Finding for a Petition to List the Yosemite Toad

AGENCY: Fish and Wildlife Service, Interior.

ACTION: Notice of 12-month petition finding.

SUMMARY: We, the U.S. Fish and Wildlife Service (Service), announce a 12-month finding for a petition to list the Yosemite toad (*Bufo canorus*) under the Endangered Species Act of 1973, as amended (Act). We find that the petitioned action is warranted, but precluded by higher priority listing actions. We will develop a proposed rule to list this species pursuant to our Listing Priority System (48 FR 43098). Upon publication of this notice of 12-month petition finding, this species will be added to our candidate species list.

DATES: The finding announced in this document was made on November 27, 2002. Comments and information may be submitted until further notice.

ADDRESSES: You may send data, information, comments, or questions concerning this finding to the Field Supervisor, U.S. Fish and Wildlife Service, Sacramento Fish and Wildlife Office, 2800 Cottage Way, Room W-2605, Sacramento, CA 95825. You may inspect the petition, administrative finding, supporting information, and comments received, by appointment, during normal business hours at the above address.

FOR FURTHER INFORMATION CONTACT: Susan Moore at the Sacramento Fish and Wildlife Office (see ADDRESSES above) (telephone 916/414-6600; facsimile 916/414-6712).

SUPPLEMENTARY INFORMATION:**Background**

Section 4(b)(3)(B) of the Endangered Species Act of 1973, as amended (Act) (16 U.S.C. 1531 *et seq.*), requires that, for any petition to revise the List of Threatened and Endangered Species containing substantial scientific and commercial information that listing may be warranted, we conduct a status review and make a finding within 12 months of the date of the receipt of the petition on whether the petitioned action is: (a) Not warranted, (b) warranted, or (c) warranted but precluded from immediate proposal by other higher priority proposals. Section 4(b)(3)(C) of the Act requires that

petitions for which a requested action is found to be warranted but precluded should be treated as though resubmitted on the date of such finding, *i.e.*, requiring a subsequent finding to be made within 12 months. Such 12-month findings are to be published promptly in the **Federal Register**.

Section 4(b) of the Act states that we may make warranted but precluded findings only if we can demonstrate that: (1) An immediate proposed rule is precluded by other pending actions, and (2) expeditious progress is being made on other listing actions. Due to the large amount of litigation over critical habitat, we are working on numerous court orders and settlement agreements. Complying with these orders and settlement agreements will consume all of our listing budget for fiscal year 2003. However, we can continue to place species on the candidate species list, as that work activity is funded separately from our listing program.

Taxonomy

The Yosemite toad was originally described by Camp (1916), and given the common name Yosemite Park toad. Subsequent detections of this species indicated that its range extends beyond the boundaries of Yosemite National Park, and Grinnel and Storer (1924) referred to this species as the Yosemite toad.

Similarities in appearance of the Yosemite toad and the western toad (*Bufo boreas*) were noted by Camp (1916). Based on general appearance, structure and distribution, it appears that these two species are closely related (Myers 1942; Stebbins 1951; Mullally 1956; Savage 1958). The close relationship between *B. boreas* and *B. canorus* is also supported by studies of bone structure (Tihen 1962a,b), and by the survivorship of hybrid toads produced by artificially crossing the two species (Blair 1959, 1963, 1964).

Camp (1916), using characteristics of the skull, concluded that *Bufo boreas*, *B. canorus*, and *B. nestor* (extinct) are more closely related to each other than to other North American toads, and that these species comprise the most primitive group of *Bufo* in North America. Blair (1972) grouped *B. boreas*, *B. canorus*, black toads (*B. exsul*), and Amargosa toads (*B. nelsoni*), together taxonomically as the "boreas group."

Feder (1977) found *Bufo canorus* to be genetically distinctive based on samples from a limited geographic range. However, Yosemite toads are thought to hybridize with western toads in the northern part of their range (Karlstrom 1962; Morton and Sokolski 1978). Shaffer *et al.* (2000) performed genetic

analysis of a segment of mitochondrial DNA from 372 Yosemite toads found in Yosemite and Kings Canyon National Parks. Their data showed significant genetic differences in Yosemite toads between the two National Parks. They also found significant genetic variability within Yosemite National Park between drainages, and within both Parks between breeding sites. Their data also indicated that black toads are a subgroup within Yosemite toads rather than a separate species. Stephens (2001) examined mitochondrial DNA from 8 Yosemite toads (selected from the samples examined by Shaffer *et al.* (2000) to represent the range of variability found in that study) and 173 western toads. Stephens' data indicate that *Bufo* in the Sierra Nevada occur in northern and southern evolutionary groups, each of which include both Yosemite and western toads (*i.e.*, toads of both species are more closely related to each other within a group than they are to members of their own species in the other group). Further genetic analysis of Yosemite toads sampled from throughout their range, and from other toad species surrounding their range is needed to fully understand the evolutionary history and appropriate taxonomic status of the Yosemite toad (Stephens 2001).

Description and Natural History

Yosemite toads are moderately sized, with a snout-urostyle length (measured from the tip of the snout to the posterior edge of the urostyle, a bony structure at the posterior end of the spinal column) of 30 to 71 millimeters (mm) (1.2 to 2.8 inches (in)) with rounded to slightly oval paratoid glands (a pair of glands, one on each side of the head, that produce toxins) (Karlstrom 1962). The paratoid glands are less than the width of a gland apart (Stebbins 1985). A thin mid-dorsal (middle of the back) stripe is present in juveniles of both sexes. The stripe disappears or is reduced with age, and more quickly in males (Jennings and Hayes 1994). The iris of the eye is dark brown with gold iridophores (reflective pigment cells) (Jennings and Hayes 1994). Males are smaller than females, with less conspicuous warts (Stebbins 1951). Differences in coloration between males and females are more pronounced in the Yosemite toad than in any other North American frog or toad (Stebbins 1951). Females have black spots or blotches edged with white or cream that are set against a grey, tan or brown background color (Jennings and Hayes 1994). Males have a nearly uniform dorsal coloration of yellow-green to olive drab to darker greenish brown (Jennings and Hayes

1994). Karlstrom (1962) suggested that differences in coloration between the sexes evolved because they provide the Yosemite toad with protective coloration. The uniform coloration of the adult male matches and blends with the silt and grasses that they frequent during the breeding season, whereas the young and females with disruptive coloration tend to use a wider range of habitats with broken backgrounds; thus coloration may help conceal individual toads from predators.

Yosemite toads overwinter in rodent burrows (Jennings and Hayes 1994). They emerge from hibernation as soon as snowmelt pools form near their overwintering sites (Karlstrom 1962; Kagarise Sherman 1980; Jennings and Hayes 1994). Observed emergence times range from early May to the middle of June (Kagarise Sherman 1980).

Males form breeding choruses, and breeding begins soon after emergence (Jennings and Hayes 1994). Males call during the day and early evening (Stebbins 1951). The breeding call is a mellow long sustained trill with 10 to more than 20 notes (Stebbins 1951). Males have been observed to attack other males to prevent them from calling, to amplex (amplexus is a characteristic clasping of the female by the male during mating) other toads in trial and error search for females, and to attack amplexed pairs and attempt to take over the female (Kagarise Sherman 1980). In studies by Kagarise Sherman (1980), males that mated successfully were more likely to be larger, have arrived at breeding sites earlier, and have stayed at breeding sites longer.

Eggs are typically deposited in shallow water with silty bottoms (Karlstrom 1962). Ideal habitat for egg development is between 2–4 centimeters (cm) (0.8–1.6 in) deep, and eggs do not survive in water deeper than 6 cm (2.4 in) (David Martin, University of California, Santa Barbara, pers. comm. 2002). Eggs are deposited in gelatinous strings (Stebbins 1951; Karlstrom and Livezey 1955) which are intertwined with vegetation and buried in silt (Karlstrom 1962). Eggs are relatively large (2.1 mm (0.08 in) average diameter) and brownish black to jet black over the upper three quarters, and gray to tannish gray over the lower one quarter (Jennings and Hayes 1994). Females are estimated to deposit between 1,000 to 1,500 eggs (Kagarise Sherman 1980).

When not breeding, adults feed in meadow or moist upland habitat until they hibernate (Kagarise Sherman 1980; D. Martin, pers. comm. 2002). Although they are largely diurnal (active during the day) (Jennings and Hayes 1994),

especially while breeding, recent evidence shows that they primarily feed and move at night (D. Martin, pers. comm. 2002).

Eggs generally hatch within 3 to 6 days depending on water temperature (Jennings and Hayes 1994), although they may take over 15 days (Kagarise Sherman 1980). Tadpoles typically transform within 40 to 50 days after fertilization. Tadpoles are not known to overwinter (Jennings and Hayes 1994), although immature tadpoles have been observed well into September (Mullally 1956). Tadpoles tend to congregate (Brattstrom 1962) and use warm shallow water during the day (Cunningham 1963), then retreat to deeper water at night (Mullally 1953). The tadpoles are uniformly black, the snout is blunt, the intestines are scarcely or not at all visible, and the dorsal fin is transparent and marked with few relatively large melanophores (dark-colored pigment cells) (Stebbins 1951). Tadpoles measure 10 to 37 mm (0.39 to 1.45 in) in length (Stebbins 1951, 1985).

Newly metamorphosed juveniles are around 10 mm (0.39 in) in snout-urostyle length (Jennings and Hayes 1994). Some individuals may reproduce at 2 years of age, but growth is slow in both sexes and most individuals require more time to reach maturity (Jennings and Hayes 1994). Males have been observed to first breed at 3 to 5 years and females at 4 to 6 years (Kagarise Sherman 1980; Kagarise Sherman and Morton 1984). Females probably do not breed every year (Morton 1981). Yosemite toads are long lived, with females documented as reaching 15 years old and males 12 years old (Kagarise Sherman and Morton 1984).

Kagarise Sherman (1980) observed one female Yosemite toad move 270 meters (m) (885 feet (ft)) in 65 days and one male move 150 m (492 ft) in 9 days. Toads in her study generally moved 150 to 230 m (492 to 755 ft) each spring from their hibernation sites to their breeding sites. In studies in which toads were repeatedly located using radiotelemetry equipment (D. Martin, pers. comm. 2002), adult toads were observed to moving up to approximately 610 m (2,000 ft) in a single night. During the active season (spring-summer), females generally spend less time at, and travel further away from, breeding ponds than males (Kagarise Sherman 1980). Young of year metamorphs (young toads that have just transformed from tadpoles) probably hibernate closer to the ponds in which they were born than adult toads (Kagarise Sherman 1980). Stebbins (1951) suggested that isolation or semi-isolation of subpopulations of Yosemite toads is likely because they are unlikely

to cross large, dry, forested areas between meadows.

Adult and juvenile Yosemite toads are lie-and-wait predators. They remain motionless until a prey item approaches, then strike and capture the prey with their sticky tongues (Kagarise Sherman and Morton 1984). The examined stomach contents of Yosemite toads have included beetles, ants, centipedes, spiders, dragonfly larvae, mosquitos, and moth and butterfly larvae (Grinnel and Storer 1924; Mullally 1953). They will also prey on flies, bees, wasps, millipedes (Kagarise Sherman and Morton 1984), spider mites, crane flies, springtails, owl flies, and damsel flies (Martin 1991).

Yosemite toad tadpoles graze on detritus and plant material such as algae and will also eat other items such as lodgepole pine pollen. Yosemite toad tadpoles can also be carnivorous and will eat other Yosemite toad tadpoles (see Natural Mortality, below), Pacific chorus frog (previously Pacific treefrog) (*Pseudacris regilla*, previously *Hyla regilla*) tadpoles, diving beetle larvae, and dead mammals (Martin 1991).

Habitat Requirements

Yosemite toads use meadow habitats surrounded by lodgepole pine (*Pinus contorta*) or whitebark pine (*P. albicaula*) (Camp 1916). They are most likely to be found in areas with thick meadow vegetation or patches of low willows (*Salix* spp.) (Mullally 1953). They are most often seen near water, but only occasionally in water (Mullally and Cunningham 1956), and use rodent burrows for overwintering and probably for temporary refuge during the summer (Jennings and Hayes 1994). They also use spaces under surface objects, including logs and rocks, for temporary refuge (Stebbins 1951; Karlstrom 1962). Breeding habitat includes the edges of wet meadows and slow-flowing streams (Jennings and Hayes 1994). Tadpoles have also been observed in shallow ponds and shallow areas of lakes (Mullally 1953). Moist upland areas such as seeps and springheads are important summer non-breeding habitats for adult toads (D. Martin, pers. comm. 2002).

Natural Mortality

Mountain yellow-legged frogs (*Rana muscosa*) (Mullally 1953), aquatic dragonfly larvae (Jennings and Hayes 1994), diving beetles (*Dytiscus* spp.) (Kagarise Sherman and Morton 1984), and possibly larval long-toed salamanders (*Ambystoma macrodactylum*) (Jennings and Hayes 1994) prey on the young life stages of Yosemite toads. American robins

(*Turdus migratorius*) prey on Yosemite toad tadpoles (Jennings and Hayes 1994). Garter snakes (*Thamnophis spp.*) have been observed to eat yearling Yosemite toads (D. Martin, pers. comm. 2002), and are probably the most significant predator on tadpoles and metamorphs (Karlstrom 1962; Jennings and Hayes 1994). California gulls (*Larus californicus*) and Clark's nutcrackers (*Nucifraga columbiana*) have been observed killing adult toads (Mulder *et al.* 1978; Kagarise Sherman 1980; Kagarise Sherman and Morton 1993). Cannibalism has been recorded in Yosemite toad tadpoles (Martin 1991; Chan 2001). The tadpoles have not been observed to kill each other, but they do wound each other in feeding frenzies, and have been observed eating dead tadpoles of their own species (Martin 1991; Chan 2001; D. Martin, pers. comm. 2002).

Dessication of breeding habitat before tadpoles metamorphose is a major cause of mortality (Zeiner *et al.* 1988; Kagarise Sherman and Morton 1993; Jennings and Hayes 1994). Eggs are sometimes killed by freezing (Kagarise Sherman and Morton 1984). Fungal growth has also been observed on eggs (Kagarise Sherman 1980), but it is unclear whether the fungus causes mortality or grows after the eggs die from other causes.

Toads may die of exposure when crossing snow or ice (Kagarise Sherman 1980). Toads that emerge from hibernation early may suffer from exposure and inability to feed if there are late-season storms (Kagarise Sherman 1980).

Adult toads of either sex may drown or asphyxiate when multiple males attempt to amplex a single female. Kagarise Sherman (1980) documented the death of a single female in this manner, and found three additional females and two males that may also have died during multiple amplexus.

Historic and Current Range and Status

The historic range of Yosemite toads in the Sierra Nevada occurs from the Blue Lakes region north of Ebbetts Pass (Alpine County) to 5 kilometers (km) (3.1 miles (mi)) south of Kaiser Pass in the Evolution Lake/Darwin Canyon area (Fresno County) (Jennings and Hayes 1994). The historic elevational range of Yosemite toads is 1,460 to 3,630 m (4,790 to 11,910 ft) (Stebbins 1985).

Pre-1990 historic records of Yosemite toad localities are primarily from museum records and incidental sightings. Systematic habitat surveys looking specifically for Yosemite toad populations have only been conducted since the early 1990s. Therefore, it is

impossible to know how many populations have declined or become extinct, because we do not know how many populations originally existed. Sites first documented after 1990 are useful to illustrate the current range of the species, but are not useful in discussing its decline, due to lack of baseline data. Based on the number of historic sites that are no longer occupied (*see below*), it is possible that many populations have disappeared without ever having been documented.

Since 1990, 292 sites throughout Yosemite toads' historic range have been surveyed, and 229 sites have been confirmed to be occupied. Known Yosemite toad locations by area is based on the most comprehensive dataset on Yosemite toad localities available, which was collected by the U.S. Forest Service (USFS) for use in their conservation assessment of the species (as required by the Sierra Nevada Forest Plan Amendment (U.S. Department of Agriculture (USDA) 2001f)). This dataset was compiled by the USFS and came from various sources, including University of California and California State University researchers, the California Academy of Science, the National Park Service (NPS), the U.S. Geological Survey, the California Department of Fish and Game (CDFG), and the California Natural Diversity Data Base. The following discussion on the number of Yosemite toad sites should be considered an approximation, based on best available information, because surveys are ongoing and some sites may have not yet been reported and added to the database. Also, multiple sightings in close proximity to each other have been considered as a single site for the purposes of this discussion.

The historic and current acreage of Yosemite toad habitat (wet meadows, shallow breeding waters, and moist uplands) within the historic range of Yosemite toads is unknown, although these habitats have been degraded and may be decreasing in area as a result of conifer encroachment and livestock grazing (*see Factor A below*). The vast majority of land within the range of the Yosemite toad is federally managed, with 919,011 hectares (ha) (2,270,918 acre (ac)) (99 percent of the range) on USFS, NPS, and Bureau of Land Management lands. Much of this land is within designated wilderness. The remaining land within the species' range is a mix of State, local government, and private lands.

The following known site discussion is based on the California Wildlife Habitat Relations range map, obtained as a geographic information system data

from CDFG for the species, although this map includes large areas of unsuitable habitat. However, this map is the best available range map for the species, although the species has been detected in a few locations outside its boundaries, primarily at the southern end of the range. The site specific information is based on localized studies that do not represent a comprehensive range-wide assessments of the species status.

(1) Yosemite toads are known from three sites in the southeast corner of the El Dorado National Forest where it borders with the Toiyabe and Stanislaus National Forests. Two of these three sites have been confirmed as occupied since 1990.

(2) Yosemite toads are known from 25 locations along the west side of the Toiyabe National Forest, 15 of which have been confirmed as occupied since 1990.

(3) Yosemite toads are known from 28 sites on the Stanislaus National Forest, 22 of which have been confirmed as occupied since 1990. These sites occur primarily in two groups, one on the northern edge of the forest, where it borders with the El Dorado and Toiyabe National Forests, and the other in a band extending west across the Stanislaus National Forest, from its southeast border with Yosemite National Park and the Toiyabe National Forest.

(4) Yosemite toads are known from 49 sites along the west side of Inyo National Forest, 35 of which have been confirmed as occupied since 1990.

(5) Yosemite toads are known from 91 locations throughout Sierra National Forest, of which 84 have been confirmed as occupied since 1990.

(6) Yosemite toads are known from 78 sites scattered throughout Yosemite National Park, 57 of which have been confirmed occupied since 1990.

(7) Yosemite toads are known from 18 sites throughout the northern half of Kings Canyon National Park, 14 of which have been confirmed as occupied since 1990.

It is impossible to fully determine the extent to which Yosemite toads have declined, because baseline data on the number and size of historic populations are few. The following studies, which reassess the current status of historically documented populations, give the most insight into the species' decline.

Jennings and Hayes (1994) reviewed the current status of Yosemite toads using museum records of historic and recent sightings, published data, and unpublished data and field notes from biologists working with the species. They mapped 55 historically

documented general localities throughout the range of the species where the toad had been present (based on 144 specific sites), and found that Yosemite toads are now absent from 29 of those localities, a decline of over 50 percent.

In 1990, David Martin surveyed 75 sites throughout the range of the Yosemite toad for which there are historic records of the species' presence, and found that 47 percent of those sites showed no evidence of any life stage of the species (Stebbins and Cohen 1997), a decline of about 63 percent.

Grinnell and Storer (1924) surveyed for vertebrates at 40 sites along a 143-km (89-mi) west-to-east transect across the Sierra Nevada, through Yosemite National Park, in 1915 and 1919. Drost and Fellers (1996) conducted more thorough surveys, specifically for amphibians, at 38 of those sites in 1992. They found that Yosemite toads were absent from 6 of 13 sites in which they had been found in the original survey. At sites where Drost and Fellers (1996) found Yosemite toads, the toads occurred in low numbers (only 15 total adult and juvenile toads at all sites), with documented declines in relative abundance in three of the Grinnell and Storer (1924) sites, as based on their generalized abundance categories such as rare, common, and abundant. Therefore, the species has declined or disappeared completely from at least 9 of 13 (69 percent) of the Grinnell and Storer (1924) sites.

The only long-term study on the size of a population of Yosemite toads indicates that the population has declined dramatically. Kagarise Sherman and Morton (1993) studied Yosemite toads at Tioga Pass Meadow (Mono County, California) intensively from 1971 to 1982, and made less systematic observations from 1983 to 1991. To estimate the adult population size, they captured and marked toads entering breeding pools. From 1974 to 1978, an average of 258 males entered the breeding pools. In 1979, the number of male toads began to decline, and by 1982, the number of males had dropped to 28. During the same time period, the number of females varied between 45 and 100, but there was no obvious trend in number observed. In periodic surveys between 1983 and 1991, it appeared that both males and females continued to decline, and breeding activity became sporadic. In 1990, the researchers were only able to locate one female, two males, and four to six egg masses. In 1991, they found only one male and two egg masses. The researchers also surveyed non-breeding habitat in the same area and found similar population

declines. To date, the population at Tioga Pass Meadow has not recovered (Roland Knapp, Sierra Nevada Aquatic Research Laboratory, pers. comm. 2002).

Kagarise Sherman and Morton (1993) also conducted occasional surveys of six other populations in the eastern Sierra Nevada. Five of these populations showed serious, apparently long-term, declines between 1978 and 1981, while the sixth population held relatively steady until the final survey in 1990, at which time it dropped precipitously. In 1991, E.L. Karlstrom revisited the site at which he had studied a breeding population of Yosemite toads from 1954 to 1958, just south of Tioga Pass Meadow within Yosemite National Park (Tuolumne County, California), and found no evidence of toads or signs of breeding (Kagarise Sherman and Morton 1993).

Previous Federal Action

On April 3, 2000, we received a petition to list the Yosemite toad as endangered from the Center for Biological Diversity and Pacific Rivers Council. On October 12, 2000, we announced a 90-day petition finding in the **Federal Register** (65 FR 60607) concluding that the petition presented substantial scientific or commercial information to indicate that the listing of the Yosemite toad may be warranted.

This 12-month finding is made in accordance with a settlement agreement which requires us to complete a finding by November 30, 2002 (*Center for Biological Diversity and Pacific Rivers Council v. Norton and Jones*, No. C-01-2106 (N.D. Calif.)).

Summary of Factors Affecting the Species

Section 4 of the Act and regulations (50 CFR part 424) promulgated to implement the listing provisions of the Act describe the procedures for adding species to the Federal lists. A species may be determined to be an endangered or threatened species due to one or more of the five factors described in section 4(a)(1). In the case of the Yosemite toad, the specific relationship between the potential threats under each factor and the continued decline of the species remains unclear. These factors, and their application to the Yosemite toad, are as follows:

A. The present or threatened destruction, modification, or curtailment of its habitat or range. The following discussion presents several threats to the species' habitat or range.

Grazing

Livestock grazing began in Sierra Nevada meadow and riparian areas with

the settlement of California by the Spanish in the mid-1700s, and rose to a level that caused significant impacts in the mid-1800s following the gold rush (Menke *et al.* 1996). In general, livestock grazing within the range of the Yosemite toad was at a high, but undocumented, level until the establishment of National Parks (beginning in 1890) and National Forests (beginning in 1905) (Menke *et al.* 1996) in the Sierra Nevada area. Within established National Parks, livestock grazing was gradually eliminated, but packstock grazing was permitted and has increased over time (Menke *et al.* 1996).

Over time within established National Forests, the amount of grazing was gradually reduced, better documented, and the type of animals grazed shifted from predominantly sheep to cattle and packstock (Menke *et al.* 1996). In general, livestock grazing within the National Forests in the Sierra Nevada has continued with gradual reductions since the 1920s, except for an increase during World War II (Menke *et al.* 1996). Currently, there are numerous active and inactive livestock grazing allotments on the five National Forests that occur within the range of the Yosemite toad. Approximately 71 active and 36 inactive allotments occur across the Eldorado, Toiyabe, Inyo, Stanislaus, and Sierra National Forests (Laura Conway, Stanislaus National Forest, pers. comm. 2002; Holly Eddinger, Sierra National Forest, *in litt.*, 2002; Aimee Smith, Sierra National Forest, *in litt.*, 2002).

Since 1970, the continuing decrease in grazing permitted on the National Forests has been motivated by concern for resource protection (Menke *et al.* 1996). National Forests have conducted projects to minimize or rehabilitate areas impacted by grazing, including exclosures around some sensitive areas, erosion control structures, and replanting of riparian species.

Packstock grazing is the only grazing currently allowed in National Parks, and it is also allowed in National Forests. There has been very little monitoring of the impacts of packstock use in the Sierra Nevada, which increased after World War II due to increased road access, and increases in leisure time and disposable income (Menke *et al.* 1996). The recreational use of packstock and horsebackriding in the Sierra Nevada can be expected to increase further as human populations increase (State of California 2001; USDA 2001g).

Mule deer (*Odocoileus hemionus*) and bighorn sheep (*Ovis canadensis*) have always occurred within the habitats used by the Yosemite toad (Ingles 1965).

However, grazing by dense groups of large herbivores such as cattle and horses is not a natural situation in those habitats, and these habitats are vulnerable to degradation. Because Yosemite toad breeding habitat is shallow, that habitat is very vulnerable to changes in hydrology caused by grazing (D. Martin, pers. comm. 2002; R. Knapp, pers. comm. 2002).

Direct and indirect mortality of Yosemite toads have occurred as a result of livestock grazing. Cattle have been observed to trample Yosemite toad eggs and disturb eggs such that they fall into hoofprints or other deeper water and die. Metamorph Yosemite toads have been observed to fall into cattle hoofprints or to be defecated on by cattle, become trapped, and die, and adult Yosemite toads have been observed trampled to death in cattle hoofprints (D. Martin, pers. comm. 2002). Preliminary research data indicate that Yosemite toad tadpoles in grazed areas take longer to metamorphose and produce smaller metamorphs than those in areas being rested from grazing, presumably due to high bacteria and nutrient levels, causing low water quality in the grazed areas (D. Martin, pers. comm. 2002).

Grazing removes vegetative cover, and before/after surveys have shown reductions in the number of Yosemite toads using an area after the herbaceous cover was grazed (D. Martin, pers. comm. 2002). Grazing can also cause erosion by disturbing the ground, removing vegetation, and destroying peat layers in meadows, which lowers the groundwater table and summer flows (Armour *et al.* 1994; D. Martin, pers. comm. 2002). Consequently, this may increase the stranding and mortality of tadpoles, or make these areas completely unsuitable for Yosemite toads (D. Martin, pers. comm. 2002). Grazing can also degrade or destroy moist upland areas used as non-breeding habitat by Yosemite toads (D. Martin, pers. comm. 2002), especially when nearby meadow and riparian areas have been fenced to exclude livestock. Livestock may also collapse rodent burrows used by Yosemite toads as cover and hibernation sites, or disturb toads and disrupt their behavior.

The impacts of grazing on habitat can be inferred by observing the recovery of vegetation, ground stability, and water flow that occurs when riparian areas are fenced to exclude livestock (Kattelmann and Embury 1996). An example of this, from a drainage occupied by Yosemite toads, is provided by a study of fish habitat on Silver King and Coyote Valley Creeks (tributaries of the Carson River, Alpine County, California). In

this study, stream reaches were fenced to exclude cattle and, over time, bank stability increased and stream channels became deeper and narrower than the unfenced reaches. This indicated that streambank sloughing had been reduced and vegetation was stabilizing soils and reducing erosion (Overton *et al.* 1994; Kattelmann and Embury 1996).

Livestock grazing in the Sierra Nevada has been so widespread for so long that, in most places, no ungrazed areas are available to illustrate the natural condition of the habitat (Kattelmann and Embury 1996). Due to the long, and historically unregulated history (Menke *et al.* 1996) of livestock and packstock grazing in the Sierra Nevada, and the lack of historic Yosemite toad population size estimates, it is difficult to make a quantitative link between grazing and reductions in Yosemite toad populations. However, because of the documented negative effects of livestock on Yosemite toad habitat, and documented direct mortality of the species caused by livestock, the decline of some populations of Yosemite toad has been attributed to the effects of livestock grazing (Jennings and Hayes 1994; Jennings 1996).

Roads and Timber Harvest

Any activity that severely alters the terrestrial environment, such as road construction and timber harvest, is likely to result in the reduction and occasional extirpation of amphibian populations in the Sierra Nevada (Jennings 1996). By creating gaps in the natural vegetation, roads and harvested areas may act as dispersal barriers and contribute to the fragmentation of Yosemite toad habitat and populations. Habitat fragmentation has been shown to have a negative effect on amphibian species richness (Lehtinen *et al.* 1999). Timber harvest removes vegetation and causes ground disturbance and soil compaction, which makes that ground more susceptible to erosion (Helms and Tappeiner 1996). Much of the erosion caused by timber harvests is from logging roads (Helms and Tappeiner 1996). This erosion could damage Yosemite toad breeding habitat by lowering the water table, and drying out riparian habitats used by the species.

Prior to the formation of National Parks and National Forests, timber harvest was widespread and unregulated, but primarily took place at low elevations on the west slope of the Sierra Nevada, below the elevational range of the Yosemite toad (University of California (UC) 1996). Between 1900 and 1950, the majority of timber harvest took place on old growth forests on private land (UC 1996). The majority of

roads in National Forests of the Sierra Nevada were built between 1950 and 1990 to allow access to the forests for timber harvest (USDA 2001h). Between 1950 and the early 1990s, the USFS allowed major increases in timber harvest on National Forests and at higher elevations, and the majority of impacts on Yosemite toads probably took place during this period.

Roads may cause the potential for direct mortality of amphibians through roadkill (deMaynadier and Hunter 2000), and the possible introduction of contaminants such as petroleum products, herbicides, and pesticides. The levels of timber harvest and road construction have declined substantially since implementation of the California Spotted Owl Sierran Province Interim Guidelines in 1993, and some existing roads have been, or are scheduled for, decommissioning (USDA 2001h). Therefore, the risks posed by new roads and timber harvests have declined, but those already existing still pose risks to the species and its habitat through erosion, roadkill, and contaminant introduction.

Vegetation and Fire Management Activities

Vegetation management includes the removal of small trees and brush to reduce fuels, and to reduce competition which allows faster growth of desired tree species (Helms and Tappeiner 1996). These activities may disturb the ground and increase erosion, which could cause damage to Yosemite toad habitat through siltation and lowering of groundwater levels. Brush removal sometimes includes the use of herbicides, which may run off into Yosemite toad habitat, causing lethal or sublethal effects on individuals (*see* Factor D and E below).

Long-term fire suppression has influenced changes in forest structure and dynamics in the Sierra Nevada. In general, the fire return interval is now much longer than it was historically, and live and dead fuels are more abundant and continuous (USDA 2001c). Fire is thought to be important in maintaining open aquatic and riparian habitats for amphibians in some systems (Russel *et al.* 1999).

Fire suppression, and changes in fire frequency and hydrology, has probably contributed to the decline of Yosemite toads through habitat loss caused by conifer encroachment on meadows (Chang 1996; NPS 2002). Under natural conditions, conifers are excluded from meadows by fire and soils too saturated for their survival. But as conifers begin to encroach on a meadow, if they are not occasionally set back by fire, they

transpire water out of the meadow, reducing the saturation of the soils, and facilitating further conifer encroachment. Therefore, some vegetation treatment may be needed to maintain or restore Yosemite toad habitat.

Increases in fuel abundance have created the potential for catastrophic fires which could cause direct mortality of Yosemite toads; however, data on the direct effects of fire on Yosemite toads are lacking. Fires and mechanical fire suppression activities (such as cutting fire lines) could cause erosion and siltation that could negatively impact Yosemite toad habitat. However, amphibians in general are thought to retreat to moist or subterranean refuges and thereby suffer low mortality during natural fires (Russel *et al.* 1999).

Fire retardant chemicals contain nitrogen compounds or surfactants (soaps). Laboratory tests of these chemicals have shown that after surfactants and ammonia are released when they are added to water, they cause mortality in fish and aquatic invertebrates (Hamilton *et al.* 1996), and likely have similar effects on amphibians. Therefore, if fire retardant chemicals were dropped in or near Yosemite toad habitat, they could have negative effects on individual toads. The majority of vegetation and fire management activities take place at lower elevations, but they do pose a threat to the species when they take place within the species' elevational range.

Recreation

Recreational activities take place throughout the Sierra Nevada and can have significant negative impacts on wildlife and their habitats (USDA 2001a). Recreation is the fastest growing use of National Forests (USDA 2001f). Heavy foot traffic in riparian areas tramples vegetation, compacts soils, and can physically damage streambanks. Trails (foot, horse, bicycle, or off-highway motor vehicle) compact soil in riparian habitat, which increases erosion, replaces vegetation, and can lower the water table (Kondolph *et al.* 1996). Trampling or the collapsing of rodent burrows by recreationists, pets, and vehicles could lead to direct mortality of all life stages of the Yosemite toad. Recreational activity may also disturb toads and disrupt their behavior (Karlstrom 1962).

Dams and Water Diversion

Several artificial lakes are located in or above Yosemite toad habitat, most notably Edison, Florence, Huntington, Courtright, and Wishon Reservoirs. By

altering the timing and magnitude of water flows, these reservoirs have caused changes in hydrology which may have negatively altered Yosemite toad habitat. Changes in water flows have caused increased water levels upstream of the reservoirs, which may have reduced the suitability of shallow water habitats necessary for egg laying, or allowed the invasion of predatory fish into those habitats. Water flow changes may have contributed to the mortality of eggs and tadpoles either by stranding during low water or inundation during high water. The reservoirs themselves probably cover what was once Yosemite toad habitat. Most native Sierra Nevada amphibians cannot live in or move through reservoirs (Jennings 1996). Therefore, reservoirs represent both a loss of habitat and a barrier to dispersal and gene flow. These factors have probably contributed to the decline of Yosemite toads and continue to pose a risk to the species.

B. Overutilization for commercial, recreational, scientific, or educational purposes. There is no known commercial market for Yosemite toads. There is also no documented recreational or educational use for Yosemite toads, although it is likely that they have been handled by curious members of the public and collected as pets.

Scientific research may cause some stress to Yosemite toads through disturbance and disruption of behavior, handling, and injuries associated with marking individuals. Scientific research has resulted in the death of a few individuals through accidental trampling (Green and Kagarise Sherman 2001), irradiation where Karlstrom (1957) collected data on Yosemite toad movements by implanting them with radioactive tags, and collection for museum specimens (Jennings and Hayes 1994). Given the current reduced size and number of populations (Jennings and Hayes 1994), further collection could pose a serious threat to Yosemite toad populations.

C. Disease or predation. Prior to the stocking of high Sierra Nevada lakes with salmonid fishes, which began over a century ago, fish were entirely absent from most of this region (Bradford 1989). Introduced fish, such as rainbow and golden trout (*Oncorhynchus mykiss* ssp.), brown trout (*Salmo trutta*), and brook trout (*Salvelinus fontinalis*), have been shown to have a negative impact, primarily through predation, on native populations of Sierra Nevada amphibians, including the mountain yellow-legged frog (Bradford 1989; Knapp and Matthews 2000) and Pacific chorus frog (Matthews *et al.* 2001).

Data on the effects of introduced fish on Yosemite toads are less clear, although re-surveys of historic Yosemite toad sites have shown that the species had disappeared from several lakes where they formally bred and which are now occupied by fish (Stebbins and Cohen 1997; D. Martin, pers. comm. 2002). Drost and Fellers (1994) state that Yosemite toads are less vulnerable to fish predation than frogs because they breed primarily in ephemeral waters that do not support fish. The palatability of Yosemite toad tadpoles to fish predators is unknown (Jennings and Hayes 1994), but is often assumed to be low based on the unpalatability of western toads (Drost and Fellers 1994; Kiesecker *et al.* 1996), to which Yosemite toads are closely related. Brook trout have been observed to prey on Yosemite toad tadpoles and to "pick at" Yosemite toad eggs, which later became infected with fungus (D. Martin, pers. comm. 2002). Brook trout have been observed to swim near, but ignore, Yosemite toad tadpoles, which gives evidence towards tadpoles being unpalatable, at least in some situations. If Yosemite toad tadpoles are unpalatable to trout, some tadpoles may still be taken by trout that have not learned to avoid them yet (R. Knapp, pers. comm. 2002). The palatability of metamorph Yosemite toads to trout is also unknown, but metamorph western toads have been observed in golden trout stomach contents (R. Knapp, pers. comm. 2002).

At a site where Yosemite toads normally breed in small meadow ponds, they have been observed to successfully switch breeding activities to stream habitat containing fish during years of low water (Phil Strand, Sierra National Forest, pers. comm. 2002). Thus, drought conditions can increase the toads' exposure to predatory fish. Also, although the number of lake breeding sites used by Yosemite toads is small relative to the number of ephemeral sites, lake sites may be especially important because they are more likely to be useable during years with low water (R. Knapp, pers. comm. 2002).

The effects of introduced fish on Yosemite toads needs further study, especially palatability experiments to determine the level of predation. Because Yosemite toads primarily breed in ephemeral waters, fish are probably less of an impact on them than on amphibians that breed primarily in perennial lakes and streams. However, the observed predation of Yosemite toad tadpoles by trout (Martin 1992; D. Martin, pers. comm. 2002) indicate that introduced fish do pose a risk to the species in some situations, which may

be accentuated during drought years. Therefore, introduced fish have probably contributed to the decline of the species. As Yosemite toad populations become smaller and more fragmented, the impacts of predation may be significant.

Various diseases have been confirmed in dead Yosemite toads (Green and Kagarise Sherman 2001). Those diseases, in concert with other factors, are likely to have contributed to the decline of Yosemite toads and continue to be a risk to the species. Mass die-offs of amphibians have been attributed to: chytrid fungal infections of metamorphs and adults (Carey *et al.* 1999); *Saprolegnia* fungal infections of eggs (Blaustein *et al.* 1994); iridovirus infection of larvae, metamorphs, or adults; and bacterial infections (Carey *et al.* 1999). Humans, pets, livestock, packstock, vehicles, and wild animals may all act as disease vectors. Although it has not been observed in the Sierra Nevada, introduced fish may also serve as disease vectors to amphibians. Infection of both fish and amphibians by the same pathogen has been documented with viral (Mao *et al.* 1999) and fungal (Blaustein *et al.* 1994) pathogens.

Tissue samples from dead or dying adults and from healthy tadpoles were collected during a die-off of adult Yosemite toads at Tioga Pass Meadow and Saddlebag Lake and analyzed for disease (Green and Kagarise Sherman 2001). Several infections were found in the adults, including: chytridiomycosis (chytrid fungal infection), bacillary bacterial septicemia (red-leg disease), *Dermosporidium* (a fungal infection), myxozoan infection (parasitic cnidarians (relatives of jellyfish)), *Rhabdias* spp. (a parasitic roundworm) infection, and several species of trematode (parasitic flatworm) infection. However, no single infectious disease was found in more than 25 percent of individuals, and some dead toads showed no infection that would explain their death. No evidence of infection was found in tadpoles. The authors concluded that the die-off was caused by suppression of the immune system caused by an undiagnosed viral infection or chemical contamination that made the toads susceptible to the diagnosed infections. This seems likely considering the evidence suggesting environmental contaminants as a factor contributing to the decline of Yosemite toads (see Factor E).

Carey (1993) developed a model to explain the disappearance of boreal toads (*Bufo boreas boreas*) in the Rocky Mountains. In that model, she hypothesized that the toads were

stressed by some unknown environmental factor. This stress caused a physiological response that suppressed the immune system, which was further hindered by cold temperatures typical of the toads' high-elevation environment. The toads then died of infection by pathogens normally found in their environment. This model may fit Yosemite toad die-offs, given the close relationship between the two toads and their occupation of similar habitats.

Saprolegnia ferax is a species of water mold that commonly infects fish. This mold has been documented to cause massive lethal infection of eggs of western toads in Oregon (Blaustein *et al.* 1994). However, it is unclear whether the infection was caused by the introduction of the fungal pathogen via fish stocking, or if the fungus was already present and the eggs' ability to resist infection was inhibited by some unknown environmental factor. Subsequent laboratory experiments (Kiesecker *et al.* 2001), showed that the fungus could be passed from hatchery fish to western toads. Fungal growth on Yosemite toad eggs was observed by Kagarise Sherman (1980), but the fungal species was not determined, and it was unclear whether the fungus killed the eggs or grew on them after they died of some other cause.

D. The inadequacy of existing regulatory mechanisms. The Yosemite toad occurs on Federal, State, and private lands. Existing regulatory mechanisms do not fully protect this species or its habitat on these lands. Federal, State, and local laws have been insufficient to prevent past and ongoing losses of the limited habitat of the Yosemite toad.

Under section 404 of the Clean Water Act (CWA), the U.S. Army Corps of Engineers (Corps) regulates the discharge of fill material into waters of the United States, including wetlands. However, 99 percent of the Yosemite toad's range is on Federal land, so few projects that include fill of wetlands are likely in these areas. Therefore, section 404 of the CWA is not likely to be relevant to the Yosemite toad in most cases.

Yosemite toads may not be taken or possessed within a National Park without a special permit from the NPS. In addition, cattle grazing, stocking of invasive fish, and most timber harvest are prohibited within National Park boundaries (NPS 2001). However, Yosemite toads have continued to decline within the National Parks in which the species occurs. This may be, in part, due to the Parks allowing such activities as packstock grazing and

recreation in Yosemite toad habitat, as well as chemical contamination of the species and its habitat from sources outside the Parks.

The Wilderness Act of 1964 calls for designated wilderness land "to be protected and managed so as to preserve its natural conditions." Timber harvest and the use of motor vehicles are generally prohibited within wilderness areas, but cattle grazing and invasive fish stocking are permitted within National Forest wilderness lands and pose a threat to the species and its habitat. The species has declined sharply (Jennings and Hayes 1994) regardless of wilderness designation in large portions of its range.

The Yosemite toad is considered a sensitive species by the USFS. Each National Forest was required to complete a Land and Resource Management Plan (LRMP) by the Forest and Rangeland Renewable Resources Planning Act of 1974, as amended by the National Forest Management Act of 1976 (NFMA). Those acts require that the LRMPs provide for multiple use and sustained yield of the products and services obtained from the National Forests, including wildlife. The Sierra Nevada Forest Plan Amendment (Amendment) (USDA 2001d) amends the LRMPs of National Forests within the Sierra Nevada to address issues pertaining to: old forest ecosystems and associated species; aquatic, riparian, and meadow ecosystems and associated species; fire and fuels; noxious weeds; and lower westside hardwood ecosystems. The Amendment calls for the preparation of a conservation assessment, activity-related standards and guidelines, and conservation measures by the USFS to protect Yosemite toads and their habitat occurring in National Forests within the Sierra Nevada.

Under the Amendment to the LRMPs of National Forests within the Sierra Nevada, (USDA 2001f), the USFS is to provide the following conservation measures for Yosemite toads under: (A) Exclude livestock (including pack and saddle stock) from standing water and saturated soils in wet meadows and associated streams and springs occupied by Yosemite toads, or identified as "essential habitat" in the conservation assessment for the Yosemite toad during the breeding and rearing season (as determined locally). If physical exclusion of livestock, such as fencing, is impractical, then exclude grazing from the entire meadow until the meadow has been dry for 2 weeks. Wet meadows are defined as relatively open meadows with low to moderate amounts of woody vegetation that have standing

water and saturated soils after the first of June; if these conditions do not persist in the meadow for more than 2 weeks, allow grazing only in those portions of the meadow where dry conditions exist; (B) Monitor a sample of occupied Yosemite toad sites to assess: (1) Habitat conditions, and (2) Yosemite toad occupancy and population dynamics. Based on the monitoring data, modify or suspend grazing if Yosemite toad conservation is not being accomplished. These grazing restrictions may be modified through formal adaptive management studies, developed in cooperation with the USFS's Pacific Southwest Research Station, designed to assess the effects of grazing intensity and frequency on Yosemite toad habitat conditions and site occupancy; and (C) Conduct surveys of unoccupied suitable habitat for the Yosemite toad within this species' historic range to determine presence of Yosemite toads. Complete surveys of these areas within 3 years of January 2001. If surveys are not completed within the 3-year period, consider unsurveyed meadows as occupied habitat and apply restrictions for excluding livestock described in (A).

Conservation measures also include direction to avoid application of pesticides within 152 m (500 ft) of known Yosemite toad sites, and the removal of invasive fish from some areas of mountain yellow-legged frog habitat, which could benefit Yosemite toads if they are also using those areas (USDA 2001d). The conservation measures also set limits for grazing utilization of grasses and shrubs, livestock use and road construction in willow flycatcher (*Empidonax trailii*) habitat (which includes areas that may also be inhabited by Yosemite toads), packstock use of Yosemite toad habitat during the breeding and rearing season, and disturbance of streambanks and lakeshores. The conservation measures also recommend removing livestock gathering and handling facilities from riparian and meadow areas and providing off-stream watering devices for livestock. The Amendment also includes requirements for monitoring to review how well the objectives established by the Amendment have been met, and how closely management standards and guidelines have been applied (USDA 2001e).

The USFS has been implementing these conservation measures since 2001, but they have not yet been fully implemented. The Amendment is currently being reviewed, and it remains unknown if these measures will be changed, or if any additional protection of the Yosemite toad will be included.

Therefore, the Amendment has not yet provided sufficient protection for the Yosemite toad and its habitat, and it is not known if it will in the future. Also, the effect of the LRMPs in place on National Forests within the Sierra Nevada is unknown. Yosemite toads have continued to decline (Jennings and Hayes 1994).

The State of California considers the Yosemite toad a species of special concern, but it is not State listed as a threatened or endangered species under the California Endangered Species Act. California Sport Fishing Regulations include the Yosemite toad as a protected species that may not be taken or possessed at any time except under special permit from the CDFG. This gives the Yosemite toad some legal protection from collecting, but does not protect it from other causes of mortality or alterations to its habitat.

The California Environmental Quality Act (CEQA) requires review of any project that is undertaken, funded, or permitted by a State or local governmental agency. If a project with potential impacts on Yosemite toad were reviewed, CDFG personnel could determine that, although not listed, the toad is a *de facto* endangered, threatened, or rare species under section 15380 of CEQA. Once significant effects are identified, the lead agency has the option of requiring mitigation for effects through changes in the project or to decide that overriding considerations make mitigation infeasible (CEQA Sec. 21002). In the latter case, projects may be approved that cause significant environmental damage, such as destruction of listed endangered species or their habitat. Protection of listed species through CEQA is, therefore, dependent upon the discretion of the agency involved.

The California Forest Practice rules set guidelines for the design of timber harvests on private land to reduce impacts on non-listed species. However, these rules have little application to the protection of Yosemite toad because approximately 99 percent of the species' range is on Federal land.

The California Department of Pesticide Regulation has authority to restrict the use of pesticides. Their Toxic Air Contaminant (TAC) Program includes assessment of the risks posed by airborne pesticides by collecting air samples near sites of pesticide application and in communities near those sites. If air samples indicate that reductions in exposure are needed, mitigation measures are developed to bring about those reductions (California Department of Pesticide Regulation 2001). However, the TAC program is

intended primarily to protect human health, and air samples are not taken at far distant locations from application sites, like those inhabited by Yosemite toads.

E. Other natural or manmade factors affecting its continued existence.

Yosemite toads probably are exposed to a variety of pesticides and other chemicals throughout their range. Environmental contaminants could negatively affect the species by causing direct mortality; suppressing the immune system; disrupting breeding behavior, fertilization, growth or development of young; and disrupting the ability to avoid predation (Carey and Bryant 1995). Hydrocarbon and other contamination from oil production and road runoff; the application of numerous chemicals for agricultural production; roadside maintenance; and rodent and vector control programs may all have negative effects on Yosemite toad populations. Also, the airborne transport of pesticides as a result of drift from agricultural applications, including chlorothalonil, malathion, diazinon, and chlorpyrifos, from the Central Valley of California to the Sierra Nevada, has been documented (Aston and Seiber 1997; McConnell *et al.* 1998) in samples of air, rain, snow, lake water, and pine needles.

Cholinesterase is an enzyme that functions in the nervous system and is disrupted by organophosphorus pesticides, including malathion, chlorpyrifos, and diazinon (Sparling *et al.* 2001). Reduced cholinesterase activity and pesticide residues have been found in Pacific chorus frog larvae collected in the Sierra Nevada downwind of the Central Valley (Sparling *et al.* 2001). Cholinesterase activity was significantly lower in samples from the Sierra Nevada than from samples taken from coastal California, upwind of the Central Valley. No samples were taken above approximately 1,500 m (4,900 ft) elevation (Sparling *et al.* 2001), which barely overlaps the 1,460 to 3,630 m (4,790 to 11,910 ft) elevational range (Stebbins 1985) of Yosemite toads. However, significant amounts of pesticide residues have been documented as high as 1,920 m (6,300 ft) in Sequoia National Park, south of Yosemite and Kings Canyon National Parks (Aston and Seiber 1997; McConnell *et al.* 1998). In addition to interfering with nerve function, contaminants may act as estrogen mimics (Jennings 1996), or may otherwise disrupt endocrine function (Carey and Bryant 1995), and may have a negative effect on amphibian populations.

Dichlorodiphenyltrichloroethane (DDT) and its residues were found in frogs throughout the Sierra Nevada during the late 1960s (Corey *et al.* 1970), and those residues still appear in Pacific chorus frog larvae collected in the late 1990s (Sparling *et al.* 2001), over 25 years after DDT was banned for use in the United States.

Spatial analysis of populations of Yosemite toads shows a trend towards greater decline in populations downwind of areas of the Central Valley with more agriculture, where there is presumably more pesticide use; however this trend is not statistically significant (Carlos Davidson, California State University, Sacramento, *in litt.*, 2002).

Snow core samples from the Sierra Nevada contain a variety of contaminants from industrial and automotive sources including: hydrogen ions (indicative of acidic precipitation), nitrogen and sulfur compounds (NH₄, NO₃, SO₂, and SO₄), and heavy metals (Pb, Fe, Mn, Cu, and Cd) (Laird *et al.* 1986). The pattern of recent frog extinctions in the southern Sierra Nevada corresponds with the pattern of highest concentration of air pollutants from automotive exhaust, possibly due to increases in nitrification (or other changes), caused by those pollutants (Jennings 1996).

The effects of contaminants on amphibians needs further research (Hall and Henry 1992), and there are few, if any, studies on the direct effect of contaminants on Yosemite toads. However, we know of one study which shows that there are significant levels of contaminants that have been deposited in the Sierra Nevada, and the correlative evidence between areas of contamination in the Sierra Nevadas and areas of amphibian decline (Jennings 1996; Sparling *et al.* 2001; C. Davidson, *in litt.*, 2002), and the significant evidence of an adverse physiologic effect of pesticides on Sierra Nevada amphibians in the field (Sparling *et al.* 2001), indicate that contaminants may be a severe risk to the Yosemite toad and may have contributed to the species' decline.

Rodent control programs probably have an adverse indirect effect on Yosemite toad populations. Control of rodents that create burrows, such as ground squirrels, could significantly reduce the number of burrows available for use by Yosemite toads that require them for hibernation. Because the burrow density required to support Yosemite toads in an area is not known, the loss of burrows as a result of control programs cannot be quantified at this time. Active rodent colonies probably

are needed to sustain Yosemite toads because inactive burrow systems become progressively unsuitable over time. Loredó *et al.* (1996) found that burrow systems collapsed within 18 months following abandonment by, or loss of, the ground squirrels. Rodent control programs must be analyzed and implemented carefully in Yosemite toad habitat so the persistence of the species is not threatened. Much of the species' range is occupied by livestock, primarily cattle, and most livestock owners seek to eliminate rodent burrows because of the threat of cows breaking their legs if they accidentally step into a burrow.

The last century has included some of the most variable climate reversals, at both the annual (extremes and high frequency of El Niño and La Niña events) and near decadal scales (periods of 5- to 8-year drought and wet periods) that has been documented (USDA 2001b). These events may have negative effects on Yosemite toads. Severe winters (El Niño) would force longer hibernation times, and could stress the toads by reducing the time available for them to feed and breed. Severe winters may also depress reproductive effort. Morton (1981) theorized that fluctuations in energy storage from year to year may explain why many female Yosemite toads do not breed on a yearly basis. Alternately, during mild winters (La Niña), precipitation is reduced. This reduction in precipitation could lead to stranding and death of Yosemite toad eggs and tadpoles, a major documented source of mortality (Zeiner *et al.* 1988; Kagarise Sherman and Morton 1993; Jennings and Hayes 1994), or to increased exposure to predatory fish.

Changes in climate that occur faster than the ability of endangered species to adapt could cause local extinctions (U.S. Environmental Protection Agency (EPA) 1989). Analysis of the Antarctic Vostok ice core has shown that over the past 160,000 years, temperatures have varied with the concentrations of greenhouse gasses such as carbon dioxide and methane (Harte 1996). Since the pre-industrial era, atmospheric concentrations of carbon dioxide have increased nearly 30 percent, methane concentrations have more than doubled, and nitrous oxide (another greenhouse gas) levels have risen approximately 15 percent (EPA 1997). The burning of fossil fuels is the primary source of these increases (EPA 1997). Global mean surface temperatures have increased 0.3 to 0.7 Celsius (0.6–1.2 Fahrenheit) since the late 19th century (EPA 1997). Climate modeling indicates that the overall effects of global warming on California

will include higher average temperatures in all seasons, higher total annual precipitation, and decreased spring and summer runoff due to decreases in snowpacks (EPA 1989, 1997). Decreases in spring and summer runoff could lead to the loss of breeding habitat for Yosemite toads and an increase in stranding mortality of eggs and tadpoles.

Changes in temperature may also affect virulence of pathogens to a different degree than the immune systems of amphibians (Carey *et al.* 1999), and may make Yosemite toads more susceptible to disease. An experimental increase in stream water temperature was shown to decrease density and biomass in invertebrates (Hogg and Williams 1996), thus global warming might have a negative impact on the Yosemite toad prey base.

Drought has contributed to the decline of Yosemite toads (Jennings and Hayes 1994), and the effects of climate change may also have contributed to that decline. These effects pose an ongoing, range-wide risk to the species.

Acid precipitation has been hypothesized as a cause of amphibian declines in the Sierra Nevada, because waters there are extremely low in acid neutralizing capacity, and therefore susceptible to changes in water chemistry due to acidic deposition (Bradford *et al.* 1994). Precipitation acidity in the Sierra Nevada has been documented to have significantly increased at a collection station at approximately 2,100 m (6,900 ft) elevation near Lake Tahoe (Byron *et al.* 1991). In addition to raising the acidity of water, acidic deposition may also cause increases in dissolved aluminum, because aluminum is more soluble at higher acidity. These increases in dissolved aluminum may be toxic to amphibians (Bradford *et al.* 1992). In laboratory experiments (Bradford *et al.* 1992; Bradford and Gordon 1992), high acidity and high aluminum concentrations did not have significant effects on survival of Yosemite toad embryos or newly hatched tadpoles. However, at pH 5.0 (pH represents acidity on a negative scale, with 7 being neutral and lower numbers being more acidic) and at high aluminum concentrations, Yosemite toad embryos hatched earlier and the tadpoles showed a reduction in body size. In a complementary field study of 235 randomly selected potential amphibian breeding sites (Bradford *et al.* 1994), no significant difference was found in pH between sites occupied and unoccupied by Yosemite toads. These data indicate that acid precipitation is an unlikely cause of decline in Yosemite toad

populations (Bradford *et al.* 1994). Therefore, acid deposition is considered a low risk to the species at this time, but should still be considered in conservation efforts because of the possibility of sublethal effects (Bradford *et al.* 1992), of its interaction with other factors, and the potential for more severe acidic deposition in the future.

Ambient ultraviolet-b (UV-B) radiation (280 to 320 nanometers (11.0 to 12.6 microinches)) has increased at north temperate latitudes in the past two decades (Adams *et al.* 2001). Ambient levels of UV-b were demonstrated to cause significant decreases in survival of western toad eggs in field experiments (Blaustein 1994). In a laboratory experiment (Kats *et al.* 2000), metamorph western toads exposed to levels of uv-b below those found in ambient sunlight showed a lower alarm response to chemical cues of injured toads than metamorphs that were completely shielded from UV-B. This indicates that ambient levels of UV-B may cause sublethal effects on toad behavior that may increase their vulnerability to predation. In a field experiment (Kiesecker and Blaustein 1995), the synergistic effects of exposure to ambient levels of UV-B radiation, and exposure to a pathogenic fungus (*Saprolegnia*), were shown to cause significantly higher mortality of western toad embryos than either factor alone.

Sadinsky *et al.* (1997) observed a high percentage of embryo mortality in Yosemite toads at six breeding sites in Yosemite National Park, but in a preliminary field experiment this mortality did not appear to be related to UV-B. In spatial statistical analysis of extant and extinct populations, higher elevation was shown to have a positive effect on the likelihood that populations of Yosemite toads were extant. This is counter to what would be expected if UV-B were the primary cause of decline (C. Davidson, *in litt.*, 2002), as sites at higher elevations would be expected to

receive more solar radiation due to the thinner atmosphere. The increase in UV-B at high elevations in the Sierra Nevada has not been more than 5 percent in the past several decades (Jennings 1996). These data indicate that UV-B has probably not contributed significantly to the decline of Yosemite toads and is probably currently a low risk to the species. However, as with acid precipitation, UV-B should still be considered as a risk to the species because of the potential for sublethal effects, synergistic effects with other factors, and the potential for further increases in UV-B radiation in the future.

Finding

We have carefully assessed the best scientific and commercial information available regarding the past, present, and future threats faced by this species. We reviewed the petition, information available in our files, and other published and unpublished information submitted to us during the public comment period following our 90-day petition finding. We also consulted with recognized Yosemite toad experts and other Federal and State resource agencies. On the basis of the best scientific and commercial information available, we find that proposing to list the Yosemite toad is warranted, but is precluded by higher priority listing actions.

In making this finding, we recognize that there have been declines in the distribution and abundance of Yosemite toads, primarily attributed to habitat degradation, airborne contaminants, and drought.

We conclude that the overall magnitude of threats to the Yosemite toad is moderate, and that the overall immediacy of these threats is non-imminent. Pursuant to our Listing Priority System (48 FR 43098), a species for which threats are moderate and non-imminent is assigned a Listing Priority

Number of 11. While we conclude that proposing to list the Yosemite toad is warranted, an immediate proposal to list is precluded by other higher priority listing actions. During fiscal year 2003, we must spend all of our Listing Program funding to comply with court orders and judicially approved settlement agreements, which are now our highest priority actions. The Yosemite toad will be added to the list of candidate species upon publication of this notice of 12-month finding. We will continue to monitor the status of this species and other candidate species. Should an emergency situation develop concerning this species, we will act to provide immediate protection, if warranted.

We intend that any proposed listing action for the Yosemite toad will be as accurate as possible. Therefore, we will continue to accept additional information and comments from all concerned governmental agencies, the scientific community, industry, or any other interested party concerning this finding. We are especially interested in further genetic information on the proper taxonomic status of the Yosemite toad and further information on the current range and status of the species, factors contributing to its decline, and conservation efforts.

References Cited

A complete list of all references cited is available on request from the Sacramento Fish and Wildlife Office (*see ADDRESSES* section, above).

Authority

The authority for this action is the Endangered Species Act of 1973, as amended (16 U.S.C. 1531 *et seq.*).

Dated: November 27, 2002.

Steve Williams,

Director, Fish and Wildlife Service.

[FR Doc. 02-30800 Filed 12-9-02; 8:45 am]

BILLING CODE 4310-55-P