



FEDERAL REGISTER

Vol. 79

Friday,

No. 2

January 3, 2014

Part III

Department of Commerce

National Oceanic and Atmospheric Administration

Takes of Marine Mammals Incidental to Specified Activities; Low-Energy Marine Geophysical Survey in the Dumont d'Urville Sea Off the Coast of East Antarctica, January to March 2013; Notice

DEPARTMENT OF COMMERCE**National Oceanic and Atmospheric Administration**

RIN 0648–XC779

Takes of Marine Mammals Incidental to Specified Activities; Low-Energy Marine Geophysical Survey in the Dumont d'Urville Sea Off the Coast of East Antarctica, January to March 2013

AGENCY: National Marine Fisheries Service (NMFS), National Oceanic and Atmospheric Administration (NOAA), Commerce.

ACTION: Notice; proposed Incidental Harassment Authorization; request for comments.

SUMMARY: NMFS has received an application from the National Science Foundation (NSF) Division of Polar Programs, and Antarctic Support Contract (ASC) on behalf of five research institutions: Colgate University, Columbia University, Texas A&M Research Foundation, University of South Florida, and University of Texas at Austin, for an Incidental Harassment Authorization (IHA) to take marine mammals, by harassment, incidental to conducting a low-energy marine geophysical (seismic) survey in the Dumont d'Urville Sea off the coast of East Antarctica, January to March 2014. Pursuant to the Marine Mammal Protection Act (MMPA), NMFS is requesting comments on its proposal to issue an IHA to NSF to incidentally harass, by Level B harassment only, 14 species of marine mammals during the specified activity.

DATES: Comments and information must be received no later than February 3, 2014.

ADDRESSES: Comments on the application should be addressed to P. Michael Payne, Chief, Permits and Conservation Division, Office of Protected Resources, National Marine Fisheries Service, 1315 East-West Highway, Silver Spring, MD 20910. The mailbox address for providing email comments is ITP.Goldstein@noaa.gov. NMFS is not responsible for email comments sent to addresses other than the one provided here. Comments sent via email, including all attachments, must not exceed a 10-megabyte file size.

All comments received are a part of the public record and will generally be posted to <http://www.nmfs.noaa.gov/pr/permits/incidental.htm#applications> without change. All Personal Identifying Information (for example, name, address, etc.) voluntarily submitted by the commenter may be publicly

accessible. Do not submit confidential business information or otherwise sensitive or protected information.

A copy of the application containing a list of the references used in this document may be obtained by writing to the above address, telephoning the contact listed here (see **FOR FURTHER INFORMATION CONTACT**) or visiting the internet at: <http://www.nmfs.noaa.gov/pr/permits/incidental.htm#applications>.

NSF and ASC have provided a "Draft Initial Environmental Evaluation/Environmental Assessment to Conduct a Marine-Based Studies of the Totten Glacier System and Marine Record of Cryosphere—Ocean Dynamics" (IEE/EA), prepared by AECOM, on behalf of NSF and ASC, which is also available at the same Internet address. Documents cited in this notice may be viewed, by appointment, during regular business hours, at the aforementioned address.

FOR FURTHER INFORMATION CONTACT: Howard Goldstein or Jolie Harrison, Office of Protected Resources, NMFS, 301–427–8401.

SUPPLEMENTARY INFORMATION:**Background**

Section 101(a)(5)(D) of the MMPA, as amended (16 U.S.C. 1371 (a)(5)(D)), directs the Secretary of Commerce (Secretary) to authorize, upon request, the incidental, but not intentional, taking of small numbers of marine mammals of a species or population stock, by United States citizens who engage in a specified activity (other than commercial fishing) within a specified geographical region if certain findings are made and, if the taking is limited to harassment, a notice of a proposed authorization is provided to the public for review.

Authorization for the incidental taking of small numbers of marine mammals shall be granted if NMFS finds that the taking will have a negligible impact on the species or stock(s), and will not have an unmitigable adverse impact on the availability of the species or stock(s) for subsistence uses (where relevant). The authorization must set forth the permissible methods of taking, other means of effecting the least practicable adverse impact on the species or stock and its habitat, and requirements pertaining to the mitigation, monitoring and reporting of such takings. NMFS has defined "negligible impact" in 50 CFR 216.103 as ". . . an impact resulting from the specified activity that cannot be reasonably expected to, and is not reasonably likely to, adversely affect the species or stock through effects on annual rates of recruitment or survival."

Section 101(a)(5)(D) of the MMPA established an expedited process by which citizens of the United States can apply for an authorization to incidentally take small numbers of marine mammals by harassment. Section 101(a)(5)(D) of the MMPA establishes a 45-day time limit for NMFS's review of an application followed by a 30-day public notice and comment period on any proposed authorizations for the incidental harassment of small numbers of marine mammals. Within 45 days of the close of the public comment period, NMFS must either issue or deny the authorization.

Except with respect to certain activities not pertinent here, the MMPA defines "harassment" as: Any act of pursuit, torment, or annoyance which (i) has the potential to injure a marine mammal or marine mammal stock in the wild [Level A harassment]; or (ii) has the potential to disturb a marine mammal or marine mammal stock in the wild by causing disruption of behavioral patterns, including, but not limited to, migration, breathing, nursing, breeding, feeding, or sheltering [Level B harassment].

Summary of Request

On July 3, 2013, NMFS received an application from the NSF and ASC requesting that NMFS issue an IHA for the take, by Level B harassment only, of small numbers of marine mammals incidental to conducting a low-energy marine seismic survey in International Waters (i.e., high seas) and in the Southern Ocean off the coast of East Antarctica during January to March 2014. We received an addendum to the application from the NSF and ASC on December 18, 2013 which reflected updates to incidental take requests for marine mammals related to icebreaking activities.

The research would be conducted by five research institutions: Colgate University, Columbia University, Texas A&M Research Foundation, University of South Florida, and University of Texas at Austin. The NSF and ASC plans to use one source vessel, the R/VIB *Nathaniel B. Palmer* (*Palmer*), and a seismic airgun array to collect seismic data in the Southern Ocean. The vessel would be operated by ASC, which operates the United States Antarctic Program under contract to the NSF. In support of the United States Antarctic Program, the NSF and ASC plans to use conventional low-energy, seismic methodology to perform marine-based studies in the Dumont d'Urville Sea to include evaluation of geophysical and physical oceanographic features in two

areas along the coast of East Antarctica (see Figures 1, 2, and 3 of the IHA application). The primary area proposed for the study is the Totten Glacier system (preferred study area) including the Moscow University Ice Shelf along the Sabrina Coast, and a secondary area, the Mertz Glacier and Cook Ice Shelf, along the Oates Coast. In addition to the proposed operations of the seismic airgun array and hydrophone streamer, NSF and ASC intend to operate a single-beam echosounder, multi-beam echosounder, acoustic Doppler current profiler (ADCP), and sub-bottom profiler continuously throughout the survey.

Acoustic stimuli (i.e., increased underwater sound) generated during the operation of the seismic airgun array and from icebreaking activities may have the potential to cause a behavioral disturbance for marine mammals in the survey area. This is the principal means of marine mammal taking associated with these activities, and NSF and ASC has requested an authorization to take 14 species of marine mammals by Level B harassment. Take is not expected to result from the use of the single-beam echosounder, multi-beam echosounder, ADCP, acoustic locator, and sub-bottom profiler, as the brief exposure of marine mammals to one pulse, or small numbers of signals, in this particular case is not likely to result in the harassment of marine mammals. Also, NMFS does not expect take to result from collision with the source vessel because it is a single vessel moving at a relatively slow, constant cruise speed of 5 knots [kts]; 9.3 kilometers per hour [km/hr]; 5.8 miles per hour [mph] during seismic acquisition within the survey, for a relatively short period of time (approximately 45 operational days). It is likely that any marine mammal would be able to avoid the vessel.

Description of the Proposed Specified Activity

NSF and ASC propose to conduct a low-energy seismic survey in the Dumont d'Urville Sea in the Southern Ocean off the coast of East Antarctica from January to March 2014. In addition to the low-energy seismic survey, scientific activities would include conducting a bathymetric profile survey of the seafloor using transducer based instruments such as a multi-beam echosounder and sub-bottom profiler; conducting magnetometry and imaging surveys using an underwater camera assembly; collecting sediment cores and dredge sampling; and collecting water samples and conductivity (salinity), temperature, depth (CTD) and current data through the deployment and

recovery of short-term (in place for approximately one month) and long-term (in place for approximately one year) instrumentation moorings, CTD equipment casts, and the use of transducer-based ADCP instruments. Sea ice conditions will dictate areas where the ship and airguns can operate. Due to dynamic ice conditions, which cannot be predicted on a local scale, it is not possible to develop tracklines a priori. The seismic survey would be conducted in one or both of the two study areas depending on the sea ice conditions; however, the preferred study area is the Totten Glacier region (see Figure 2 of the IHA application). Water depths in the survey area range from 100 to 1,000 meters (m) (328.1 to 3,280.1 feet [ft]), and possibly exceeding 1,000 m in some areas. The seismic surveys are scheduled to occur for a total of less than or equal to 300 hours at one or both of the two study areas for approximately 45 operational days in January to March 2014. The operation hours and survey length would include equipment testing, ramp-up, line changes, and repeat coverage. The long transit time between port and the study site constrains how long the ship can be in the study area and effectively limits the maximum amount of time the airguns can operate. Some minor deviation from these dates would be possible, depending on logistics and weather.

The proposed survey of Totten Glacier and Moscow University Ice Shelf along the Sabrina Coast continental shelf is designed to address several critical questions. The Totten Glacier system, which drains one-eighth of the East Antarctic Ice Sheet and contains more ice volume than the entire West Antarctic Ice Sheet, remains the single largest and least understood glacial system which possesses a potentially unsteady dynamic. If it were to melt, sea-level would rise by more than 5 m (16.4 ft) worldwide. The proposed marine studies would help to understand both the dynamics and the controls of the Totten Glacier system, and to resolve ambiguity in large ice mass dynamic behavior. This research would be accomplished via the collection of glaciological, geological, and physical oceanographic data. In order to place the modern system, as well as more recent changes to the system, into a longer-term perspective, researchers would collect and interpret marine geologic, geochemical, and geophysical records of the longer term behavior and response of this system.

The proposed research would complement fieldwork studying other Antarctic ice shelves oceanographic

studies near the Antarctic Peninsula, and ongoing development of ice sheet and other ocean models. It would facilitate learning at sea and ashore by students, help to fill important spatial and temporal gaps in a sparsely sampled region of coastal Antarctica, and communicate its findings via publications and outreach. Obtaining records of currents and oceanographic properties in this region are consistent with the objectives of the Southern Ocean Observing System for climate change. The work would enhance general understanding of air-sea-ice interactions, ocean circulation, ice shelf sensitivity to climate change, and the present and future roles of East Antarctic Ice Sheet on sea level.

The Principal Investigators are Dr. Amy Leventer of Colgate University, Dr. Donald Blankenship and Dr. Sean Gulick of the University of Texas at Austin, Dr. Eugene Domack of the University of South Florida, Mr. Bruce Huber of Columbia University, and Dr. Alejandro Orsi of Texas A&M Research Foundation.

The procedures to be used for the surveys would be similar to those used during previous low-energy seismic surveys by NSF and would use conventional seismic methodology. The proposed survey will involve one source vessel, the R/V *Nathaniel B. Palmer* (*Palmer*). NSF and ASC will deploy two (each with a discharge volume of 45 cubic inch [in³] with a total volume of 90 in³ or each with a discharge volume of 105 in³ with a total volume of 210 in³) Sercel Generator Injector (GI) airgun array as an energy source at a tow depth of up to 3 m (9.8 ft) below the surface (more information on the airguns can be found in Appendix B of the IHA application). The receiving system will consist of one 100 m (328.1 ft) long, 24-channel, solid-state hydrophone streamer towed behind the vessel. As the GI airguns are towed along the survey lines, the hydrophone streamer will receive the returning acoustic signals and transfer the data to the onboard processing system. All planned seismic data acquisition activities will be conducted by technicians provided by NSF and ASC with onboard assistance by the scientists who have proposed the study. The vessel will be self-contained, and the crew will live aboard the vessel for the entire cruise.

The planned seismic survey (e.g., equipment testing, start-up, line changes, repeat coverage of any areas, and equipment recovery) will consist of approximately 2,800 kilometer (km) (1,511.9 nautical miles [nmi]) of transect lines (including turns) in the survey area in the Dumont d'Urville Sea of the

Southern Ocean (see Figures 1, 2, and 3 of the IHA application). In addition to the operation of the airgun array, a single-beam and multi-beam echosounder, ADCP, and a sub-bottom profiler will also likely be operated from

the *Palmer* continuously throughout the cruise between the first and last survey sites. There will be additional seismic operations associated with equipment testing, ramp-up, and possible line changes or repeat coverage of any areas

where initial data quality is sub-standard. In NSF and ASC's estimated take calculations, 25% has been added for those additional operations.

TABLE 1—PROPOSED LOW-ENERGY SEISMIC SURVEY ACTIVITIES IN THE DUMONT D'URVILLE SEA OFF THE COAST OF EAST ANTARCTICA

Survey length (km)	Cumulative duration (hr) ¹	Airgun array total volume	Time between airgun shots (distance)	Streamer length (m)
2,800 (1,511.9 nmi)	≤300	2 × 45 in ³ (2 × 737 cm ³) or 2 × 105 in ³ (2 × 1,720 cm ³)	5 seconds (12.5 m or 41 ft) ...	100 (328.1 ft).

¹ Airgun operations are planned for no more than 16 continuous hours at a time.

Vessel Specifications

The *Palmer*, a research vessel owned by Edison Chouest Offshore, Inc. and operated by NSF and ACS (under a long-term charter with Edison Chouest Offshore, Inc.), will tow the two GI airgun array, as well as the hydrophone streamer. When the *Palmer* is towing the airgun array and the relatively short hydrophone streamer, the turning rate of the vessel while the gear is deployed is much higher than the limit of 5 degrees per a minute for a seismic vessel towing a streamer of more typical length (much greater than 1 km [0.5 nmi]), which is approximately 20 degrees. Thus, the maneuverability of the vessel is not limited much during operations with the streamer.

The U.S.-flagged vessel has a length of 94 m (308.5 ft); a beam of 18.3 m (60 ft); a maximum draft of 6.8 m (22.5 ft); and a gross tonnage of 6,174. The ship is powered by four Caterpillar 3608 diesel engines (3,300 brake horsepower [hp] at 900 rotations per minute [rpm]) and a 1,400 hp flush-mounted, water jet azimuthing bowthruuster. Electrical power is provided by four Caterpillar 3512, 1,050 kiloWatt (kW) diesel generators. The *Palmer's* operation speed during seismic acquisition is typically approximately 9.3 km/hr (5 kts) (varying between 7.4 to 11.1 km/hr [4 to 6 kts]). When not towing seismic survey gear, the *Palmer* typically cruises at 18.7 km/hr (10.1 kts) and has a maximum speed of 26.9 km/hr (14.5 kts). The *Palmer* has an operating range of approximately 27,780 km (15,000 nmi) (the distance the vessel can travel without refueling), which is approximately 70 to 75 days. The vessel can accommodate 37 scientists and 22 crew members.

The vessel also has two locations as likely observation stations from which Protected Species Observers (PSO) will watch for marine mammals before and during the proposed airgun operations

on the *Palmer*. Observing stations will be at the bridge level with PSO's eye level approximately 16.5 m (54.1 ft) above sea level with an approximately 270° view around the vessel, and an aloft observation tower that is approximately 24.4 m (80.1 ft) above sea level that is protected from the weather and has an approximately 360° view around the vessel. More details of the *Palmer* can be found in the IHA application and online at: <http://www.nsf.gov/geo/plr/support/nathpalm.jsp> and <http://www.usap.gov/vesselScienceAndOperations/contentHandler.cfm?id=1561>.

Acoustic Source Specifications

Seismic Airguns

The *Palmer* will deploy an airgun array, consisting of two 45 in³ or two 105 in³ GI airguns as the primary energy source and a 100 m streamer containing hydrophones. The airgun array will have a supply firing pressure of 2,000 pounds per square inch (psi) and 2,200 psi when at high pressure stand-by (i.e., shut-down). The regulator is adjusted to ensure that the maximum pressure to the GI airguns is 2,000 psi, but there are times when the GI airguns may be operated at pressures as low as 1,750 to 1,800 psi. Seismic pulses for the GI airguns will be emitted at intervals of approximately 5 seconds. At speeds of approximately 9.3 km/hr, the shot intervals correspond to spacing of approximately will be 12.5 m (41 ft) during the study. There would be approximately 720 shots per hour. During firing, a brief (approximately 0.03 second) pulse sound is emitted; the airguns will be silent during the intervening periods. The dominant frequency components range from two to 188 Hertz (Hz).

The GI airguns would be used in harmonic mode, that is, the volume of the injector chamber (I) of each GI airgun is equal to that of its generator

chamber (G): 45 in³ and 105 in³ for each airgun array. Each airgun would be initially configured to a displacement volume of 45 in³ for the generator and injector. The generator chamber of each GI airgun in the primary source, the one responsible for introducing the sound pulse into the ocean, is 45 in³. The injector chamber injects air into the previously-generated bubble to maintain its shape, and does not introduce more sound into the water. The airguns would fire the compressed air volume in unison in a harmonic mode. In harmonic mode, the injector volume is designed to destructively interfere with the reverberations of the generator (source component). Firing the airguns in harmonic mode maximizes resolution in the data and minimizes any excess noise in the water column or data caused by the reverberations (or bubble pulses). The two GI airguns will be spaced approximately 3 or 6 m (9.8 or 19.7 ft) apart, side-by-side, between 15 and 40 m (49.2 and 131.2 ft) behind the *Palmer*, at a depth of up to 3 m during the surveys. If needed to improve penetration of the strata, the two airguns may be reconfigured to a displacement volume of 105 in³ each and would still be considered a low-energy acoustic source as defined in the NSF/USGS PEIS. Therefore, there are three possible two airgun array configurations: Two 45/45 in³ airguns separated by 3 m, two 45/45 in³ airguns separated by 6 m, and two 105/105 in³ airguns separated by 3 m. The two 45/45 in³ airguns separated by 3 m layout is preferred, the two 45/45 in³ separated by 6 m layout would be used in the event the middle of the three 45/45 in³ airgun fails, and the two 105/105 in³ airguns separated by 3 m would be used only if additional penetration is needed. To summarize, two strings of GI airguns would be available: (1) Three 45/45 in³ airguns on a single string where one of these is used as a "hot spare" in the event of

failure of one of the other two airguns, these three GI airguns are separated by 3 m; and (2) two 105/105 in³ airguns on a second string without a "hot spare." The total effective volume will be 90 or 210 in³. The two strings would be spaced 14 m (45.9 ft) apart, on either side of the midline of the vessel, however, only one string at a time would be used.

The Nucleus modeling software used at Lamont-Doherty Earth Observatory of Columbia University (L-DEO) does not include GI airguns as part of its airgun library, however signatures and mitigation models have been obtained for two 45 in³ G airguns at 2 m tow depth and two 105 in³ G airguns at 3 m tow depth that are close approximations. For the two 45 in³ airgun array, the source output (downward) is 230.6 dB re: 1 μPam for 0-to-peak and 235.9 dB re: 1 μPam for peak-to-peak. For the two 105 in³ airgun array, the source output (downward) is 234.4 dB re: 1 μPam 0-to-peak and 239.8 dB re: 1 μPam for peak-to-peak. These numbers were determined using the aforementioned G-airgun approximation to the GI airgun and using signatures filtered with DFS V out-256 Hz 72 dB/octave. The dominant frequency range would be 20 to 160 Hz for a pair of GI airguns towed at 3 m depth and 35 to 230 Hz for a pair of GI airguns towed at 2 m depth.

During the low-energy seismic survey, the vessel would attempt to maintain a constant cruise speed of approximately 5 knots. The airguns would operate continuously for no more than 16 hours at a time and duration of continuous operation is dependent on ice concentration. The cumulative duration of the airgun operations will not exceed 200 hrs. The relatively short, 24-channel hydrophone streamer would provide operational flexibility to allow the seismic survey to proceed along the designated cruise track with minimal interruption due to variable sea ice conditions. The design of the seismic equipment is to achieve high-resolution images of the glacial marine sequence stratigraphy with the ability to correlate to the ultra-high frequency sub-bottom profiling data and provide cross-sectional views to pair with the seafloor bathymetry. The cruise path would be designated once in the study area and would take care to avoid heavy ice conditions such as icebergs or dense areas of pack ice that could potentially damage the airguns or streamer and minimize proximity to potential marine receptors.

Weather conditions that could affect the movement of sea ice and hinder the hydrophone streamer would be closely

monitored, as well as conditions that could limit visibility. If situations are encountered which pose a risk to the equipment, impede data collection, or require the vessel to stop forward progress, the seismic survey equipment would be shut-down and retrieved until conditions improve. In general, the hydrophone streamer and sources could be retrieved in less than 30 minutes.

Metrics Used in This Document

This section includes a brief explanation of the sound measurements frequently used in the discussions of acoustic effects in this document. Sound pressure is the sound force per unit area, and is usually measured in micropascals (μPa), where 1 pascal (Pa) is the pressure resulting from a force of one newton exerted over an area of one square meter. Sound pressure level (SPL) is expressed as the ratio of a measured sound pressure and a reference level. The commonly used reference pressure level in underwater acoustics is 1 μPa, and the units for SPLs are dB re: 1 μPa. SPL (in decibels [dB]) = 20 log (pressure/reference pressure).

SPL is an instantaneous measurement and can be expressed as the peak, the peak-to-peak (p-p), or the root mean square (rms). Root mean square, which is the square root of the arithmetic average of the squared instantaneous pressure values, is typically used in discussions of the effects of sounds on vertebrates and all references to SPL in this document refer to the root mean square unless otherwise noted. SPL does not take the duration of a sound into account.

Characteristics of the Airgun Pulses

Airguns function by venting high-pressure air into the water which creates an air bubble. The pressure signature of an individual airgun consists of a sharp rise and then fall in pressure, followed by several positive and negative pressure excursions caused by the oscillation of the resulting air bubble. The oscillation of the air bubble transmits sounds downward through the seafloor and the amount of sound transmitted in the near horizontal directions is reduced. However, the airgun array also emits sounds that travel horizontally toward non-target areas.

The nominal downward-directed source levels of the airgun arrays used by NSF and ASC on the *Palmer* do not represent actual sound levels that can be measured at any location in the water. Rather, they represent the level that would be found 1 m (3.3 ft) from a hypothetical point source emitting the

same total amount of sound as is emitted by the combined GI airguns. The actual received level at any location in the water near the GI airguns will not exceed the source level of the strongest individual source. In this case, that will be about 224.6 dB re 1 μPam peak, or 229.8 dB re 1 μPam peak-to-peak for the two 45 in³ airgun array, and 228.2 dB re 1 μPam peak or 233.5 dB re 1 μPam peak-to-peak for the two 105 in³ airgun array. However, the difference between rms and peak or peak-to-peak values for a given pulse depends on the frequency content and duration of the pulse, among other factors. Actual levels experienced by any organism more than 1 m from either GI airgun will be significantly lower.

Accordingly, Lamont-Doherty Earth Observatory of Columbia University (L-DEO) has predicted and modeled the received sound levels in relation to distance and direction from the two GI airgun array. A detailed description of L-DEO's modeling for this survey's marine seismic source arrays for protected species mitigation is provided in the NSF/USGS PEIS. These are the nominal source levels applicable to downward propagation. The NSF/USGS PEIS discusses the characteristics of the airgun pulses. NMFS refers the reviewers to those documents for additional information.

Predicted Sound Levels for the Airguns

To determine exclusion zones for the airgun array to be used in the intermediate and deep water of the Gulf of Mexico (GOM), received sound levels have been modeled by L-DEO for a number of airgun configurations, including two 45 in³ and two 105 in³ G airguns, in relation to distance and direction from the airguns (see Figure 2 and 3 in Attachment B of the IHA application). The model does not allow for bottom interactions, and is most directly applicable to deep water. Because the model results are for G airguns, which have more energy than GI airguns of the same size, those distances overestimate (by approximately 10%) the distances for the two 45 in³ GI airguns and two 105 in³ GI airguns, respectively. Although the distances are overestimated, no adjustments for this have been made to the radii distances in Table 2 (below). Based on the modeling, estimates of the maximum distances from the GI airguns where sound levels of 190, 180, and 160 dB re 1 μPa (rms) are predicted to be received in shallow, intermediate, and deep water are shown in Table 2 (see Table 1 of Attachment B of the IHA application).

Empirical data concerning the 190, 180, and 160 dB (rms) distances were acquired for various airgun arrays based on measurements during the acoustic verification studies conducted by L-DEO in the northern GOM in 2003 (Tolstoy *et al.*, 2004) and 2007 to 2008 (Tolstoy *et al.*, 2009; Diebold *et al.*, 2010). Results of the 18 and 36 airgun array are not relevant for the two GI airguns to be used in the proposed survey. The empirical data for the 6, 10, 12, and 20 airgun arrays indicate that, for deep water, the L-DEO model tends to overestimate the received sound levels at a given distance (Tolstoy *et al.*, 2004). Measurements were not made for the two GI airgun array in deep water; however, NSF and ASC proposes to use

the buffer and exclusion zones predicted by L-DEO's model for the proposed GI airgun operations in deep water, although they are likely conservative given the empirical results for the other arrays. Using the L-DEO model, Table 2 (below) shows the distances at which three rms sound levels are expected to be received from the two GI airguns. The 180 and 190 dB re 1 μ Pam (rms) distances are the safety criteria for potential Level A harassment as specified by NMFS (2000) and are applicable to cetaceans and pinnipeds, respectively. If marine mammals are detected within or about to enter the appropriate exclusion zone, the airguns will be shut-down immediately.

Table 2 summarizes the predicted distances at which sound levels (160, 180, and 190 dB [rms]) are expected to be received from the two airgun array (45 in³ or 105 in³) operating in shallow (less than 100 m [328 ft]), intermediate (100 to 1,000 m [328 to 3,280 ft]), and deep water (greater than 1,000 m [3,280 ft]) depths.

Table 2— Predicted and modeled (two 45 in³ and two 105 in³ GI airgun array) distances to which sound levels \geq 190, 180 and 160 dB re: 1 μ Pa (rms) could be received in shallow, intermediate, and deep water during the proposed low-energy seismic survey in the Dumont d'Urville Sea of the Southern Ocean, January to March 2014.

Source and total volume	Tow depth (m)	Water depth (m)	Predicted RMS radii distances (m) for 2 GI airgun array		
			160 dB	180 dB	190 dB
Two GI Airguns (45 in ³)	3	Shallow (<100)	1,176 (3,858.3 ft)	296 (971.1 ft)	147 (482.3 ft)
Two GI Airguns (45 in ³)	3	Intermediate (100 to 1,000)	600 (1,968.5 ft)	100 (328ft)	15 (49.2 ft)
Two GI Airguns (45 in ³)	3	Deep (\leq 1,000)	400 (1,312.3 ft)	100 (328 ft)	10 (32.8 ft)
Two GI Airguns (105 in ³)	3	Shallow (<100)	1,970 (6,463.3 ft)	511 (1,676.5 ft)	294 (964.6 ft)
Two GI Airguns (105 in ³)	3	Intermediate (100 to 1,000)	1,005 (3,297.2 ft)	100	30 (98.4 ft)
Two GI Airguns (105 in ³)	3	Deep (>1,000)	670 (2,198.2 ft)	100	20 (65.6 ft)

NMFS expects that acoustic stimuli resulting from the proposed operation of the two GI airgun array has the potential to harass marine mammals. NMFS does not expect that the movement of the *Palmer*, during the conduct of the low-energy seismic survey, has the potential to harass marine mammals because of the relatively slow operation speed of the vessel (approximately 5 kts; 9.3 km/hr; 5.8 mph) during seismic acquisition.

Bathymetric Survey

Along with the low-energy airgun operations, other additional geophysical measurements would be made using swath bathymetry, backscatter sonar imagery, high-resolution sub-bottom profiling ("CHIRP"), imaging, and magnetometer instruments. In addition, several other transducer-based instruments onboard the vessel would be operated continuously during the cruise for operational and navigational purposes. Operating characteristics for the instruments to be used are described below.

Single-Beam Echosounder (Knudsen 3260)—The hull-mounted CHIRP sonar would be operated continuously during all phases of the cruise. This instrument

is operated at 12 kHz for bottom-tracking purposes or at 3.5 kHz in the sub-bottom profiling mode. The sonar emits energy in a 30° beam from the bottom of the ship.

Single-Beam Echosounder (Bathy 2000)—The hull-mounted sonar characteristics of the Bathy 2000 are similar to the Knudsen 3260. Only one hull-mounted echosounder can be operated a time, and this source would be operated instead of the Knudsen 3260 only if needed (i.e., only one would be in continuous operation during the cruise).

Multi-Beam Sonar (Simrad EM120)—The hull-mounted multi-beam sonar would be operated continuously during the cruise. This instrument operates at a frequency of 12 kHz, has an estimated maximum source energy level of 242 dB re 1 μ Pa (rms), and emits a very narrow (<2°) beam fore to aft and 150° in cross-track. The multi-beam system emits a series of nine consecutive 15 ms pulses.

Acoustic Doppler Current Profiler (ADCP Teledyne RDI VM-150)—The hull-mounted ADCP would be operated continuously throughout the cruise. The ADCP operates at a frequency of 150 kHz with an estimated acoustic output

level at the source of 223.6 dB re 1 μ Pa (rms). Sound energy from the ADCP is emitted as a 30° conically-shaped beam.

Acoustic Doppler Current Profiler (ADCP Ocean Surveyor OS-38)—The characteristics of this backup hull-mounted ADCP unit are similar to the Teledyne VM-150 and would be continuously operated.

Acoustic Locator (Pinger)—An acoustic locator (i.e., pinger) would be deployed when using the Smith-McIntyre grab sampler and multi-corer (Mega-corer) to enable these devices to be located in the event they become detached from their lines. A pinger typically operates at a frequency of 12 kHz, generates a 5 ms pulse per second, and has an acoustical output of 162 dB re 1 μ Pa (rms). A maximum total of 30 samples would be obtained using these devices and require approximately one hour per sample; therefore, the pinger would operate for a total of 30 hours.

Passive Instruments—During the seismic survey in the Dumont d'Urville Sea, a precession magnetometer and Air-Sea gravity meter would be deployed. In addition, numerous (approximately 24) expendable bathythermograph (XBTs) probes would

also be released (and none would be recovered) over the course of the cruise to obtain temperature data necessary to calculate sound velocity profiles used by the multi-beam sonar.

Core and Dredge Sampling

The primary sampling goals involve the acquisition of marine sediment cores of various lengths up to 25 m (82 ft). It is anticipated that up to 65 sediment cores and grab samples and 12 rock dredge samples would be collected as summarized in Table 3 (Table 3 of the IHA application). Each core or grab sample would require approximately one hour per sample. All cores and dredges would be deployed using a steel cable/winch system.

Approximately 75 m² (807.3 ft²) of seafloor would be disturbed by each of four deployments of the dredge at three different sites (resulting in a total of 900 m² [9,687.5 ft²] of affected seafloor for the project). The selection of the bottom sampling locations and sampling method would be based on observations of the seafloor, subsurface reflectivity, sediment type, and accessibility due to ice and weather conditions. Bottom sampling in the Mertz Glacier area would be limited to strategically selected locations including possible re-sampling at a previous core site.

TABLE 3—PROPOSED CORING AND DREDGING ACTIVITIES IN THE DUMONT D'URVILLE SEA

Sampling device	Number of deployments
Smith-McIntyre grab sampler	10 to 15.
Multi-corer (Mega-corer)	10 to 15.
Kasten corer (regular or jumbo).	20 to 25.
Jumbo piston corer	8 to 10.
Box cage dredge	10 to 12.

Limited sampling of rock material would be conducted using a dredge that would be towed along the seafloor for short distances (approximately 50 m [164 ft]) to collect samples of bedrock and ice rafted debris. The available dredges, which have openings of 0.5 to 1.5 m (1.6 to 4.9 ft), would be deployed on rocky substrates. The locations of the proposed dredge sites are limited to the inner shelf (southern) perimeter of three areas: The Mertz Trough and two regions along the Sabrina Coast. Final selection of dredge sites will include review to ensure that the seamounts or corals in the area are avoided (AOA, 2011).

The Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR) has adopted conservation measures (i.e., 22–06, 22–07, and 22–09)

to protect vulnerable marine ecosystems (VME), which include seamounts, hydrothermal vents, cold water corals, and sponge fields. The conservation measure 22–07 includes mitigation and reporting requirements if VME are encountered. The science team would follow these requirements (see Attachment C of the IHA application) if VME's are encountered while sampling the sea bottom.

In addition, a camera and towed video system would be deployed at up to 25 sites. This device would lightly touch the seafloor to establish a baseline and rise to an optimum elevation to obtain the desired images.

Water Sampling and Current Measurements

High-resolution conductivity, depth, and temperature (CTD) measurements would be collected to characterize the summer regional water mass stratification and circulation, and the meridional exchange of waters between the oceanic and shelf regimes. These physical measurements would involve approximately SeaBird CTD system casts including the use of a lowered ADCP (LADCP).

The LADCP would consist of two Teledyne RDI Workhorse Monitor ADCPs mounted on the CTD/rosette frame and one oriented upward and the other downward. The LADCP and frame would be raised and lowered by cable and winch. The LADCPs would operate at a frequency of 307.2 kHz, with an estimated output acoustic pressure along each 4 beams of 216.3 dB re 1μPa at 1 m. The beams are angled at 20 degrees from the centerline of the ADCP head, with a beam angle of 4 degrees for the individual beams. Typical pulse duration is 5.7 ms, with a typical repetition rate of 1.75 s. The upward and downward-looking ADCPs are operated in master-slave mode so that only one head pings at a time. The LADCP would be operated approximately one hour at every CTD/rosette station (maximum of 100 stations) for a total of 100 hours of operation.

These instruments would be used to profile the full water column for temperature, salinity (conductivity), dissolved oxygen and currents at a series of transects in the study area. Discrete water samples would be collected for salinity and dissolved oxygen to monitor CTD/rosette performance, and for oxygen isotopes to assess meltwater content. Water samples would also be collected for development and interpretation of marine sediment proxies using Niskin bottles.

Observations of the thermal structure along other portions of the cruise track would be made using an underway CTD system and XBTs while the seafloor is swath-mapped. The number and spacing of stations would be adjusted according to ocean features discovered through multi-beam swath mapping and the sea ice conditions. If portions of the study area are inaccessible to the NBP, a contingency sampling focused on the inflows of MDCW would be pursued in adjacent shelf troughs.

It is noted that underway ADCP on the *Palmer* can, under ideal conditions, obtain profiles of ocean currents to depths greater than 800 m (2,624.7 ft). On continental shelves where depths may be less than the range of the ADCP, the underway profiles cannot resolve the deepest 15% of the water column due to side lobe reflections from the bottom which contaminate the water column Doppler returns. For a depth of 800 m, expected in the MCDW, currents in the lower 120 m (393.7 ft) could not be measured by the ship ADCP; therefore, the lowered ADCP can provide accurate current profiles to within a few meters of the bottom and provide complete coverage of the velocity field at each CTD station.

Instrumentation Moorings

Four instrumented moorings would be deployed during the proposed cruise to measure current, temperature, and salinity (conductivity) continuously. Two of the moorings would be deployed for approximately one month (short-term moorings) and two moorings would be deployed for approximately one year (long-term moorings). The two short-term moorings and one long-term mooring would include ADCP paired with CTD recorders, and additional intermediate T (i.e., temperature) recorders. The characteristics of the ADCP units deployed on the moorings are similar to the Teledyne VM–150; the moored ADCPs operate at frequencies of 75 kHz (one unit) and 300 kHz (two units). The fourth mooring would be equipped with sediment traps, a CTD recorder and intermediate T recorders, and be deployed for approximately one year (long-term mooring). The two long-term moorings would be retrieved approximately one year later by a U.S. Arctic Program (USAP) vessel or collaborators from other countries.

Subject to sea ice conditions, these moorings would preferably be placed in front of Totten Glacier, but otherwise as close as possible inside adjacent cross-shelf troughs. If access to the inner shelf is not allowed by sea ice conditions we would attempt mooring deployments within the outer shelf close to the

troughs mouth, where the Totten Glacier is more directly connected to inflows from the oceanic domain offshore. The two long-term moorings would be deployed within 16 km of each other. The short-term moorings would be within a few kilometers of each other and no farther than 32 km (17.3 nmi) from the long-term moorings. All instruments would be kept at depths below 250 m (820.2 ft) to minimize damage or loss by icebergs.

The moorings would temporarily attached to anchors and be recovered using acoustic release mechanisms. The mooring recovery process would be similar regardless of mooring type or when they would be retrieved. Locating the moorings and releasing the moorings from the steel railroad wheel anchors (which would not be recovered) would be accomplished by transmitting sound over a period of several seconds. This is done with an acoustic deck command unit that sends a sequence of coded pulses to the receiving units, the acoustic releases, connected to the mooring anchors. The acoustic releases response to acknowledge the receipt of commands from the deck unit is by transmitting a short sequence of pulses back. Both of the acoustic units (onboard deck unit and moored releases) operate at frequencies between approximately 7 and 15 kHz. The beam pattern is approximately omnidirectional. The acoustic source level is less than 192 dB re 1 μ Pa at 1 m.

In addition to the U.S. moorings described above, three new moorings would be deployed on behalf of Australia's national science agency the Commonwealth of Scientific and Industrial Research Organisation (CSIRO) Physical Oceanography group in the Totten Glacier region by the project team. These moorings would be retrieved approximately one year later by collaborators from other countries. Also, during this cruise, three CSIRO moorings that were deployed over a year ago in the western outlet of the Mertz-Ninnis Trough would be recovered. The recovery process and acoustic sources described above for the U.S. moorings would be used for recovery of the CSIRO moorings.

Icebreaking

Icebreaking is considered by NMFS to be a continuous sound and NMFS estimates that harassment occurs when marine mammals are exposed to continuous sounds at a received sound level of 120 dB SPL or above. Potential takes of marine mammals may ensue from icebreaking activity in which the *Palmer* is expected to engage in

Antarctic waters (i.e., along the George V and Oates Coast of East Antarctica, >65° South, between 140° and 165° East). While breaking ice, the noise from the ship, including impact with ice, engine noise, and propeller cavitation, will exceed 120 dB (rms) continuously. If icebreaking does occur in Antarctic waters, NMFS, NSF and ASC expect it will occur during transit and non-seismic operations to gain access to coring, dredging, or other sampling locations and not during seismic airgun operations. The research activities and associated contingencies are designed to avoid areas of heavy sea ice condition. The buffer zone (160 dB [rms]) for the marine mammal Level B harassment threshold during the proposed activities is greater than the calculated radius during icebreaking. Therefore, if the *Palmer* breaks ice during seismic operations within the Antarctic waters (within the Dumont d'Urville Sea or other areas of the Southern Ocean), the more conservative and larger radius (i.e., that for seismic operations) will be used and supercede the buffer zone for icebreaking.

In 2008, acousticians from Scripps Institution of Oceanography Marine Physical Laboratory and University of New Hampshire Center for Coastal and Ocean Mapping conducted measurements of SPLs of the *Healy* icebreaking under various conditions (Roth and Schmidt, 2010). The results indicated that the highest mean SPL (185 dB) was measured at survey speeds of 4 to 4.5 kts in conditions of 5/10 ice and greater. Mean SPL under conditions where the ship was breaking heavy ice by backing and ramming was actually lower (180 dB). In addition, when backing and ramming, the vessel is essentially stationary, so the ensonified area is limited for a short period (on the order of minutes to tens of minutes) to the immediate vicinity of the vessel until the ship breaks free and once again makes headway.

The 120 dB received sound level radius around the *Healy* while icebreaking was estimated by researchers (USGS, 2010). Using a spherical spreading model, a source level of 185 dB decays to 120 dB in about 1,750 m (5,741.5 ft). This model is corroborated by Roth and Schmidt (2010). Therefore, as the ship travels through the ice, a watch 3,500 m (11,482.9 ft) wide would be subject to sound levels greater than or equal to 120 dB. This results in potential exposure of 3,500 km² (1,020.4 nmi²) to sounds greater than or equal to 120 dB from icebreaking.

Data characterizing the sound levels generated by icebreaking activities

conducted by the *Palmer* are not available; therefore, data for noise generating from an icebreaking vessel such as the U.S. Coast Guard Cutter (USCGC) *Healy* will be used as a proxy. It is noted that the *Palmer* is a smaller vessel and has less icebreaking capability than the U.S. Coast Guard's other polar icebreakers, being only capable of breaking ice up to 1 m thick at speeds of 3 kts (5.6 km/hr or 3 nmi). Therefore, the sound levels that may be generated by the *Palmer* are expected to be lower than the conservative levels estimated and measured for the *Healy*. Researchers will work to minimize time spent breaking ice as science operations are more difficult to conduct in icy conditions since the ice noise degrades the quality of the seismic and ADCP data and time spent breaking ice takes away from time supporting scientific research. Logistically, if the vessel were in heavy ice conditions, researchers would not tow the airgun array and streamer, as this would likely damage equipment and generate noisy data. It is possible that the seismic survey can be performed in low ice conditions if the *Palmer* could generate an open path behind the vessel.

Because the *Palmer* is not rated to break multi-year ice routinely, operations generally avoid transiting through older ice (i.e., 2 years or older, thicker than 1 m). If sea ice is encountered during the cruise, it is anticipated the *Palmer* will proceed primarily through one year sea ice, and possibly some new, very thin ice, and would follow leads wherever possible. Satellite imagery from the Totten region documents that sea ice is at its minimum extent during the month of February. The most recent image for the region, from November 21, 2013, shows that the sea ice is currently breaking up, with a significant coastal lead of open water. Based on a maximum sea ice extent of 250 km (135 nmi) and estimating that NSF and ASC will transit to the innermost shelf and back into open water twice, a round trip transit in each of the potential work regions, NSF and ASC estimate that the *Palmer* will actively break ice up to a distance of 1,000 km (540 nmi). Based on a ship's speed of 5 kts under moderate ice conditions, this distance represents approximately 108 hrs of icebreaking operations. It is noted that typical transit through areas primarily open water and containing brash ice or pancake ice will not be considered icebreaking.

Dates, Duration, and Specified Geographic Region

The proposed project and survey sites are located in selected regions of the Dumont d'Urville Sea in the Southern Ocean off the coast of East Antarctica and focus on the Totten Glacier and Moscow University Ice Shelf, located on the Sabrina Coast, from greater than approximately 64° South and between approximately 95 to 135° East (see Figure 2 of the IHA application), and the Mertz Glacier and Cook Ice Shelf systems located on the George V and Oates Coast, from greater than approximately 65° South and between approximately 140 to 165° East in International Waters. The proposed study sites are characterized by heavy ice cover, with a seasonal break-up in the ice that structures biological patterns. The proposed studies would occur in both areas, or entirely in one or the other, depending on ice conditions. Figure 3 of the IHA application illustrates the limited detailed bathymetry of the two study areas. Ice conditions encountered during the previous surveys in the region limited the area where bathymetric data could be collected. Water depths in the survey area range from approximately 100 to 1,000 m, and possibly exceeding 1,000 m in some areas. There is limited information on the depths in the study area and therefore more detailed information on bathymetry is not available. Figures 2 and 3 of the IHA application illustrate the limited available detailed bathymetry of the two proposed study areas due to ice conditions encountered during previous surveys in the region. The proposed seismic survey would be within an area of approximately 5,628 km² (1,640.9 nmi²). This estimate is based on the maximum number of kilometers for the seismic survey (2,800 km) times the predicted rms radii (m) based on modeling and empirical measurements (assuming 100% use of the two 105 in³ GI airguns in 100 to 1,000 m water depths) which was calculated to be 1,005 m (3,297.2 ft).

The icebreaking will occur, as necessary, between approximately 66 to 70° South and between 140 to 165° East. The total distance in the region of the vessel will travel include the proposed seismic survey and transit to dredging or sampling locations and will represent approximately 5,600 km (3,023.8 nmi). Based on a maximum sea ice extent of 250 km (135 nmi) and estimating that NSF and ASC will transit to the innermost shelf and back into open water twice, a round trip transit in each of the potential work regions, NSF and

ASC estimate that the Palmer will actively break ice up to a distance of 1,000 km (540 nmi). Based on a ship's speed of 5 kts under moderate ice conditions, this distance represents approximately 108 hrs of icebreaking operations.

The *Palmer* is expected to depart from Hobart, Tasmania on approximately January 29, 2014 and arrive at Hobart, Tasmania on approximately March 16, 2014. Research operations would be over a span of 45-days, including to and from port. Ice-free or very low concentrations of sea ice are required in order to collect high quality seismic data and not impede passage of the vessel between sampling locations. This requirement restricts the cruise to operating in mid to late austral summer when the ice concentrations are typically the lowest. Some minor deviation from this schedule is possible, depending on logistics and weather (i.e., the cruise may depart earlier or be extended due to poor weather; there could be additional days of seismic operations if collected data are deemed to be of substandard quality).

Description of the Marine Mammals in the Area of the Proposed Specified Activity

The marine mammals that generally occur in the proposed action area belong to three taxonomic groups: Mysticetes (baleen whales), odontocetes (toothed whales), and pinnipeds (seals and sea lions). The marine mammal species that potentially occur within the Southern Ocean in proximity to the proposed action area in the Dumont d'Urville Sea include 28 species of cetaceans and 6 species of pinnipeds.

The Dumont d'Urville Sea may be a feeding ground for many of these marine mammals. Many of the species that may be potentially present in the study area seasonally migrate to higher latitudes along the east coast of Antarctica. In general, most species (except for the killer whale) migrate north in the middle of the austral winter and return to Antarctica in the early austral summer. Some species, particularly Antarctic minke (*Balaenoptera bonaerensis*) and killer whales (*Orcinus orca*), are expected to be present in higher concentrations along the ice edge (SCAR, 2002). The 6 species of pinnipeds that are found in the Southern Ocean and which may be present in the proposed study area include the crabeater (*Lebodon carcinophagus*), leopard (*Hydrurga leptonyx*), Wedell (*Leptonychotes weddellii*), Ross (*Ommatophoca rossii*), southern elephant (*Mirounga leonina*), and Antarctic fur seal (*Arctocephalus*

gazella). Many of these pinniped species breed on either the pack ice or sub-Antarctic islands. Since the southern elephant seal and Antarctic fur seal haul-outs and rookeries are located on sub-Antarctic islands and prefer beaches, they are more common north of the seasonally shifting pack ice found in the proposed study area; therefore, these two species have not been considered further. Marine mammal species listed as endangered under the U.S. Endangered Species Act of 1973 (ESA; 16 U.S.C. 1531 *et seq.*), includes the southern right (*Eubalaena australis*), humpback (*Megaptera novaeangliae*), sei (*Balaenoptera borealis*), fin (*Balaenoptera physalus*), blue (*Balaenoptera musculus*), and sperm (*Physeter macrocephalus*) whale. Of those endangered species, the humpback, sei, fin, blue, and sperm whale are likely to be encountered in the proposed survey area.

Various national Antarctic research programs along the coast of East Antarctica have conducted scientific cruises that included data on marine mammal sightings. These observations were made primarily between 30° East and 170° East and north to 60° South. The reported cetacean sightings are summarized in Tables 5 to 7 of the IHA application. For pinnipeds, observations made during a scientific cruise over a 13-day period in East Antarctica are summarized in Table 8 of the IHA application. These observations were made below 60° South and between 110° East to 165° East and include sightings of individual animals in the water as well as individuals that were hauled-out (i.e., resting on the surface of the sea ice).

Records from the International Whaling Commission's Southern Ocean Whale and Ecosystem Research (IWC-SOWER) circumpolar cruises were also considered. In addition to the 14 species known to occur in the Dumont d'Urville Sea of the Southern Ocean, there are 18 cetacean species with ranges that are known to occur in the sub-Antarctic waters of the study area which may also feed and/or migrate to the Southern Ocean during the austral summer, these include the southern right, pygmy right (*Caperea marginata*), Bryde's (*Balaenoptera brydeii*), dwarf minke (*Balaenoptera acutorostrata* spp.), pygmy blue (*Balaenoptera musculus breviceps*), pygmy dwarf sperm whale (*Kogia breviceps*), Arnoux's beaked (*Berardius arnuxii*), Blainville's beaked whale (*Mesoplodon densirostris*), Cuvier's beaked (*Ziphiopsis cavirostris*), Shepherd's beaked (*Tasmacetus shepherdii*), Southern bottlenose (*Hyperoodon planifrons*), Andrew's

beaked (*Mesoplodon bowdoini*), Hector's beaked (*Mesoplodon hectori*), Gray's beaked (*Mesoplodon grayi*), strap-toothed beaked (*Mesoplodon layardii*), spade-toothed beaked (*Mesoplodon traversii*), southern right whale dolphin (*Lissodelphis peronii*),

Dusky (*Lagenorhynchus obscurus*), and bottlenose dolphin (*Tursiops truncatus*). However, these species have not been sighted and are not expected to occur where the proposed activities would take place. These species are not considered further in this document.

Table 4 (below) presents information on the abundance, distribution, population status, conservation status, and population trend of the species of marine mammals that may occur in the proposed study area during February to March 2014.

TABLE 4—THE HABITAT, REGIONAL ABUNDANCE, AND CONSERVATION STATUS OF MARINE MAMMALS THAT MAY OCCUR IN OR NEAR THE PROPOSED LOW-ENERGY SEISMIC SURVEY AREA IN THE ANTARCTIC AREA OF THE SOUTHERN OCEAN
[See text and Tables 4 in NSF and ASC's application for further details]

Species	Habitat	Population estimate	ESA ¹	MMPA ²	Population trend
Mysticetes:					
Southern right whale (<i>Eubalaena australis</i>).	Coastal, pelagic	8,000 ³ to 15,000 ⁴	EN	D	Increasing.
Pygmy right whale (<i>Caperea marginata</i>).	Coastal, pelagic	NA	NL	NC	NA.
Humpback whale (<i>Megaptera novaeangliae</i>).	Pelagic, nearshore waters, and banks.	35,000 to 40,000 ³ —Worldwide. 9,484 ⁵ —Scotia Sea and Antarctica Peninsula.	EN	D	Increasing.
Dwarf minke whale (<i>Balaenoptera acutorostrata</i> sub-species).	Pelagic and coastal	NA	NL	NC	NA.
Antarctic minke whale (<i>Balaenoptera bonaerensis</i>).	Pelagic, ice floes	Several 100,000 ³ —Worldwide. 18,125 ⁵ —Scotia Sea and Antarctica Peninsula.	NL	NC	Stable.
Bryde's whale (<i>Balaenoptera brydei</i>).	Pelagic and coastal	NA	NL	NC	NA.
Sei whale (<i>Balaenoptera borealis</i>).	Primarily offshore, pelagic ...	80,000 ³ —Worldwide	EN	D	NA.
Fin whale (<i>Balaenoptera physalus</i>).	Continental slope, pelagic ...	140,000 ³ —Worldwide	EN	D	NA.
Blue whale (<i>Balaenoptera musculus</i>).	Pelagic, shelf, coastal	8,000 to 9,000 ³ —Worldwide 1,700 ⁶ —Southern Ocean ...	EN	D	NA.
Odontocetes:					
Sperm whale (<i>Physeter macrocephalus</i>).	Pelagic, deep sea	360,000 ³ —Worldwide	EN	D	NA.
Pygmy sperm whale (Kogia breviceps).	Pelagic, slope	9,500 ³ —Antarctic	NL	NC	NA.
Arnoux's beaked whale (<i>Berardius arnuxii</i>).	Pelagic	NA	NL	NC	NA.
Blainville's beaked whale (<i>Mesoplodon densirostris</i>).	Pelagic	NA	NL	NC	NA.
Cuvier's beaked whale (<i>Ziphius cavirostris</i>).	Pelagic	NA	NL	NC	NA.
Shepherd's beaked whale (Tasmacetus shepherdi).	Pelagic	NA	NL	NC	NA.
Southern bottlenose whale (<i>Hyperoodon planifrons</i>).	Pelagic	500,000 ³ —South of Antarctic Convergence.	NL	NC	NA.
Andrew's beaked whale (<i>Mesoplodon bowdoini</i>).	Pelagic	NA	NL	NC	NA.
Hector's beaked whale (<i>Mesoplodon hectori</i>).	Pelagic	NA	NL	NC	NA.
Gray's beaked whale (<i>Mesoplodon grayi</i>).	Pelagic	NA	NL	NC	NA.
Strap-toothed beaked whale (<i>Mesoplodon layardii</i>).	Pelagic	NA	NL	NC	NA.
Spade-toothed beaked whale (<i>Mesoplodon traversii</i>).	Pelagic	NA	NL	NC	NA.

TABLE 4—THE HABITAT, REGIONAL ABUNDANCE, AND CONSERVATION STATUS OF MARINE MAMMALS THAT MAY OCCUR IN OR NEAR THE PROPOSED LOW-ENERGY SEISMIC SURVEY AREA IN THE ANTARCTIC AREA OF THE SOUTHERN OCEAN—Continued

[See text and Tables 4 in NSF and ASC's application for further details]

Species	Habitat	Population estimate	ESA ¹	MMPA ²	Population trend
Killer whale (<i>Orcinus orca</i>).	Pelagic, shelf, coastal, pack ice.	80,000 ³ —South of Antarctic Convergence.	NL	NC	NA.
Long-finned pilot whale (<i>Globicephala melas</i>).	Pelagic, shelf, coastal	25,000 ⁷ —Southern Ocean ..	NL	NC	NA.
Bottlenose dolphin (<i>Tursiops truncatus</i>).	Offshore, inshore, coastal, estuaries.	200,000 ^{3,8} —South of Antarctic Convergence.	NL	NC	NA.
Southern right whale dolphin (<i>Lissodelphis peronii</i>).	Pelagic	>625,500 ³ —Worldwide	NL	NC	NA.
Dusky dolphin (<i>Lagenorhynchus obscurus</i>).	Coastal, continental shelf and slope.	NA	NL	NC	NA.
Hourglass dolphin (<i>Lagenorhynchus cruciger</i>).	Pelagic, ice edge	144,000 ³	NL	NC	NA.
Spectacled porpoise (<i>Phocoena dioptrica</i>).	Coastal, pelagic	NA	NL	NC	NA.
Pinnipeds:					
Crabeater seal (<i>Lobodon carcinophaga</i>).	Coastal, pack ice	5,000,000 to 15,000,000 ^{3,9} ..	NL	NC	Increasing.
Leopard seal (<i>Hydrurga leptonyx</i>).	Pack ice, sub-Antarctic islands.	220,000 to 440,000 ^{3,10}	NL	NC	NA.
Ross seal (<i>Ommatophoca rossii</i>).	Pack ice, smooth ice floes, pelagic.	130,000 ³	NL	NC	NA.
Wedell seal (<i>Leptonychotes weddellii</i>).	Fast ice, pack ice, sub-Antarctic islands.	500,000 to 1,000,000 ^{3,11}	NL	NC	NA.
Southern elephant seal (<i>Mirounga leonina</i>).	Coastal, pelagic, sub-Antarctic waters.	640,000 ¹² to 650,000 ³	NL	NC	Decreasing, increasing or stable depending on breeding population.
Antarctic fur seal (<i>Arctocephalus gazella</i>).	Shelf, rocky habitats	1,600,000 ¹³ to 3,000,000 ³ ..	NL	NC	Increasing.

NA = Not available or not assessed.

¹ U.S. Endangered Species Act: EN = Endangered, T = Threatened, DL = Delisted, NL = Not listed.² U.S. Marine Mammal Protection Act: D = Depleted, S = Strategic, NC = Not Classified.³ Jefferson *et al.*, 2008.⁴ Kenney, 2009.⁵ Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR) survey area (Reilly *et al.*, 2004).⁶ Sears and Perrin, 2009.⁷ Ford, 2009.⁸ Olson, 2009.⁹ Bengston, 2009.¹⁰ Rogers, 2009.¹¹ Thomas and Terhune, 2009.¹² Hindell and Perrin, 2009.¹³ Arnould, 2009.

Refer to sections 3 and 4 of NSF and ASC's IHA application for detailed information regarding the abundance and distribution, population status, and life history and behavior of these other marine mammal species and their occurrence in the proposed project area. The IHA application also presents how NSF and ASC calculated the estimated densities for the marine mammals in the proposed survey area. NMFS has reviewed these data and determined them to be the best available scientific information for the purposes of the proposed IHA.

Potential Effects on Marine Mammals

Acoustic stimuli generated by the operation of the airguns, which introduce sound into the marine environment, may have the potential to cause Level B harassment of marine mammals in the proposed survey area. The effects of sounds from airgun operations might include one or more of the following: Tolerance, masking of natural sounds, behavioral disturbance, temporary or permanent hearing impairment, or non-auditory physical or physiological effects (Richardson *et al.*, 1995; Gordon *et al.*, 2004; Nowacek *et al.*, 2007; Southall *et al.*, 2007).

Permanent hearing impairment, in the unlikely event that it occurred, would constitute injury, but temporary threshold shift (TTS) is not an injury (Southall *et al.*, 2007). Although the possibility cannot be entirely excluded, it is unlikely that the proposed project would result in any cases of temporary or permanent hearing impairment, or any significant non-auditory physical or physiological effects. Based on the available data and studies described here, some behavioral disturbance is expected. A more comprehensive review of these issues can be found in the "Programmatic Environmental

Impact Statement/Overseas Environmental Impact Statement prepared for Marine Seismic Research that is funded by the National Science Foundation and conducted by the U.S. Geological Survey" (NSF/USGS, 2011).

Tolerance

Richardson *et al.* (1995) defines tolerance as the occurrence of marine mammals in areas where they are exposed to human activities or man-made noise. In many cases, tolerance develops by the animal habituating to the stimulus (i.e., the gradual waning of responses to a repeated or ongoing stimulus) (Richardson, *et al.*, 1995; Thorpe, 1963), but because of ecological or physiological requirements, many marine animals may need to remain in areas where they are exposed to chronic stimuli (Richardson, *et al.*, 1995).

Numerous studies have shown that pulsed sounds from airguns are often readily detectable in the water at distances of many kilometers. Several studies have shown that marine mammals at distances more than a few kilometers from operating seismic vessels often show no apparent response. That is often true even in cases when the pulsed sounds must be readily audible to the animals based on measured received levels and the hearing sensitivity of the marine mammal group. Although various baleen whales and toothed whales, and (less frequently) pinnipeds have been shown to react behaviorally to airgun pulses under some conditions, at other times marine mammals of all three types have shown no overt reactions. The relative responsiveness of baleen and toothed whales are quite variable.

Masking

The term masking refers to the inability of a subject to recognize the occurrence of an acoustic stimulus as a result of the interference of another acoustic stimulus (Clark *et al.*, 2009). Introduced underwater sound may, through masking, reduce the effective communication distance of a marine mammal species if the frequency of the source is close to that used as a signal by the marine mammal, and if the anthropogenic sound is present for a significant fraction of the time (Richardson *et al.*, 1995).

The airguns for the proposed low-energy seismic survey have dominant frequency components of 2 to 188 Hz. This frequency range fully overlaps the lower part of the frequency range of odontocete calls and/or functional hearing (full range about 150 Hz to 180 kHz). Airguns also produce a small portion of their sound at mid and high

frequencies that overlap most, if not all, frequencies produced by odontocetes. While it is assumed that mysticetes can detect acoustic impulses from airguns and vessel sounds (Richardson *et al.*, 1995a), sub-bottom profilers, pingers, and most of the multi-beam echosounders would likely be detectable by some mysticetes based on presumed mysticete hearing sensitivity. Odontocetes are presumably more sensitive to mid to high frequencies produced by the multi-beam echosounders, sub-bottom profilers, and pingers than to the dominant low frequencies produced by the airguns and vessel. A more comprehensive review of the relevant background information for odontocetes appears in Section 3.6.4.3, Section 3.7.4.3 and Appendix E of the NSF/USGS PEIS (2011).

Masking effects of pulsed sounds (even from large arrays of airguns) on marine mammal calls and other natural sounds are expected to be limited. Because of the intermittent nature and low duty cycle of seismic airgun pulses, animals can emit and receive sounds in the relatively quiet intervals between pulses. However, in some situations, reverberation occurs for much or the entire interval between pulses (e.g., Simard *et al.*, 2005; Clark and Gagnon, 2006) which could mask calls. Some baleen and toothed whales are known to continue calling in the presence of seismic pulses, and their calls can usually be heard between the seismic pulses (e.g., Richardson *et al.*, 1986; McDonald *et al.*, 1995; Greene *et al.*, 1999; Nieuwirth *et al.*, 2004; Smultea *et al.*, 2004; Holst *et al.*, 2005a,b, 2006; and Dunn and Hernandez, 2009). However, Clark and Gagnon (2006) reported that fin whales in the North Atlantic Ocean went silent for an extended period starting soon after the onset of a seismic survey in the area. Similarly, there has been one report that sperm whales ceased calling when exposed to pulses from a very distant seismic ship (Bowles *et al.*, 1994). However, more recent studies found that they continued calling in the presence of seismic pulses (Madsen *et al.*, 2002; Tyack *et al.*, 2003; Smultea *et al.*, 2004; Holst *et al.*, 2006; and Jochens *et al.*, 2008). Dilorio and Clark (2009) found evidence of increased calling by blue whales during operations by a lower-energy seismic source (i.e., sparker). Dolphins and porpoises commonly are heard calling while airguns are operating (e.g., Gordon *et al.*, 2004; Smultea *et al.*, 2004; Holst *et al.*, 2005a, b; and Potter *et al.*, 2007). The sounds important to small odontocetes are predominantly at much

higher frequencies than are the dominant components of airgun sounds, thus limiting the potential for masking.

Pinnipeds have the most sensitive hearing and/or produce most of their sounds in frequencies higher than the dominant components of airgun sound, but there is some overlap in the frequencies of the airgun pulses and the calls. However, the intermittent nature of airgun pulses presumably reduces the potential for masking.

Marine mammals are thought to be able to compensate for masking by adjusting their acoustic behavior through shifting call frequencies, increasing call volume, and increasing vocalization rates. For example blue whales are found to increase call rates when exposed to noise from seismic surveys in the St. Lawrence Estuary (Dilorio and Clark, 2009). The North Atlantic right whales (*Eubalaena glacialis*) exposed to high shipping noise increased call frequency (Parks *et al.*, 2007), while some humpback whales respond to low-frequency active sonar playbacks by increasing song length (Miller *et al.*, 2000). In general, NMFS expects the masking effects of seismic pulses to be minor, given the normally intermittent nature of seismic pulses.

Behavioral Disturbance

Marine mammals may behaviorally react to sound when exposed to anthropogenic noise. Disturbance includes a variety of effects, including subtle to conspicuous changes in behavior, movement, and displacement. Reactions to sound, if any, depend on species, state of maturity, experience, current activity, reproductive state, time of day, and many other factors (Richardson *et al.*, 1995; Wartzok *et al.*, 2004; Southall *et al.*, 2007; Weilgart, 2007). These behavioral reactions are often shown as: Changing durations of surfacing and dives, number of blows per surfacing, or moving direction and/or speed; reduced/increased vocal activities; changing/cessation of certain behavioral activities (such as socializing or feeding); visible startle response or aggressive behavior (such as tail/fluke slapping or jaw clapping); avoidance of areas where noise sources are located; and/or flight responses (e.g., pinnipeds flushing into the water from haul-outs or rookeries). If a marine mammal does react briefly to an underwater sound by changing its behavior or moving a small distance, the impacts of the change are unlikely to be significant to the individual, let alone the stock or population. However, if a sound source displaces marine mammals from an important feeding or breeding area for a

prolonged period, impacts on individuals and populations could be significant (e.g., Lusseau and Bejder, 2007; Weilgart, 2007).

The biological significance of many of these behavioral disturbances is difficult to predict, especially if the detected disturbances appear minor. However, the consequences of behavioral modification could be expected to be biologically significant if the change affects growth, survival, and/or reproduction. Some of these significant behavioral modifications include:

- Change in diving/surfacing patterns (such as those thought to be causing beaked whale stranding due to exposure to military mid-frequency tactical sonar);
- Habitat abandonment due to loss of desirable acoustic environment; and
- Cessation of feeding or social interaction.

The onset of behavioral disturbance from anthropogenic noise depends on both external factors (characteristics of noise sources and their paths) and the receiving animals (hearing, motivation, experience, demography) and is also difficult to predict (Richardson *et al.*, 1995; Southall *et al.*, 2007). Given the many uncertainties in predicting the quantity and types of impacts of noise on marine mammals, it is common practice to estimate how many mammals would be present within a particular distance of industrial activities and/or exposed to a particular level of sound. In most cases, this approach likely overestimates the numbers of marine mammals that would be affected in some biologically-important manner.

Baleen Whales—Baleen whales generally tend to avoid operating airguns, but avoidance radii are quite variable (reviewed in Richardson *et al.*, 1995; Gordon *et al.*, 2004). Whales are often reported to show no overt reactions to pulses from large arrays of airguns at distances beyond a few kilometers, even though the airgun pulses remain well above ambient noise levels out to much longer distances. However, baleen whales exposed to strong noise pulses from airguns often react by deviating from their normal migration route and/or interrupting their feeding and moving away. In the cases of migrating gray (*Eschrichtius robustus*) and bowhead (*Balaena mysticetus*) whales, the observed changes in behavior appeared to be of little or no biological consequence to the animals (Richardson *et al.*, 1995). They simply avoided the sound source by displacing their migration route to varying degrees, but within the natural boundaries of the migration corridors.

Studies of gray, bowhead, and humpback whales have shown that seismic pulses with received levels of 160 to 170 dB re 1 μ Pa (rms) seem to cause obvious avoidance behavior in a substantial fraction of the animals exposed (Malme *et al.*, 1986, 1988; Richardson *et al.*, 1995). In many areas, seismic pulses from large arrays of airguns diminish to those levels at distances ranging from 4 to 15 km (2.2 to 8.1 nmi) from the source. A substantial proportion of the baleen whales within those distances may show avoidance or other strong behavioral reactions to the airgun array. Subtle behavioral changes sometimes become evident at somewhat lower received levels, and studies have shown that some species of baleen whales, notably bowhead, gray, and humpback whales, at times, show strong avoidance at received levels lower than 160 to 170 dB re 1 μ Pa (rms).

Researchers have studied the responses of humpback whales to seismic surveys during migration, feeding during the summer months, breeding while offshore from Angola, and wintering offshore from Brazil. McCauley *et al.* (1998, 2000a) studied the responses of humpback whales off western Australia to a full-scale seismic survey with a 16 airgun array (2,678 in³) and to a single airgun (20 in³) with source level of 227 dB re 1 μ Pa (p-p). In the 1998 study, they documented that avoidance reactions began at 5 to 8 km (2.7 to 4.3 nmi) from the array, and that those reactions kept most pods approximately 3 to 4 km (1.6 to 2.2 nmi) from the operating seismic boat. In the 2000 study, they noted localized displacement during migration of 4 to 5 km (2.2 to 2.7 nmi) by traveling pods and 7 to 12 km (3.8 to 6.5 nmi) by more sensitive resting pods of cow-calf pairs. Avoidance distances with respect to the single airgun were smaller but consistent with the results from the full array in terms of the received sound levels. The mean received level for initial avoidance of an approaching airgun was 140 dB re 1 μ Pa (rms) for humpback pods containing females, and at the mean closest point of approach distance the received level was 143 dB re 1 μ Pa (rms). The initial avoidance response generally occurred at distances of 5 to 8 km (2.7 to 4.3 nmi) from the airgun array and 2 km (1.1 nmi) from the single airgun. However, some individual humpback whales, especially males, approached within distances of 100 to 400 m (328 to 1,312 ft), where the maximum received level was 179 dB re 1 μ Pa (rms).

Data collected by observers during several seismic surveys in the

Northwest Atlantic showed that sighting rates of humpback whales were significantly greater during non-seismic periods compared with periods when a full array was operating (Moulton and Holst, 2010). In addition, humpback whales were more likely to swim away and less likely to swim towards a vessel during seismic vs. non-seismic periods (Moulton and Holst, 2010).

Humpback whales on their summer feeding grounds in southeast Alaska did not exhibit persistent avoidance when exposed to seismic pulses from a 1.64–L (100 in³) airgun (Malme *et al.*, 1985). Some humpbacks seemed “startled” at received levels of 150 to 169 dB re 1 μ Pa. Malme *et al.* (1985) concluded that there was no clear evidence of avoidance, despite the possibility of subtle effects, at received levels up to 172 dB re 1 μ Pa (rms). However, Moulton and Holst (2010) reported that humpback whales monitored during seismic surveys in the Northwest Atlantic had lower sighting rates and were most often seen swimming away from the vessel during seismic periods compared with periods when airguns were silent.

Studies have suggested that South Atlantic humpback whales wintering off Brazil may be displaced or even strand upon exposure to seismic surveys (Engel *et al.*, 2004). The evidence for this was circumstantial and subject to alternative explanations (IAGC, 2004). Also, the evidence was not consistent with subsequent results from the same area of Brazil (Parente *et al.*, 2006), or with direct studies of humpbacks exposed to seismic surveys in other areas and seasons. After allowance for data from subsequent years, there was “no observable direct correlation” between strandings and seismic surveys (IWC, 2007: 236).

Reactions of migrating and feeding (but not wintering) gray whales to seismic surveys have been studied. Malme *et al.* (1986, 1988) studied the responses of feeding eastern Pacific gray whales to pulses from a single 100 in³ airgun off St. Lawrence Island in the northern Bering Sea. They estimated, based on small sample sizes, that 50 percent of feeding gray whales stopped feeding at an average received pressure level of 173 dB re 1 μ Pa on an (approximate) rms basis, and that 10 percent of feeding whales interrupted feeding at received levels of 163 dB re 1 μ Pa (rms). Those findings were generally consistent with the results of experiments conducted on larger numbers of gray whales that were migrating along the California coast (Malme *et al.*, 1984; Malme and Miles, 1985), and western Pacific gray whales

feeding off Sakhalin Island, Russia (Wursig *et al.*, 1999; Gailey *et al.*, 2007; Johnson *et al.*, 2007; Yazvenko *et al.*, 2007a, b), along with data on gray whales off British Columbia (Bain and Williams, 2006).

Various species of *Balaenoptera* (blue, sei, fin, and minke whales) have occasionally been seen in areas ensounded by airgun pulses (Stone, 2003; MacLean and Haley, 2004; Stone and Tasker, 2006), and calls from blue and fin whales have been localized in areas with airgun operations (e.g., McDonald *et al.*, 1995; Dunn and Hernandez, 2009; Castellote *et al.*, 2010). Sightings by observers on seismic vessels off the United Kingdom from 1997 to 2000 suggest that, during times of good sightability, sighting rates for mysticetes (mainly fin and sei whales) were similar when large arrays of airguns were shooting versus silent (Stone, 2003; Stone and Tasker, 2006). However, these whales tended to exhibit localized avoidance, remaining significantly further (on average) from the airgun array during seismic operations compared with non-seismic periods (Stone and Tasker, 2006). Castellote *et al.* (2010) reported that singing fin whales in the Mediterranean moved away from an operating airgun array.

Ship-based monitoring studies of baleen whales (including blue, fin, sei, minke, and humpback whales) in the Northwest Atlantic found that overall, this group had lower sighting rates during seismic vs. non-seismic periods (Moulton and Holst, 2010). Baleen whales as a group were also seen significantly farther from the vessel during seismic compared with non-seismic periods, and they were more often seen to be swimming away from the operating seismic vessel (Moulton and Holst, 2010). Blue and minke whales were initially sighted significantly farther from the vessel during seismic operations compared to non-seismic periods; the same trend was observed for fin whales (Moulton and Holst, 2010). Minke whales were most often observed to be swimming away from the vessel when seismic operations were underway (Moulton and Holst, 2010).

Data on short-term reactions by cetaceans to impulsive noises are not necessarily indicative of long-term or biologically significant effects. It is not known whether impulsive sounds affect reproductive rate or distribution and habitat use in subsequent days or years. However, gray whales have continued to migrate annually along the west coast of North America with substantial increases in the population over recent

years, despite intermittent seismic exploration (and much ship traffic) in that area for decades (Appendix A in Malme *et al.*, 1984; Richardson *et al.*, 1995; Allen and Angliss, 2010). The western Pacific gray whale population did not seem affected by a seismic survey in its feeding ground during a previous year (Johnson *et al.*, 2007). Similarly, bowhead whales have continued to travel to the eastern Beaufort Sea each summer, and their numbers have increased notably, despite seismic exploration in their summer and autumn range for many years (Richardson *et al.*, 1987; Allen and Angliss, 2010). The history of coexistence between seismic surveys and baleen whales suggests that brief exposures to sound pulses from any single seismic survey are unlikely to result in prolonged effects.

Toothed Whales—Little systematic information is available about reactions of toothed whales to noise pulses. Few studies similar to the more extensive baleen whale/seismic pulse work summarized above have been reported for toothed whales. However, there are recent systematic studies on sperm whales (e.g., Gordon *et al.*, 2006; Madsen *et al.*, 2006; Winsor and Mate, 2006; Jochens *et al.*, 2008; Miller *et al.*, 2009). There is an increasing amount of information about responses of various odontocetes to seismic surveys based on monitoring studies (e.g., Stone, 2003; Smultea *et al.*, 2004; Moulton and Miller, 2005; Bain and Williams, 2006; Holst *et al.*, 2006; Stone and Tasker, 2006; Potter *et al.*, 2007; Hauser *et al.*, 2008; Holst and Smultea, 2008; Weir, 2008; Barkaszi *et al.*, 2009; Richardson *et al.*, 2009; Moulton and Holst, 2010).

Seismic operators and PSOs on seismic vessels regularly see dolphins and other small toothed whales near operating airgun arrays, but in general there is a tendency for most delphinids to show some avoidance of operating seismic vessels (e.g., Goold, 1996a,b,c; Calambokidis and Osmeck, 1998; Stone, 2003; Moulton and Miller, 2005; Holst *et al.*, 2006; Stone and Tasker, 2006; Weir, 2008; Richardson *et al.*, 2009; Barkaszi *et al.*, 2009; Moulton and Holst, 2010). Some dolphins seem to be attracted to the seismic vessel and floats, and some ride the bow wave of the seismic vessel even when large arrays of airguns are firing (e.g., Moulton and Miller, 2005). Nonetheless, small toothed whales more often tend to head away, or to maintain a somewhat greater distance from the vessel, when a large array of airguns is operating than when it is silent (e.g., Stone and Tasker, 2006; Weir, 2008; Barry *et al.*, 2010; Moulton and Holst, 2010). In most

cases, the avoidance radii for delphinids appear to be small, on the order of one km or less, and some individuals show no apparent avoidance. Captive bottlenose dolphins and beluga whales (*Delphinapterus leucas*) exhibited changes in behavior when exposed to strong pulsed sounds similar in duration to those typically used in seismic surveys (Finneran *et al.*, 2000, 2002, 2005). However, the animals tolerated high received levels of sound before exhibiting aversive behaviors.

Results of porpoises depend on species. The limited available data suggest that harbor porpoises (*Phocoena phocoena*) show stronger avoidance of seismic operations than do Dall's porpoises (*Phocoenoides dalli*) (Stone, 2003; MacLean and Koski, 2005; Bain and Williams, 2006; Stone and Tasker, 2006). Dall's porpoises seem relatively tolerant of airgun operations (MacLean and Koski, 2005; Bain and Williams, 2006), although they too have been observed to avoid large arrays of operating airguns (Calambokidis and Osmeck, 1998; Bain and Williams, 2006). This apparent difference in responsiveness of these two porpoise species is consistent with their relative responsiveness to boat traffic and some other acoustic sources (Richardson *et al.*, 1995; Southall *et al.*, 2007).

Most studies of sperm whales exposed to airgun sounds indicate that the sperm whale shows considerable tolerance of airgun pulses (e.g., Stone, 2003; Moulton *et al.*, 2005, 2006a; Stone and Tasker, 2006; Weir, 2008). In most cases the whales do not show strong avoidance, and they continue to call. However, controlled exposure experiments in the Gulf of Mexico indicate that foraging behavior was altered upon exposure to airgun sound (Jochens *et al.*, 2008; Miller *et al.*, 2009; Tyack, 2009). There are almost no specific data on the behavioral reactions of beaked whales to seismic surveys. However, some northern bottlenose whales (*Hyperoodon ampullatus*) remained in the general area and continued to produce high-frequency clicks when exposed to sound pulses from distant seismic surveys (Gosselin and Lawson, 2004; Laurinolli and Cochrane, 2005; Simard *et al.*, 2005). Most beaked whales tend to avoid approaching vessels of other types (e.g., Wursig *et al.*, 1998). They may also dive for an extended period when approached by a vessel (e.g., Kasuya, 1986), although it is uncertain how much longer such dives may be as compared to dives by undisturbed beaked whales, which also are often quite long (Baird *et al.*, 2006; Tyack *et al.*, 2006). Based on a single observation,

Aguilar-Soto *et al.* (2006) suggested that foraging efficiency of Cuvier's beaked whales may be reduced by close approach of vessels. In any event, it is likely that most beaked whales would also show strong avoidance of an approaching seismic vessel, although this has not been documented explicitly. In fact, Moulton and Holst (2010) reported 15 sightings of beaked whales during seismic studies in the Northwest Atlantic; seven of those sightings were made at times when at least one airgun was operating. There was little evidence to indicate that beaked whale behavior was affected by airgun operations; sighting rates and distances were similar during seismic and non-seismic periods (Moulton and Holst, 2010).

There are increasing indications that some beaked whales tend to strand when naval exercises involving mid-frequency sonar operation are ongoing nearby (e.g., Simmonds and Lopez-Jurado, 1991; Frantzis, 1998; NOAA and USN, 2001; Jepson *et al.*, 2003; Hildebrand, 2005; Barlow and Gisiner, 2006; see also the "Stranding and Mortality" section in this notice). These strandings are apparently a disturbance response, although auditory or other injuries or other physiological effects may also be involved. Whether beaked whales would ever react similarly to seismic surveys is unknown. Seismic survey sounds are quite different from those of the sonar in operation during the above-cited incidents.

Odontocete reactions to large arrays of airguns are variable and, at least for delphinids, seem to be confined to a smaller radius than has been observed for the more responsive of some mysticetes. However, other data suggest that some odontocete species, including harbor porpoises, may be more responsive than might be expected given their poor low-frequency hearing. Reactions at longer distances may be particularly likely when sound propagation conditions are conducive to transmission of the higher frequency components of airgun sound to the animals' location (DeRuiter *et al.*, 2006; Goold and Coates, 2006; Tyack *et al.*, 2006; Potter *et al.*, 2007).

Pinnipeds—Pinnipeds are not likely to show a strong avoidance reaction to the airgun array. Visual monitoring from seismic vessels has shown only slight (if any) avoidance of airguns by pinnipeds, and only slight (if any) changes in behavior. In the Beaufort Sea, some ringed seals avoided an area of 100 m to (at most) a few hundred meters around seismic vessels, but many seals remained within 100 to 200 m (328 to 656 ft) of the trackline as the operating

airgun array passed by (e.g., Harris *et al.*, 2001; Moulton and Lawson, 2002; Miller *et al.*, 2005.). Ringed seal (*Pusa hispida*) sightings averaged somewhat farther away from the seismic vessel when the airguns were operating than when they were not, but the difference was small (Moulton and Lawson, 2002). Similarly, in Puget Sound, sighting distances for harbor seals (*Phoca vitulina*) and California sea lions (*Zalophus californianus*) tended to be larger when airguns were operating (Calambokidis and Osmeck, 1998). Previous telemetry work suggests that avoidance and other behavioral reactions may be stronger than evident to date from visual studies (Thompson *et al.*, 1998).

During seismic exploration off Nova Scotia, gray seals (*Halichoerus grypus*) exposed to noise from airguns and linear explosive charges did not react strongly (J. Parsons in Greene *et al.*, 1985). Pinnipeds in both water and air, sometimes tolerate strong noise pulses from non-explosive and explosive scaring devices, especially if attracted to the area for feeding and reproduction (Mate and Harvey, 1987; Reeves *et al.*, 1996). Thus pinnipeds are expected to be rather tolerant of, or habituate to, repeated underwater sounds from distant seismic sources, at least when the animals are strongly attracted to the area.

Hearing Impairment and Other Physical Effects

Exposure to high intensity sound for a sufficient duration may result in auditory effects such as a noise-induced threshold shift—an increase in the auditory threshold after exposure to noise (Finneran, Carder, Schlundt, and Ridgway, 2005). Factors that influence the amount of threshold shift include the amplitude, duration, frequency content, temporal pattern, and energy distribution of noise exposure. The magnitude of hearing threshold shift normally decreases over time following cessation of the noise exposure. The amount of threshold shift just after exposure is called the initial threshold shift. If the threshold shift eventually returns to zero (i.e., the threshold returns to the pre-exposure value), it is called temporary threshold shift (TTS) (Southall *et al.*, 2007). Researchers have studied TTS in certain captive odontocetes and pinnipeds exposed to strong sounds (reviewed in Southall *et al.*, 2007). However, there has been no specific documentation of TTS let alone permanent hearing damage, i.e., permanent threshold shift (PTS), in free-ranging marine mammals exposed to

sequences of airgun pulses during realistic field conditions.

Temporary Threshold Shift—TTS is the mildest form of hearing impairment that can occur during exposure to a strong sound (Kryter, 1985). While experiencing TTS, the hearing threshold rises and a sound must be stronger in order to be heard. At least in terrestrial mammals, TTS can last from minutes or hours to (in cases of strong TTS) days. For sound exposures at or somewhat above the TTS threshold, hearing sensitivity in both terrestrial and marine mammals recovers rapidly after exposure to the noise ends. Few data on sound levels and durations necessary to elicit mild TTS have been obtained for marine mammals, and none of the published data concern TTS elicited by exposure to multiple pulses of sound. Available data on TTS in marine mammals are summarized in Southall *et al.* (2007). Table 2 (above) presents the estimated distances from the Palmer's airguns at which the received energy level (per pulse, flat-weighted) would be expected to be greater than or equal to 180 and 190 dB re 1 μ Pa (rms).

To avoid the potential for injury, NMFS (1995, 2000) concluded that cetaceans and pinnipeds should not be exposed to pulsed underwater noise at received levels exceeding 180 and 190 dB re 1 μ Pa (rms). NMFS believes that to avoid the potential for Level A harassment, cetaceans and pinnipeds should not be exposed to pulsed underwater noise at received levels exceeding 180 and 190 dB re 1 μ Pa (rms), respectively. The established 180 and 190 dB (rms) criteria are not considered to be the levels above which TTS might occur. Rather, they are the received levels above which, in the view of a panel of bioacoustics specialists convened by NMFS before TTS measurements for marine mammals started to become available, one could not be certain that there would be no injurious effects, auditory or otherwise, to marine mammals. NMFS also assumes that cetaceans and pinnipeds exposed to levels exceeding 160 dB re 1 μ Pa (rms) may experience Level B harassment.

For toothed whales, researchers have derived TTS information from odontocetes from studies on the bottlenose dolphin and beluga. The experiments show that exposure to a single impulse at a received level of 207 kPa (or 30 psi, p-p), which is equivalent to 228 dB re 1 Pa (p-p), resulted in a 7 and 6 dB TTS in the beluga whale at 0.4 and 30 kHz, respectively. Thresholds returned to within 2 dB of the pre-exposure level within 4 minutes of the exposure (Finneran *et al.*, 2002). For the

one harbor porpoise tested, the received level of airgun sound that elicited onset of TTS was lower (Lucke *et al.*, 2009). If these results from a single animal are representative, it is inappropriate to assume that onset of TTS occurs at similar received levels in all odontocetes (*cf.* Southall *et al.*, 2007). Some cetaceans apparently can incur TTS at considerably lower sound exposures than are necessary to elicit TTS in the beluga or bottlenose dolphin.

For baleen whales, there are no data, direct or indirect, on levels or properties of sound that are required to induce TTS. The frequencies to which baleen whales are most sensitive are assumed to be lower than those to which odontocetes are most sensitive, and natural background noise levels at those low frequencies tend to be higher. As a result, auditory thresholds of baleen whales within their frequency band of best hearing are believed to be higher (less sensitive) than are those of odontocetes at their best frequencies (Clark and Ellison, 2004). From this, it is suspected that received levels causing TTS onset may also be higher in baleen whales than those of odontocetes (Southall *et al.*, 2007).

In pinnipeds, researchers have not measured TTS thresholds associated with exposure to brief pulses (single or multiple) of underwater sound. Initial evidence from more prolonged (non-pulse) exposures suggested that some pinnipeds (harbor seals in particular) incur TTS at somewhat lower received levels than do small odontocetes exposed for similar durations (Kastak *et al.*, 1999, 2005; Ketten *et al.*, 2001). The TTS threshold for pulsed sounds has been indirectly estimated as being an SEL of approximately 171 dB re 1 $\mu\text{Pa}^2\cdot\text{s}$ (Southall *et al.*, 2007) which would be equivalent to a single pulse with a received level of approximately 181 to 186 dB re 1 μPa (rms), or a series of pulses for which the highest rms values are a few dB lower. Corresponding values for California sea lions and northern elephant seals (*Mirounga angustirostris*) are likely to be higher (Kastak *et al.*, 2005).

Permanent Threshold Shift—When PTS occurs, there is physical damage to the sound receptors in the ear. In severe cases, there can be total or partial deafness, whereas in other cases, the animal has an impaired ability to hear sounds in specific frequency ranges (Kryter, 1985). There is no specific evidence that exposure to pulses of airgun sound can cause PTS in any marine mammal, even with large arrays of airguns. However, given the possibility that mammals close to an airgun array might incur at least mild

TTS, there has been further speculation about the possibility that some individuals occurring very close to airguns might incur PTS (e.g., Richardson *et al.*, 1995, p. 372ff; Gedamke *et al.*, 2008). Single or occasional occurrences of mild TTS are not indicative of permanent auditory damage, but repeated or (in some cases) single exposures to a level well above that causing TTS onset might elicit PTS.

Relationships between TTS and PTS thresholds have not been studied in marine mammals but are assumed to be similar to those in humans and other terrestrial mammals (Southall *et al.*, 2007). PTS might occur at a received sound level at least several dBs above that inducing mild TTS if the animal were exposed to strong sound pulses with rapid rise times. Based on data from terrestrial mammals, a precautionary assumption is that the PTS threshold for impulse sounds (such as airgun pulses as received close to the source) is at least 6 dB higher than the TTS threshold on a peak-pressure basis, and probably greater than 6 dB (Southall *et al.*, 2007).

Given the higher level of sound necessary to cause PTS as compared with TTS, it is considerably less likely that PTS would occur. Baleen whales generally avoid the immediate area around operating seismic vessels, as do some other marine mammals.

Stranding and Mortality—When a living or dead marine mammal swims or floats onto shore and becomes “beached” or incapable of returning to sea, the event is termed a “stranding” (Geraci *et al.*, 1999; Perrin and Geraci, 2002; Geraci and Lounsbury, 2005; NMFS, 2007). The legal definition for a stranding under the MMPA is that “(A) a marine mammal is dead and is (i) on a beach or shore of the United States; or (ii) in waters under the jurisdiction of the United States (including any navigable waters); or (B) a marine mammal is alive and is (i) on a beach or shore of the United States and is unable to return to the water; (ii) on a beach or shore of the United States and, although able to return to the water is in need of apparent medical attention; or (iii) in the waters under the jurisdiction of the United States (including any navigable waters), but is unable to return to its natural habitat under its own power or without assistance.”

Marine mammals are known to strand for a variety of reasons, such as infectious agents, biotoxins, starvation, fishery interaction, ship strike, unusual oceanographic or weather events, sound exposure, or combinations of these stressors

sustained concurrently or in series. However, the cause or causes of most strandings are unknown (Geraci *et al.*, 1976; Eaton, 1979; Odell *et al.*, 1980; Best, 1982). Numerous studies suggest that the physiology, behavior, habitat relationships, age, or condition of cetaceans may cause them to strand or might pre-dispose them to strand when exposed to another phenomenon. These suggestions are consistent with the conclusions of numerous other studies that have demonstrated that combinations of dissimilar stressors commonly combine to kill an animal or dramatically reduce its fitness, even though one exposure without the other does not produce the same result (Chrousos, 2000; Creel, 2005; DeVries *et al.*, 2003; Fair and Becker, 2000; Foley *et al.*, 2001; Moberg, 2000; Relyea, 2005a, 2005b; Romero, 2004; Sih *et al.*, 2004).

Strandings Associated With Military Active Sonar—Several sources have published lists of mass stranding events of cetaceans in an attempt to identify relationships between those stranding events and military active sonar (Hildebrand, 2004; IWC, 2005; Taylor *et al.*, 2004). For example, based on a review of stranding records between 1960 and 1995, the International Whaling Commission (2005) identified ten mass stranding events and concluded that, out of eight stranding events reported from the mid-1980s to the summer of 2003, seven had been coincident with the use of mid-frequency active sonar and most involved beaked whales.

Over the past 12 years, there have been five stranding events coincident with military mid-frequency active sonar use in which exposure to sonar is believed to have been a contributing factor to strandings: Greece (1996); the Bahamas (2000); Madeira (2000); Canary Islands (2002); and Spain (2006). Refer to Cox *et al.* (2006) for a summary of common features shared by the strandings events in Greece (1996), Bahamas (2000), Madeira (2000), and Canary Islands (2002); and Fernandez *et al.*, (2005) for an additional summary of the Canary Islands 2002 stranding event.

Potential for Stranding From Seismic Surveys—Marine mammals close to underwater detonations of high explosives can be killed or severely injured, and the auditory organs are especially susceptible to injury (Ketten *et al.*, 1993; Ketten, 1995). However, explosives are no longer used in marine waters for commercial seismic surveys or (with rare exceptions) for seismic research. These methods have been replaced entirely by airguns or related non-explosive pulse generators. Airgun

pulses are less energetic and have slower rise times, and there is no specific evidence that they can cause serious injury, death, or stranding even in the case of large airgun arrays. However, the association of strandings of beaked whales with naval exercises involving mid-frequency active sonar (non-pulse sound) and, in one case, the co-occurrence of an L-DEO seismic survey (Malakoff, 2002; Cox *et al.*, 2006), has raised the possibility that beaked whales exposed to strong “pulsed” sounds could also be susceptible to injury and/or behavioral reactions that can lead to stranding (e.g., Hildebrand, 2005; Southall *et al.*, 2007).

Specific sound-related processes that lead to strandings and mortality are not well documented, but may include:

(1) Swimming in avoidance of a sound into shallow water;

(2) A change in behavior (such as a change in diving behavior) that might contribute to tissue damage, gas bubble formation, hypoxia, cardiac arrhythmia, hypertensive hemorrhage or other forms of trauma;

(3) A physiological change such as a vestibular response leading to a behavioral change or stress-induced hemorrhagic diathesis, leading in turn to tissue damage; and

(4) Tissue damage directly from sound exposure, such as through acoustically-mediated bubble formation and growth or acoustic resonance of tissues.

Some of these mechanisms are unlikely to apply in the case of impulse sounds. However, there are indications that gas-bubble disease (analogous to “the bends”), induced in supersaturated tissue by a behavioral response to acoustic exposure, could be a pathologic mechanism for the strandings and mortality of some deep-diving cetaceans exposed to sonar. The evidence for this remains circumstantial and associated with exposure to naval mid-frequency sonar, not seismic surveys (Cox *et al.*, 2006; Southall *et al.*, 2007).

Seismic pulses and mid-frequency sonar signals are quite different, and some mechanisms by which sonar sounds have been hypothesized to affect beaked whales are unlikely to apply to airgun pulses. Sounds produced by airgun arrays are broadband impulses with most of the energy below one kHz. Typical military mid-frequency sonar emits non-impulse sounds at frequencies of 2 to 10 kHz, generally with a relatively narrow bandwidth at any one time. A further difference between seismic surveys and naval exercises is that naval exercises can involve sound sources on more than one vessel. Thus, it is not appropriate to expect that the same to marine

mammals will result from military sonar and seismic surveys. However, evidence that sonar signals can, in special circumstances, lead (at least indirectly) to physical damage and mortality (e.g., Balcomb and Claridge, 2001; NOAA and USN, 2001; Jepson *et al.*, 2003; Fernández *et al.*, 2004, 2005; Hildebrand 2005; Cox *et al.*, 2006) suggests that caution is warranted when dealing with exposure of marine mammals to any high-intensity sound.

There is no conclusive evidence of cetacean strandings or deaths at sea as a result of exposure to seismic surveys, but a few cases of strandings in the general area where a seismic survey was ongoing have led to speculation concerning a possible link between seismic surveys and strandings.

Suggestions that there was a link between seismic surveys and strandings of humpback whales in Brazil (Engel *et al.*, 2004) were not well founded (IAGC, 2004; IWC, 2007). In September 2002, there was a stranding of two Cuvier’s beaked whales in the Gulf of California, Mexico, when the L-DEO vessel R/V *Maurice Ewing* was operating a 20 airgun (8,490 in³) array in the general area. The link between the stranding and the seismic surveys was inconclusive and not based on any physical evidence (Hogarth, 2002; Yoder, 2002). Nonetheless, the Gulf of California incident plus the beaked whale strandings near naval exercises involving use of mid-frequency sonar suggests a need for caution in conducting seismic surveys in areas occupied by beaked whales until more is known about effects of seismic surveys on those species (Hildebrand, 2005). No injuries of beaked whales are anticipated during the proposed study because of:

(1) The high likelihood that any beaked whales nearby would avoid the approaching vessel before being exposed to high sound levels, and

(2) Differences between the sound sources operated by L-DEO and those involved in the naval exercises associated with strandings.

Non-auditory Physiological Effects—Non-auditory physiological effects or injuries that theoretically might occur in marine mammals exposed to strong underwater sound include stress, neurological effects, bubble formation, resonance, and other types of organ or tissue damage (Cox *et al.*, 2006; Southall *et al.*, 2007). Studies examining such effects are limited. However, resonance effects (Gentry, 2002) and direct noise-induced bubble formations (Crum *et al.*, 2005) are implausible in the case of exposure to an impulsive broadband source like an airgun array. If seismic

surveys disrupt diving patterns of deep-diving species, this might perhaps result in bubble formation and a form of the bends, as speculated to occur in beaked whales exposed to sonar. However, there is no specific evidence of this upon exposure to airgun pulses.

In general, very little is known about the potential for seismic survey sounds (or other types of strong underwater sounds) to cause non-auditory physical effects in marine mammals. Such effects, if they occur at all, would presumably be limited to short distances and to activities that extend over a prolonged period. The available data do not allow identification of a specific exposure level above which non-auditory effects can be expected (Southall *et al.*, 2007), or any meaningful quantitative predictions of the numbers (if any) of marine mammals that might be affected in those ways. Marine mammals that show behavioral avoidance of seismic vessels, including most baleen whales, some odontocetes, and some pinnipeds, are especially unlikely to incur non-auditory physical effects.

Potential Effects of Other Acoustic Devices and Sources

Multibeam Echosounder

NSF and ASC will operate the Simrad EM120 multibeam echosounder from the source vessel during the planned study. Sounds from the multibeam echosounder are very short pulses, occurring for 15 ms, depending on water depth. Most of the energy in the sound pulses emitted by the multibeam echosounder is at frequencies near 12 kHz, and the maximum source level is 242 dB re 1 μ Pa (rms). The beam is narrow (1 to 2°) in fore-aft extent and wide (150°) in the cross-track extent. Each ping consists of nine (in water greater than 1,000 m deep) consecutive successive fan-shaped transmissions (segments) at different cross-track angles. Any given mammal at depth near the trackline would be in the main beam for only one or two of the nine segments. Also, marine mammals that encounter the Simrad EM120 are unlikely to be subjected to repeated pulses because of the narrow fore-aft width of the beam and will receive only limited amounts of pulse energy because of the short pulses. Animals close to the ship (where the beam is narrowest) are especially unlikely to be ensounded for more than one 15 ms pulse (or two pulses if in the overlap area). Similarly, Kremser *et al.* (2005) noted that the probability of a cetacean swimming through the area of exposure when a multibeam echosounder emits a

pulse is small. The animal would have to pass the transducer at close range and be swimming at speeds similar to the vessel in order to receive the multiple pulses that might result in sufficient exposure to cause TTS.

Navy sonars that have been linked to avoidance reactions and stranding of cetaceans: (1) Generally have longer pulse duration than the Simrad EM120; and (2) are often directed close to horizontally, as well as omnidirectional, versus more downward and narrowly for the multibeam echosounder. The area of possible influence of the multibeam echosounder is much smaller—a narrow band below the source vessel. Also, the duration of exposure for a given marine mammal can be much longer for naval sonar. During NSF and ASC's operations, the individual pulses will be very short, and a given mammal would not receive many of the downward-directed pulses as the vessel passes by. Possible effects of a multibeam echosounder on marine mammals are described below.

Masking—Marine mammal communications will not be masked appreciably by the multibeam echosounder signals given the low duty cycle of the echosounder and the brief period when an individual mammal is likely to be within its beam. Furthermore, in the case of baleen whales, the multibeam echosounder signals (12 kHz) do not overlap with the predominant frequencies in the calls, which would avoid any significant masking.

Behavioral Responses—Behavioral reactions of free-ranging marine mammals to sonars, echosounders, and other sound sources appear to vary by species and circumstance. Observed reactions have included silencing and dispersal by sperm whales (Watkins *et al.*, 1985), increased vocalizations and no dispersal by pilot whales (Rendell and Gordon, 1999), and the previously-mentioned beachings by beaked whales. During exposure to a 21 to 25 kHz “whale-finding” sonar with a source level of 215 dB re 1 μ Pa, gray whales reacted by orienting slightly away from the source and being deflected from their course by approximately 200 m (656.2 ft) (Frankel, 2005). When a 38 kHz echosounder and a 150 kHz acoustic Doppler current profiler were transmitting during studies in the Eastern Tropical Pacific, baleen whales showed no significant responses, while spotted and spinner dolphins were detected slightly more often and beaked whales less often during visual surveys (Gerrodette and Pettis, 2005).

Captive bottlenose dolphins and a beluga whale exhibited changes in

behavior when exposed to 1 second tonal signals at frequencies similar to those that will be emitted by the multibeam echosounder used by NSF and ASC, and to shorter broadband pulsed signals. Behavioral changes typically involved what appeared to be deliberate attempts to avoid the sound exposure (Schlundt *et al.*, 2000; Finneran *et al.*, 2002; Finneran and Schlundt, 2004). The relevance of those data to free-ranging odontocetes is uncertain, and in any case, the test sounds were quite different in duration as compared with those from a multibeam echosounder.

Hearing Impairment and Other Physical Effects—Given several stranding events that have been associated with the operation of naval sonar in specific circumstances, there is concern that mid-frequency sonar sounds can cause serious impacts to marine mammals (see above). However, the multibeam echosounder proposed for use by NSF and ASC is quite different than sonar used for Navy operations. Pulse duration of the multibeam echosounder is very short relative to the naval sonar. Also, at any given location, an individual marine mammal would be in the beam of the multibeam echosounder for much less time given the generally downward orientation of the beam and its narrow fore-aft beamwidth; Navy sonar often uses near-horizontally-directed sound. Those factors would all reduce the sound energy received from the multibeam echosounder rather drastically relative to that from naval sonar.

NMFS believes that the brief exposure of marine mammals to one pulse, or small numbers of signals, from the multi-beam echosounder in this particular case is not likely to result in the harassment of marine mammals.

Single-Beam Echosounder

NSF and ASC will operate the Knudsen 3260 and Bathy 2000 single-beam echosounders from the source vessel during the planned study. Sounds from the single-beam echosounder are very short pulses, depending on water depth. Most of the energy in the sound pulses emitted by the singlebeam echosounder is at frequencies near 12 kHz for bottom-tracking purposes or at 3.5 kHz in the sub-bottom profiling mode. The sonar emits energy in a 30° beam from the bottom of the ship. Marine mammals that encounter the Simrad EM120 are unlikely to be subjected to repeated pulses because of the narrow fore-aft width of the beam and will receive only limited amounts of pulse energy

because of the short pulses. Animals close to the ship (where the beam is narrowest) are especially unlikely to be ensounded for more than one 15 ms pulse (or two pulses if in the overlap area). Similarly, Kremser *et al.* (2005) noted that the probability of a cetacean swimming through the area of exposure when a multibeam echosounder emits a pulse is small. The animal would have to pass the transducer at close range and be swimming at speeds similar to the vessel in order to receive the multiple pulses that might result in sufficient exposure to cause TTS.

Navy sonars that have been linked to avoidance reactions and stranding of cetaceans: (1) Generally have longer pulse duration than the Simrad EM120; and (2) are often directed close to horizontally versus more downward for the echosounder. The area of possible influence of the single-beam echosounder is much smaller—a narrow band below the source vessel. Also, the duration of exposure for a given marine mammal can be much longer for naval sonar. During NSF and ASC's operations, the individual pulses will be very short, and a given mammal would not receive many of the downward-directed pulses as the vessel passes by. Possible effects of a single-beam echosounder on marine mammals are described below.

Masking—Marine mammal communications will not be masked appreciably by the single-beam echosounder signals given the low duty cycle of the echosounder and the brief period when an individual mammal is likely to be within its beam. Furthermore, in the case of baleen whales, the single-beam echosounder signals (12 or 3.5 kHz) do not overlap with the predominant frequencies in the calls, which would avoid any significant masking.

Behavioral Responses—Behavioral reactions of free-ranging marine mammals to sonars, echosounders, and other sound sources appear to vary by species and circumstance. Observed reactions have included silencing and dispersal by sperm whales (Watkins *et al.*, 1985), increased vocalizations and no dispersal by pilot whales (Rendell and Gordon, 1999), and the previously-mentioned beachings by beaked whales. During exposure to a 21 to 25 kHz “whale-finding” sonar with a source level of 215 dB re 1 μ Pa, gray whales reacted by orienting slightly away from the source and being deflected from their course by approximately 200 m (656.2 ft) (Frankel, 2005). When a 38 kHz echosounder and a 150 kHz ADCP were transmitting during studies in the Eastern Tropical Pacific, baleen whales

showed no significant responses, while spotted and spinner dolphins were detected slightly more often and beaked whales less often during visual surveys (Gerrodette and Pettis, 2005).

Captive bottlenose dolphins and a beluga whale exhibited changes in behavior when exposed to 1 second tonal signals at frequencies similar to those that will be emitted by the single-beam echosounder used by NSF and ASC, and to shorter broadband pulsed signals. Behavioral changes typically involved what appeared to be deliberate attempts to avoid the sound exposure (Schlundt *et al.*, 2000; Finneran *et al.*, 2002; Finneran and Schlundt, 2004). The relevance of those data to free-ranging odontocetes is uncertain, and in any case, the test sounds were quite different in duration as compared with those from a single-beam echosounder.

Hearing Impairment and Other Physical Effects—Given recent stranding events that have been associated with the operation of naval sonar, there is concern that mid-frequency sonar sounds can cause serious impacts to marine mammals (see above). However, the single-beam echosounder proposed for use by NSF and ASC is quite different than sonar used for Navy operations. Pulse duration of the single-beam echosounder is very short relative to the naval sonar. Also, at any given location, an individual marine mammal would be in the beam of the single-beam echosounder for much less time given the generally downward orientation of the beam and its narrow fore-aft beamwidth; Navy sonar often uses near-horizontally-directed sound. Those factors would all reduce the sound energy received from the single-beam echosounder rather drastically relative to that from naval sonar.

NMFS believes that the brief exposure of marine mammals to one pulse, or small numbers of signals, from the single-beam echosounder in this particular case is not likely to result in the harassment of marine mammals.

Acoustic Doppler Current Profilers

NSF and ASC will operate the ADCP Teledyne RDI VM-150 and ADCP Ocean Surveyor OS-38 from the source vessel during the planned study. Most of the energy in the sound pulses emitted by the ADCPs operate at frequencies near 150 kHz, and the maximum source level is 223.6 dB re 1 μ Pa (rms). Sound energy from the ADCP is emitted as a 30° conically-shaped beam. Marine mammals that encounter the ADCPs are unlikely to be subjected to repeated pulses because of the narrow fore-aft width of the beam and will receive only limited amounts of pulse energy

because of the short pulses. Animals close to the ship (where the beam is narrowest) are especially unlikely to be ensounded for more than one 15 ms pulse (or two pulses if in the overlap area). Similarly, Kremser *et al.* (2005) noted that the probability of a cetacean swimming through the area of exposure when the ADCPs emits a pulse is small. The animal would have to pass the transducer at close range and be swimming at speeds similar to the vessel in order to receive the multiple pulses that might result in sufficient exposure to cause TTS.

Navy sonars that have been linked to avoidance reactions and stranding of cetaceans: (1) Generally have longer pulse duration than the ADCPs; and (2) are often directed close to horizontally versus more downward for the ADCPs. The area of possible influence of the multibeam echosounder is much smaller—a narrow band below the source vessel. Also, the duration of exposure for a given marine mammal can be much longer for naval sonar. During NSF and ASC's operations, the individual pulses will be very short, and a given mammal would not receive many of the downward-directed pulses as the vessel passes by. Possible effects of the ADCPs on marine mammals are described below.

Masking—Marine mammal communications will not be masked appreciably by the ADCP signals given the low duty cycle of the ADCPs and the brief period when an individual mammal is likely to be within its beam. Furthermore, in the case of baleen whales, the ADCP signals (150 kHz) do not overlap with the predominant frequencies in the calls, which would avoid any significant masking.

Behavioral Responses—Behavioral reactions of free-ranging marine mammals to sonars, echosounders, and other sound sources appear to vary by species and circumstance. Observed reactions have included silencing and dispersal by sperm whales (Watkins *et al.*, 1985), increased vocalizations and no dispersal by pilot whales (Rendell and Gordon, 1999), and the previously-mentioned beachings by beaked whales. During exposure to a 21 to 25 kHz “whale-finding” sonar with a source level of 215 dB re 1 μ Pa, gray whales reacted by orienting slightly away from the source and being deflected from their course by approximately 200 m (656.2 ft) (Frankel, 2005). When a 38 kHz echosounder and a 150 kHz ADCP were transmitting during studies in the Eastern Tropical Pacific, baleen whales showed no significant responses, while spotted and spinner dolphins were detected slightly more often and beaked

whales less often during visual surveys (Gerrodette and Pettis, 2005).

Captive bottlenose dolphins and a beluga whale exhibited changes in behavior when exposed to 1 second tonal signals at frequencies similar to those that will be emitted by the multibeam echosounder used by NSF and ASC, and to shorter broadband pulsed signals. Behavioral changes typically involved what appeared to be deliberate attempts to avoid the sound exposure (Schlundt *et al.*, 2000; Finneran *et al.*, 2002; Finneran and Schlundt, 2004). The relevance of those data to free-ranging odontocetes is uncertain, and in any case, the test sounds were quite different in duration as compared with those from a multibeam echosounder.

Hearing Impairment and Other Physical Effects—Given recent stranding events that have been associated with the operation of naval sonar, there is concern that mid-frequency sonar sounds can cause serious impacts to marine mammals (see above). However, the multibeam echosounder proposed for use by NSF and ASC is quite different than sonar used for Navy operations. Pulse duration of the ADCP is very short relative to the naval sonar. Also, at any given location, an individual marine mammal would be in the beam of the multibeam echosounder for much less time given the generally downward orientation of the beam and its narrow fore-aft beamwidth; Navy sonar often uses near-horizontally-directed sound. Those factors would all reduce the sound energy received from the multibeam echosounder rather drastically relative to that from naval sonar.

NMFS believes that the brief exposure of marine mammals to one pulse, or small numbers of signals, from the multi-beam echosounder in this particular case is not likely to result in the harassment of marine mammals.

Acoustic Locator

NSF and ASC will operate the acoustic locator from the source vessel during the planned study during sampling. Sounds from the locator are very short pulses, occurring for 5 ms. Most of the energy in the sound pulses emitted by the acoustic locator is at frequencies near 12 kHz, and the maximum source level is 162 dB re 1 μ Pa (rms). Animals close to the ship (where the beam is narrowest) are especially unlikely to be ensounded for more than one 5 ms pulse (or two pulses if in the overlap area). Similarly, Kremser *et al.* (2005) noted that the probability of a cetacean swimming through the area of exposure when a

multibeam echosounder emits a pulse is small. The animal would have to pass the transducer at close range and be swimming at speeds similar to the vessel in order to receive the multiple pulses that might result in sufficient exposure to cause TTS.

Masking—Marine mammal communications will not be masked appreciably by the acoustic locator signals given the low duty cycle and the low source level. Furthermore, in the case of baleen whales, the acoustic locator signals (12 kHz) do not overlap with the predominant frequencies in the calls, which would avoid any significant masking.

Behavioral Responses—Behavioral reactions of free-ranging marine mammals to sonars, echosounders, and other sound sources appear to vary by species and circumstance. Observed reactions have included silencing and dispersal by sperm whales (Watkins *et al.*, 1985), increased vocalizations and no dispersal by pilot whales (Rendell and Gordon, 1999), and the previously-mentioned beachings by beaked whales. During exposure to a 21 to 25 kHz “whale-finding” sonar with a source level of 215 dB re 1 μ Pa, gray whales reacted by orienting slightly away from the source and being deflected from their course by approximately 200 m (656.2 ft) (Frankel, 2005). When a 38 kHz echosounder and a 150 kHz ADCP were transmitting during studies in the Eastern Tropical Pacific, baleen whales showed no significant responses, while spotted and spinner dolphins were detected slightly more often and beaked whales less often during visual surveys (Gerrodette and Pettis, 2005).

NMFS believes that the brief exposure of marine mammals to one pulse, or small numbers of signals, from the acoustic locator is not likely to result in the harassment of marine mammals.

Core and Dredge Sampling

During coring and dredging, the noise created by the mechanical action of the devices on the seafloor is expected to be perceived by nearby fish and other marine organisms and deter them from swimming toward the source. Coring and dredging activities would be highly localized and short-term in duration and would not be expected to significantly interfere with marine mammal behavior. The potential direct effects include temporary localized disturbance or displacement from associated sounds and/or physical movement/actions of the operations. Additionally, the potential indirect effects may consist of very localized and transitory/short-term disturbance of bottom habitat and associated prey in shallow-water areas

as a result of coring, dredging, and sediment sampling (NSF/USGS PEIS, 2011). NMFS believes that the brief exposure of marine mammals to noise created from the mechanical action of the devices for core and dredge sampling is not likely to result in the harassment of marine mammals.

Vessel Movement and Collisions

Vessel movement in the vicinity of marine mammals has the potential to result in either a behavioral response or a direct physical interaction. Both scenarios are discussed below in this section.

Behavioral Responses to Vessel Movement—There are limited data concerning marine mammal behavioral responses to vessel traffic and vessel noise, and a lack of consensus among scientists with respect to what these responses mean or whether they result in short-term or long-term adverse effects. In those cases where there is a busy shipping lane or where there is a large amount of vessel traffic, marine mammals (especially low frequency specialists) may experience acoustic masking (Hildebrand, 2005) if they are present in the area (e.g., killer whales in Puget Sound; Foote *et al.*, 2004; Holt *et al.*, 2008). In cases where vessels actively approach marine mammals (e.g., whale watching or dolphin watching boats), scientists have documented that animals exhibit altered behavior such as increased swimming speed, erratic movement, and active avoidance behavior (Bursk, 1983; Acevedo, 1991; Baker and MacGibbon, 1991; Trites and Bain, 2000; Williams *et al.*, 2002; Constantine *et al.*, 2003), reduced blow interval (Ritcher *et al.*, 2003), disruption of normal social behaviors (Lusseau, 2003, 2006), and the shift of behavioral activities which may increase energetic costs (Constantine *et al.*, 2003, 2004). A detailed review of marine mammal reactions to ships and boats is available in Richardson *et al.*, (1995). For each of the marine mammal taxonomy groups, Richardson *et al.*, (1995) provides the following assessment regarding reactions to vessel traffic:

Toothed whales—“In summary, toothed whales sometimes show no avoidance reaction to vessels, or even approach them. However, avoidance can occur, especially in response to vessels of types used to chase or hunt the animals. This may cause temporary displacement, but we know of no clear evidence that toothed whales have abandoned significant parts of their range because of vessel traffic.”

Baleen whales—“When baleen whales receive low-level sounds from distant or

stationary vessels, the sounds often seem to be ignored. Some whales approach the sources of these sounds. When vessels approach whales slowly and non-aggressively, whales often exhibit slow and inconspicuous avoidance maneuvers. In response to strong or rapidly changing vessel noise, baleen whales often interrupt their normal behavior and swim rapidly away. Avoidance is especially strong when a boat heads directly toward the whale.”

Behavioral responses to stimuli are complex and influenced to varying degrees by a number of factors, such as species, behavioral contexts, geographical regions, source characteristics (moving or stationary, speed, direction, etc.), prior experience of the animal and physical status of the animal. For example, studies have shown that beluga whales’ reaction varied when exposed to vessel noise and traffic. In some cases, beluga whales exhibited rapid swimming from ice-breaking vessels up to 80 km (43.2 nmi) away and showed changes in surfacing, breathing, diving, and group composition in the Canadian high Arctic where vessel traffic is rare (Finley *et al.*, 1990). In other cases, beluga whales were more tolerant of vessels, but responded differentially to certain vessels and operating characteristics by reducing their calling rates (especially older animals) in the St. Lawrence River where vessel traffic is common (Blane and Jaakson, 1994). In Bristol Bay, Alaska, beluga whales continued to feed when surrounded by fishing vessels and resisted dispersal even when purposefully harassed (Fish and Vania, 1971).

In reviewing more than 25 years of whale observation data, Watkins (1986) concluded that whale reactions to vessel traffic were “modified by their previous experience and current activity: habituation often occurred rapidly, attention to other stimuli or preoccupation with other activities sometimes overcame their interest or wariness of stimuli.” Watkins noticed that over the years of exposure to ships in the Cape Cod area, minke whales changed from frequent positive interest (e.g., approaching vessels) to generally uninterested reactions; fin whales changed from mostly negative (e.g., avoidance) to uninterested reactions; fin whales changed from mostly negative (e.g., avoidance) to uninterested reactions; right whales apparently continued the same variety of responses (negative, uninterested, and positive responses) with little change; and humpbacks dramatically changed from mixed responses that were often

negative to reactions that were often strongly positive. Watkins (1986) summarized that “whales near shore, even in regions with low vessel traffic, generally have become less wary of boats and their noises, and they have appeared to be less easily disturbed than previously. In particular locations with intense shipping and repeated approaches by boats (such as the whale-watching areas of Stellwagen Bank), more and more whales had positive reactions to familiar vessels, and they also occasionally approached other boats and yachts in the same ways.”

Although the radiated sound from the *Palmer* will be audible to marine mammals over a large distance, it is unlikely that marine mammals will respond behaviorally (in a manner that NMFS would consider harassment under the MMPA) to low-level distant shipping noise as the animals in the area are likely to be habituated to such noises (Nowacek *et al.*, 2004). In light of these facts, NMFS does not expect the *Palmer*'s movements to result in Level B harassment.

Vessel Strike—Ship strikes of cetaceans can cause major wounds, which may lead to the death of the animal. An animal at the surface could be struck directly by a vessel, a surfacing animal could hit the bottom of a vessel, or an animal just below the surface could be cut by a vessel's propeller. The severity of injuries typically depends on the size and speed of the vessel (Knowlton and Kraus, 2001; Laist *et al.*, 2001; Vanderlaan and Taggart, 2007).

The most vulnerable marine mammals are those that spend extended periods of time at the surface in order to restore oxygen levels within their tissues after deep dives (e.g., the sperm whale). In addition, some baleen whales, such as the North Atlantic right whale, seem generally unresponsive to vessel sound, making them more susceptible to vessel collisions (Nowacek *et al.*, 2004). These species are primarily large, slow moving whales. Smaller marine mammals (e.g., bottlenose dolphin) move quickly through the water column and are often seen riding the bow wave of large ships. Marine mammal responses to vessels may include avoidance and changes in dive pattern (NRC, 2003).

An examination of all known ship strikes from all shipping sources (civilian and military) indicates vessel speed is a principal factor in whether a vessel strike results in death (Knowlton and Kraus, 2001; Laist *et al.*, 2001; Jensen and Silber, 2003; Vanderlaan and Taggart, 2007). In assessing records in which vessel speed was known, Laist *et al.* (2001) found a direct relationship

between the occurrence of a whale strike and the speed of the vessel involved in the collision. The authors concluded that most deaths occurred when a vessel was traveling in excess of 13 kts (24.1 km/hr, 14.9 mph).

NSF and ASC's proposed operation of one source vessel for the proposed low-energy seismic survey is relatively small in scale compared to the number of commercial ships transiting at higher speeds in the same areas on an annual basis. The probability of vessel and marine mammal interactions occurring during the proposed low-energy seismic survey is unlikely due to the *Palmer*'s slow operational speed, which is typically 5 kts. Outside of seismic operations, the *Palmer*'s cruising speed would be approximately 10.1 to 14.5 kts, which is generally below the speed at which studies have noted reported increases of marine mammal injury or death (Laist *et al.*, 2001).

As a final point, the *Palmer* has a number of other advantages for avoiding ship strikes as compared to most commercial merchant vessels, including the following: the *Palmer*'s bridge and aloft observation tower offers good visibility to visually monitor for marine mammal presence; PSOs posted during operations scan the ocean for marine mammals and must report visual alerts of marine mammal presence to crew; and the PSOs receive extensive training that covers the fundamentals of visual observing for marine mammals and information about marine mammals and their identification at sea.

Entanglement

Entanglement can occur if wildlife becomes immobilized in survey lines, cables, nets, or other equipment that is moving through the water column. The proposed low-energy seismic survey would require towing approximately a single 100 m cable streamer. This large of an array carries the risk of entanglement for marine mammals. Wildlife, especially slow moving individuals, such as large whales, have a low probability of becoming entangled due to slow speed of the survey vessel and onboard monitoring efforts. In May 2011, there was one recorded entanglement of an olive ridley sea turtle (*Lepidochelys olivacea*) in the R/V *Marcus G. Langseth*'s barovanes after the conclusion of a seismic survey off Costa Rica. There have been cases of baleen whales, mostly gray whales (Heyning, 1990), becoming entangled in fishing lines. The probability for entanglement of marine mammals is considered not significant because of the vessel speed and the monitoring efforts onboard the survey vessel.

Icebreaking Activities

Icebreakers produce more noise while breaking ice than ships of comparable size due, primarily, to the sounds of propeller cavitating (Richardson *et al.*, 1995). Multi-year ice, which is expected to be encountered in the proposed survey area. Icebreakers commonly back and ram into heavy ice until losing momentum to make way. The highest noise levels usually occur while backing full astern in preparation to ram forward through the ice. Overall the noise generated by an icebreaker pushing ice was 10 to 15 dB greater than the noise produced by the ship underway in open water (Richardson *et al.*, 1995). In general, the Antarctic and Southern Ocean is a noisy environment. Calving and grounding icebergs as well as the break-up of ice sheets, can produce a large amount of underwater noise. Little information is available about the increased sound levels due to icebreaking.

Cetaceans—Few studies have been conducted to evaluate the potential interference of icebreaking noise with marine mammal vocalizations. Erbe and Farmer (1998) measured masked hearing thresholds of a captive beluga whale. They reported that the recording of a CCG ship, *Henry Larsen*, ramming ice in the Beaufort Sea, masked recordings of beluga vocalizations at a noise to signal pressure ratio of 18 dB, when the noise pressure level was eight times as high as the call pressure. Erbe and Farmer (2000) also predicted when icebreaker noise would affect beluga whales through software that combined a sound propagation model and beluga whale impact threshold models. They again used the data from the recording of the *Henry Larsen* in the Beaufort Sea and predicted that masking of beluga whale vocalizations could extend between 40 and 71 km (21.6 and 38.3 nmi) near the surface. Lesage *et al.* (1999) report that beluga whales changed their call type and call frequency when exposed to boat noise. It is possible that the whales adapt to the ambient noise levels and are able to communicate despite the sound. Given the documented reaction of belugas to ships and icebreakers it is highly unlikely that beluga whales would remain in the proximity of vessels where vocalizations would be masked.

Beluga whales have been documented swimming rapidly away from ships and icebreakers in the Canadian high Arctic when a ship approaches to within 35 to 50 km (18.9 to 27 nmi), and they may travel up to 80 km (43.2 nmi) from the vessel's track (Richardson *et al.*, 1995). It is expected that belugas avoid

icebreakers as soon as they detect the ships (Cosens and Dueck, 1993). However, the reactions of beluga whales to ships vary greatly and some animals may become habituated to high levels of ambient noise (Erbe and Darmber, 2000).

There is little information about the effects of icebreaking ships on baleen whales. Migrating bowhead whales appeared to avoid an area around a drill site by greater than 25 km (13.5 mi) where an icebreaker was working in the Beaufort Sea. There was intensive icebreaking daily in support of the drilling activities (Brewer *et al.*, 1993). Migrating bowheads also avoided a nearby drill site at the same time of year where little icebreaking was being conducted (LGL and Greeneridge, 1987). It is unclear as to whether the drilling activities, icebreaking operations, or the ice itself might have been the cause for the whale's diversion. Bowhead whales are not expected to occur in the proximity of the proposed action area.

Pinnipeds—Brueggeman *et al.* (1992) reported on the reactions of seals to an icebreaker during activities at two prospects in the Chukchi Sea. Reactions of seals to the icebreakers varied between the two prospects. Most (67%) seals did not react to the icebreaker at either prospect. Reaction at one prospect was greatest during icebreaking activity (running/maneuvering/jogging) and was 0.23 km (0.12 nmi) of the vessel and lowest for animals beyond 0.93 km (0.5 nmi). At the second prospect however, seal reaction was lowest during icebreaking activity with higher and similar levels of response during general (non-icebreaking) vessel operations and when the vessel was at anchor or drifting. The frequency of seal reaction generally declined with increasing distance from the vessel except during general vessel activity where it remained consistently high to about 0.46 km (0.25 nmi) from the vessel before declining.

Similarly, Kanik *et al.* (1980) found that ringed (*Pusa hispida*) and harp seals (*Pagophilus groenlandicus*) often dove into the water when an icebreaker was breaking ice within 1 km (0.5 nmi) of the animals. Most seals remained on the ice when the ship was breaking ice 1 to 2 km (0.5 to 1.1 nmi) away.

The potential effects to marine mammals described in this section of the document do not take into consideration the proposed monitoring and mitigation measures described later in this document (see the "Proposed Mitigation" and "Proposed Monitoring and Reporting" sections) which, as noted are designed to effect the least

practicable adverse impact on affected marine mammal species and stocks.

Anticipated Effects on Marine Mammal Habitat

The proposed seismic survey is not anticipated to have any permanent impact on habitats used by the marine mammals in the proposed survey area, including the food sources they use (i.e. fish and invertebrates). Additionally, no physical damage to any habitat is anticipated as a result of conducting airgun operations during the proposed low-energy seismic survey. While it is anticipated that the specified activity may result in marine mammals avoiding certain areas due to temporary ensonification, this impact to habitat is temporary and was considered in further detail earlier in this document, as behavioral modification. The main impact associated with the proposed activity will be temporarily elevated noise levels and the associated direct effects on marine mammals in any particular area of the approximately 5,628 km² proposed project area, previously discussed in this notice.

The *Palmer* is designed for continuous passage at 3 kts through ice 1 m thick. During the proposed project the *Palmer* will typically encounter first- or second-year ice while avoiding thicker ice floes, particularly large intact multi-year ice, whenever possible. In addition, the vessel will follow leads when possible while following the survey route. As the vessel passes through the ice, the ship causes the ice to part and travel alongside the hull. This ice typically returns to fill the wake as the ship passes. The effects are transitory (i.e., hours at most) and localized (i.e., constrained to a relatively narrow swath perhaps 10 m (32.1 ft) to each side of the vessel. The *Palmer's* maximum beam is 18.3 m (60 ft). Applying the maximum estimated amount of icebreaking (1,000 km), to the corridor opened by the ship, NSF and ASC anticipate that a maximum of approximately 18 km² (5.3 nmi²) of ice may be disturbed. This represents an inconsequential amount of the total ice present in the Southern Ocean.

Sea ice is important for pinniped life functions such as resting, breeding, and molting. Icebreaking activities may damage seal breathing holes and will also reduce the haul-out area in the immediate vicinity of the ship's track. Icebreaking along a maximum of 1,000 km of trackline will alter local ice conditions in the immediate vicinity of the vessel. This has the potential to temporarily lead to a reduction of suitable seal haul-out habitat. However, the dynamic sea-ice environment

requires that seals be able to adapt to changes in sea, ice, and snow conditions, and they therefore create new breathing holes and lairs throughout the winter and spring (Hammill and Smith, 1989). In addition, seals often use open leads and cracks in the ice to surface and breathe (Smith and Stirling, 1975). Disturbance of the ice will occur in a very small area relative to the Southern Ocean ice-pack and no significant impact on marine mammals is anticipated by icebreaking during the proposed low-energy seismic survey. The next section discusses the potential impacts of anthropogenic sound sources on common marine mammal prey in the proposed survey area (i.e., fish and invertebrates).

Anticipated Effects on Fish

One reason for the adoption of airguns as the standard energy source for marine seismic surveys is that, unlike explosives, they have not been associated with large-scale fish kills. However, existing information on the impacts of seismic surveys on marine fish and invertebrate populations is limited. There are three types of potential effects of exposure to seismic surveys: (1) Pathological, (2) physiological, and (3) behavioral. Pathological effects involve lethal and temporary or permanent sub-lethal injury. Physiological effects involve temporary and permanent primary and secondary stress responses, such as changes in levels of enzymes and proteins. Behavioral effects refer to temporary and (if they occur) permanent changes in exhibited behavior (e.g., startle and avoidance behavior). The three categories are interrelated in complex ways. For example, it is possible that certain physiological and behavioral changes could potentially lead to an ultimate pathological effect on individuals (i.e., mortality).

The specific received sound levels at which permanent adverse effects to fish potentially could occur are little studied and largely unknown. Furthermore, the available information on the impacts of seismic surveys on marine fish is from studies of individuals or portions of a population; there have been no studies at the population scale. The studies of individual fish have often been on caged fish that were exposed to airgun pulses in situations not representative of an actual seismic survey. Thus, available information provides limited insight on possible real-world effects at the ocean or population scale. This makes drawing conclusions about impacts on fish problematic because, ultimately, the most important issues concern effects on marine fish populations, their

viability, and their availability to fisheries.

Hastings and Popper (2005), Popper (2009), and Popper and Hastings (2009a, b) provided recent critical reviews of the known effects of sound on fish. The following sections provide a general synopsis of the available information on the effects of exposure to seismic and other anthropogenic sound as relevant to fish. The information comprises results from scientific studies of varying degrees of rigor plus some anecdotal information. Some of the data sources may have serious shortcomings in methods, analysis, interpretation, and reproducibility that must be considered when interpreting their results (see Hastings and Popper, 2005). Potential adverse effects of the program's sound sources on marine fish are noted.

Pathological Effects—The potential for pathological damage to hearing structures in fish depends on the energy level of the received sound and the physiology and hearing capability of the species in question. For a given sound to result in hearing loss, the sound must exceed, by some substantial amount, the hearing threshold of the fish for that sound (Popper, 2005). The consequences of temporary or permanent hearing loss in individual fish on a fish population are unknown; however, they likely depend on the number of individuals affected and whether critical behaviors involving sound (e.g., predator avoidance, prey capture, orientation and navigation, reproduction, etc.) are adversely affected.

Little is known about the mechanisms and characteristics of damage to fish that may be inflicted by exposure to seismic survey sounds. Few data have been presented in the peer-reviewed scientific literature. As far as NSF, ASC, and NMFS know, there are only two papers with proper experimental methods, controls, and careful pathological investigation implicating sounds produced by actual seismic survey airguns in causing adverse anatomical effects. One such study indicated anatomical damage, and the second indicated TTS in fish hearing. The anatomical case is McCauley *et al.* (2003), who found that exposure to airgun sound caused observable anatomical damage to the auditory maculae of pink snapper (*Pagrus auratus*). This damage in the ears had not been repaired in fish sacrificed and examined almost two months after exposure. On the other hand, Popper *et al.* (2005) documented only TTS (as determined by auditory brainstem response) in two of three fish species from the Mackenzie River Delta. This

study found that broad whitefish (*Coregonus nasus*) exposed to five airgun shots were not significantly different from those of controls. During both studies, the repetitive exposure to sound was greater than would have occurred during a typical seismic survey. However, the substantial low-frequency energy produced by the airguns (less than 400 Hz in the study by McCauley *et al.* [2003] and less than approximately 200 Hz in Popper *et al.* [2005]) likely did not propagate to the fish because the water in the study areas was very shallow (approximately nine m in the former case and less than two m in the latter). Water depth sets a lower limit on the lowest sound frequency that will propagate (the “cutoff frequency”) at about one-quarter wavelength (Urick, 1983; Rogers and Cox, 1988).

Wardle *et al.* (2001) suggested that in water, acute injury and death of organisms exposed to seismic energy depends primarily on two features of the sound source: (1) The received peak pressure, and (2) the time required for the pressure to rise and decay. Generally, as received pressure increases, the period for the pressure to rise and decay decreases, and the chance of acute pathological effects increases. According to Buchanan *et al.* (2004), for the types of seismic airguns and arrays involved with the proposed program, the pathological (mortality) zone for fish would be expected to be within a few meters of the seismic source. Numerous other studies provide examples of no fish mortality upon exposure to seismic sources (Falk and Lawrence, 1973; Holliday *et al.*, 1987; La Bella *et al.*, 1996; Santulli *et al.*, 1999; McCauley *et al.*, 2000a, b, 2003; Bjarti, 2002; Thomsen, 2002; Hassel *et al.*, 2003; Popper *et al.*, 2005; Boeger *et al.*, 2006).

An experiment of the effects of a single 700 in³ airgun was conducted in Lake Meade, Nevada (USGS, 1999). The data were used in an Environmental Assessment of the effects of a marine reflection survey of the Lake Meade fault system by the National Park Service (Paulson *et al.*, 1993, in USGS, 1999). The airgun was suspended 3.5 m (11.5 ft) above a school of threadfin shad in Lake Meade and was fired three successive times at a 30 second interval. Neither surface inspection nor diver observations of the water column and bottom found any dead fish.

For a proposed seismic survey in Southern California, USGS (1999) conducted a review of the literature on the effects of airguns on fish and fisheries. They reported a 1991 study of the Bay Area Fault system from the

continental shelf to the Sacramento River, using a 10 airgun (5,828 in³) array. Brezzina and Associates were hired by USGS to monitor the effects of the surveys and concluded that airgun operations were not responsible for the death of any of the fish carcasses observed. They also concluded that the airgun profiling did not appear to alter the feeding behavior of sea lions, seals, or pelicans observed feeding during the seismic surveys.

Some studies have reported, some equivocally, that mortality of fish, fish eggs, or larvae can occur close to seismic sources (Kostyuchenko, 1973; Dalen and Knutsen, 1986; Booman *et al.*, 1996; Dalen *et al.*, 1996). Some of the reports claimed seismic effects from treatments quite different from actual seismic survey sounds or even reasonable surrogates. However, Payne *et al.* (2009) reported no statistical differences in mortality/morbidity between control and exposed groups of capelin eggs or monkfish larvae. Saetre and Ona (1996) applied a ‘worst-case scenario’ mathematical model to investigate the effects of seismic energy on fish eggs and larvae. They concluded that mortality rates caused by exposure to seismic surveys are so low, as compared to natural mortality rates, that the impact of seismic surveying on recruitment to a fish stock must be regarded as insignificant.

Physiological Effects—Physiological effects refer to cellular and/or biochemical responses of fish to acoustic stress. Such stress potentially could affect fish populations by increasing mortality or reducing reproductive success. Primary and secondary stress responses of fish after exposure to seismic survey sound appear to be temporary in all studies done to date (Sverdrup *et al.*, 1994; Santulli *et al.*, 1999; McCauley *et al.*, 2000a, b). The periods necessary for the biochemical changes to return to normal are variable and depend on numerous aspects of the biology of the species and of the sound stimulus.

Behavioral Effects—Behavioral effects include changes in the distribution, migration, mating, and catchability of fish populations. Studies investigating the possible effects of sound (including seismic survey sound) on fish behavior have been conducted on both uncaged and caged individuals (e.g., Chapman and Hawkins, 1969; Pearson *et al.*, 1992; Santulli *et al.*, 1999; Wardle *et al.*, 2001; Hassel *et al.*, 2003). Typically, in these studies fish exhibited a sharp startle response at the onset of a sound followed by habituation and a return to normal behavior after the sound ceased.

The Minerals Management Service (MMS, 2005) assessed the effects of a proposed seismic survey in Cook Inlet. The seismic survey proposed using three vessels, each towing two, four-airgun arrays ranging from 1,500 to 2,500 in³. MMS noted that the impact to fish populations in the survey area and adjacent waters would likely be very low and temporary. MMS also concluded that seismic surveys may displace the pelagic fishes from the area temporarily when airguns are in use. However, fishes displaced and avoiding the airgun noise are likely to backfill the survey area in minutes to hours after cessation of seismic testing. Fishes not dispersing from the airgun noise (e.g., demersal species) may startle and move short distances to avoid airgun emissions.

In general, any adverse effects on fish behavior or fisheries attributable to seismic testing may depend on the species in question and the nature of the fishery (season, duration, fishing method). They may also depend on the age of the fish, its motivational state, its size, and numerous other factors that are difficult, if not impossible, to quantify at this point, given such limited data on effects of airguns on fish, particularly under realistic at-sea conditions.

Anticipated Effects on Invertebrates

The existing body of information on the impacts of seismic survey sound on marine invertebrates is very limited. However, there is some unpublished and very limited evidence of the potential for adverse effects on invertebrates, thereby justifying further discussion and analysis of this issue. The three types of potential effects of exposure to seismic surveys on marine invertebrates are pathological, physiological, and behavioral. Based on the physical structure of their sensory organs, marine invertebrates appear to be specialized to respond to particle displacement components of an impinging sound field and not to the pressure component (Popper *et al.*, 2001).

The only information available on the impacts of seismic surveys on marine invertebrates involves studies of individuals; there have been no studies at the population scale. Thus, available information provides limited insight on possible real-world effects at the regional or ocean scale. The most important aspect of potential impacts concerns how exposure to seismic survey sound ultimately affects invertebrate populations and their viability, including availability to fisheries.

Literature reviews of the effects of seismic and other underwater sound on invertebrates were provided by Moriyasu *et al.* (2004) and Payne *et al.* (2008). The following sections provide a synopsis of available information on the effects of exposure to seismic survey sound on species of decapod crustaceans and cephalopods, the two taxonomic groups of invertebrates on which most such studies have been conducted. The available information is from studies with variable degrees of scientific soundness and from anecdotal information. A more detailed review of the literature on the effects of seismic survey sound on invertebrates is provided in Appendix D of NSF/USGS's PEIS.

Pathological Effects—In water, lethal and sub-lethal injury to organisms exposed to seismic survey sound appears to depend on at least two features of the sound source: (1) The received peak pressure; and (2) the time required for the pressure to rise and decay. Generally, as received pressure increases, the period for the pressure to rise and decay decreases, and the chance of acute pathological effects increases. For the type of airgun array planned for the proposed program, the pathological (mortality) zone for crustaceans and cephalopods is expected to be within a few meters of the seismic source, at most; however, very few specific data are available on levels of seismic signals that might damage these animals. This premise is based on the peak pressure and rise/decay time characteristics of seismic airgun arrays currently in use around the world.

Some studies have suggested that seismic survey sound has a limited pathological impact on early developmental stages of crustaceans (Pearson *et al.*, 1994; Christian *et al.*, 2003; DFO, 2004). However, the impacts appear to be either temporary or insignificant compared to what occurs under natural conditions. Controlled field experiments on adult crustaceans (Christian *et al.*, 2003, 2004; DFO, 2004) and adult cephalopods (McCauley *et al.*, 2000a, b) exposed to seismic survey sound have not resulted in any significant pathological impacts on the animals. It has been suggested that exposure to commercial seismic survey activities has injured giant squid (Guerra *et al.*, 2004), but the article provides little evidence to support this claim. Tenera Environmental (2011b) reported that Norris and Mohl (1983, summarized in Moriyasu *et al.*, 2004) observed lethal effects in squid (*Loligo vulgaris*) at levels of 246 to 252 dB after 3 to 11 minutes.

Andre *et al.* (2011) exposed four species of cephalopods (*Loligo vulgaris*, *Sepia officinalis*, *Octopus vulgaris*, and *Ilex coindetii*), primarily cuttlefish, to two hours of continuously 50 to 400 Hz sinusoidal wave sweeps at 157±5 dB re 1 µPa while captive in relatively small tanks. They reported morphological and ultrastructural evidence of massive acoustic trauma (i.e., permanent and substantial alterations [lesions] of statocyst sensory hair cells) to the exposed animals that increased in severity with time, suggesting that cephalopods are particularly sensitive to low frequency sound. The received SPL was reported as 157±5 dB re 1 µPa, with peak levels at 175 dB re 1 µPa. As in the McCauley *et al.* (2003) paper on sensory hair cell damage in pink snapper as a result of exposure to seismic sound, the cephalopods were subjected to higher sound levels than they would be under natural conditions, and they were unable to swim away from the sound source.

Physiological Effects—Physiological effects refer mainly to biochemical responses by marine invertebrates to acoustic stress. Such stress potentially could affect invertebrate populations by increasing mortality or reducing reproductive success. Primary and secondary stress responses (i.e., changes in haemolymph levels of enzymes, proteins, etc.) of crustaceans have been noted several days or months after exposure to seismic survey sounds (Payne *et al.*, 2007). It was noted however, that no behavioral impacts were exhibited by crustaceans (Christian *et al.*, 2003, 2004; DFO, 2004). The periods necessary for these biochemical changes to return to normal are variable and depend on numerous aspects of the biology of the species and of the sound stimulus.

Behavioral Effects—There is increasing interest in assessing the possible direct and indirect effects of seismic and other sounds on invertebrate behavior, particularly in relation to the consequences for fisheries. Changes in behavior could potentially affect such aspects as reproductive success, distribution, susceptibility to predation, and catchability by fisheries. Studies investigating the possible behavioral effects of exposure to seismic survey sound on crustaceans and cephalopods have been conducted on both uncaged and caged animals. In some cases, invertebrates exhibited startle responses (e.g., squid in McCauley *et al.*, 2000a, b). In other cases, no behavioral impacts were noted (e.g., crustaceans in Christian *et al.*, 2003, 2004; DFO 2004). There have been anecdotal reports of

reduced catch rates of shrimp shortly after exposure to seismic surveys; however, other studies have not observed any significant changes in shrimp catch rate (Andriquetto-Filho *et al.*, 2005). Similarly, Parry and Gason (2006) did not find any evidence that lobster catch rates were affected by seismic surveys. Any adverse effects on crustacean and cephalopod behavior or fisheries attributable to seismic survey sound depend on the species in question and the nature of the fishery (season, duration, fishing method).

Proposed Mitigation

In order to issue an Incidental Take Authorization (ITA) under section 101(a)(5)(D) of the MMPA, NMFS must set forth the permissible methods of taking pursuant to such activity, and other means of effecting the least practicable impact on such species or stock and its habitat, paying particular attention to rookeries, mating grounds, and areas of similar significance, and the availability of such species or stock for taking for certain subsistence uses.

NSF and ASC reviewed the following source documents and have incorporated a suite of appropriate mitigation measures into their project description.

(1) Protocols used during previous NSF and USGS-funded seismic research cruises as approved by NMFS and detailed in the recently completed "Final Programmatic Environmental Impact Statement/Overseas Environmental Impact Statement for Marine Seismic Research Funded by the National Science Foundation or Conducted by the U.S. Geological Survey;"

(2) Previous IHA applications and IHAs approved and authorized by NMFS; and

(3) Recommended best practices in Richardson *et al.* (1995), Pierson *et al.* (1998), and Weir and Dolman, (2007).

To reduce the potential for disturbance from acoustic stimuli associated with the activities, NSF, ASC and/or its designees have proposed to implement the following mitigation measures for marine mammals:

(1) Proposed exclusion zones around the sound source;

(2) Speed and course alterations;

(3) Shut-down procedures; and

(4) Ramp-up procedures.

Proposed Exclusion Zones—During pre-planning of the cruise, the smallest airgun array was identified that could be used and still meet the geophysical scientific objectives. NSF and ASC use radii to designate exclusion and buffer zones and to estimate take for marine mammals. Table 2 (presented earlier in

this document) shows the distances at which one would expect to receive three sound levels (160, 180, and 190 dB) from the two GI airgun array. The 180 and 190 dB level shut-down criteria are applicable to cetaceans and pinnipeds, respectively, as specified by NMFS (2000). NSF and ASC used these levels to establish the exclusion and buffer zones.

Received sound levels have been modeled by L-DEO for a number of airgun configurations, including two 45 in³ Nucleus G airguns, in relation to distance and direction from the airguns (see Figure 2 of the IHA application). In addition, propagation measurements of pulses from two GI airguns have been reported for shallow water (approximately 30 m [98.4 ft] depth in the GOM (Tolstoy *et al.*, 2004).

However, measurements were not made for the two GI airguns in deep water. The model does not allow for bottom interactions, and is most directly applicable to deep water. Based on the modeling, estimates of the maximum distances from the GI airguns where sound levels are predicted to be 190, 180, and 160 dB re 1 μ Pa (rms) in shallow, intermediate, and deep water were determined (see Table 2 above).

Empirical data concerning the 190, 180, and 160 dB (rms) distances were acquired for various airgun arrays based on measurements during the acoustic verification studies conducted by L-DEO in the northern GOM in 2003 (Tolstoy *et al.*, 2004) and 2007 to 2008 (Tolstoy *et al.*, 2009). Results of the 36 airgun array are not relevant for the two GI airguns to be used in the proposed survey. The empirical data for the 6, 10, 12, and 20 airgun arrays indicate that, for deep water, the L-DEO model tends to overestimate the received sound levels at a given distance (Tolstoy *et al.*, 2004). Measurements were not made for the two GI airgun array in deep water; however, NSF and ASC propose to use the safety radii predicted by L-DEO's model for the proposed GI airgun operations in shallow, intermediate, and deep water, although they are likely conservative given the empirical results for the other arrays.

Based on the modeling data, the outputs from the pair of 45 in³ or 105 in³ GI airguns proposed to be used during the seismic survey are considered a low-energy acoustic source in the NSF/USGS PEIS (2011) for marine seismic research. A low-energy seismic source was defined in the NSF/USGS PEIS as an acoustic source whose received level at 100 m is less than 180 dB. The NSF/USGS PEIS also established for these low-energy sources, a standard exclusion zone of

100 m for all low-energy sources in water depths greater than 100 m. This standard 100 m exclusion zone would be used during the proposed low-energy seismic survey. The 180 and 190 dB (rms) radii are shut-down criteria applicable to cetaceans and pinnipeds, respectively, as specified by NMFS (2000); these levels were used to establish exclusion zones. Therefore, the assumed 180 and 190 dB radii are 100 m for intermediate and deep water, respectively. If the PSO detects a marine mammal(s) within or about to enter the appropriate exclusion zone, the airguns will be shut-down immediately.

Speed and Course Alterations—If a marine mammal is detected outside the exclusion zone and, based on its position and direction of travel (relative motion), is likely to enter the exclusion zone, changes of the vessel's speed and/or direct course will be considered if this does not compromise operational safety or damage the deployed equipment. This would be done if operationally practicable while minimizing the effect on the planned science objectives. For marine seismic surveys towing large streamer arrays, however, course alterations are not typically implemented due to the vessel's limited maneuverability. After any such speed and/or course alteration is begun, the marine mammal activities and movements relative to the seismic vessel will be closely monitored to ensure that the marine mammal does not approach within the exclusion zone. If the marine mammal appears likely to enter the exclusion zone, further mitigation actions will be taken, including further speed and/or course alterations, and/or shut-down of the airgun(s). Typically, during seismic operations, the source vessel is unable to change speed or course, and one or more alternative mitigation measures will need to be implemented.

Shut-down Procedures—NSF and ASC will shut-down the operating airgun(s) if a marine mammal is detected outside the exclusion zone for the airgun(s), and if the vessel's speed and/or course cannot be changed to avoid having the animal enter the exclusion zone, the seismic source will be shut-down before the animal is within the exclusion zone. Likewise, if a marine mammal is already within the exclusion zone when first detected, the seismic source will be shut-down immediately.

Following a shut-down, NSF and ASC will not resume airgun activity until the marine mammal has cleared the exclusion zone. NSF and ASC will consider the animal to have cleared the exclusion zone if:

- A PSO has visually observed the animal leave the exclusion zone, or
- A PSO has not sighted the animal within the exclusion zone for 15 minutes for species with shorter dive durations (i.e., small odontocetes and pinnipeds), or 30 minutes for species with longer dive durations (i.e., mysticetes and large odontocetes, including sperm, pygmy and dwarf sperm, killer, and beaked whales).

Although power-down procedures are often standard operating practice for seismic surveys, they are not proposed to be used during this planned seismic survey because powering-down from two airguns to one airgun would make only a small difference in the exclusion zone(s)—but probably not enough to allow continued one-airgun operations if a marine mammal came within the exclusion zone for two airguns.

Ramp-up Procedures—Ramp-up of an airgun array provides a gradual increase in sound levels, and involves a step-wise increase in the number and total volume of airguns firing until the full volume of the airgun array is achieved. The purpose of a ramp-up is to “warn” marine mammals in the vicinity of the airguns and to provide the time for them to leave the area avoiding any potential injury or impairment of their hearing abilities. NSF and ASC will follow a ramp-up procedure when the airgun array begins operating after a specified period without airgun operations or when a shut-down shut down has exceeded that period. NSF and ASC propose that, for the present cruise, this period would be approximately 15 minutes. SIO, L-DEO, and USGS have used similar periods (approximately 15 minutes) during previous low-energy seismic surveys.

Ramp-up will begin with a single GI airgun (45 or 105 in³). The second GI airgun (45 or 105 in³) will be added after 5 minutes. During ramp-up, the PSOs will monitor the exclusion zone, and if marine mammals are sighted, a shut-down will be implemented as though both GI airguns were operational.

If the complete exclusion zone has not been visible for at least 30 minutes prior to the start of operations in either daylight or nighttime, NSF and ASC will not commence the ramp-up. Given these provisions, it is likely that the airgun array will not be ramped-up from a complete shut-down at night or in thick fog, because the outer part of the exclusion zone for that array will not be visible during those conditions. If one airgun has operated, ramp-up to full power will be permissible at night or in poor visibility, on the assumption that marine mammals will be alerted to the approaching seismic vessel by the

sounds from the single airgun and could move away if they choose. A ramp-up from a shut-down may occur at night, but only where the exclusion zone is small enough to be visible. NSF and ASC will not initiate a ramp-up of the airguns if a marine mammal is sighted within or near the applicable exclusion zones during the day or close to the vessel at night.

NMFS has carefully evaluated the applicant’s proposed mitigation measures and has considered a range of other measures in the context of ensuring that NMFS prescribes the means of effecting the least practicable adverse impact on the affected marine mammal species and stocks and their habitat. NMFS’s evaluation of potential measures included consideration of the following factors in relation to one another:

- (1) The manner in which, and the degree to which, the successful implementation of the measure is expected to minimize adverse impacts to marine mammals;
- (2) The proven or likely efficacy of the specific measure to minimize adverse impacts as planned; and
- (3) The practicability of the measure for applicant implementation.

Based on NMFS’s evaluation of the applicant’s proposed measures, as well as other measures considered by NMFS or recommended by the public, NMFS has preliminarily determined that the proposed mitigation measures provide the means of effecting the least practicable adverse impacts on marine mammal species or stocks and their habitat, paying particular attention to rookeries, mating grounds, and areas of similar significance.

Proposed Monitoring and Reporting

In order to issue an ITA for an activity, section 101(a)(5)(D) of the MMPA states that NMFS must set forth “requirements pertaining to the monitoring and reporting of such taking.” The MMPA implementing regulations at 50 CFR 216.104 (a)(13) indicate that requests for IHAs must include the suggested means of accomplishing the necessary monitoring and reporting that will result in increased knowledge of the species and of the level of taking or impacts on populations of marine mammals that are expected to be present in the action area.

Proposed Monitoring

NSF and ASC proposes to sponsor marine mammal monitoring during the proposed project, in order to implement the proposed mitigation measures that require real-time monitoring, and to

satisfy the anticipated monitoring requirements of the IHA. NSF and ASC’s proposed “Monitoring Plan” is described below this section. NSF and ASC understand that this monitoring plan will be subject to review by NMFS and that refinements may be required. The monitoring work described here has been planned as a self-contained project independent of any other related monitoring projects that may be occurring simultaneously in the same regions. NSF and ASC is prepared to discuss coordination of their monitoring program with any related work that might be done by other groups insofar as this is practical and desirable.

Vessel-Based Visual Monitoring

PSOs will be based aboard the seismic source vessel and will watch for marine mammals near the vessel during icebreaking activities, daytime airgun operations (austral summer) and during any ramp-ups of the airguns at night. Nighttime operations of the airguns are not anticipated. PSOs will also watch for marine mammals near the seismic vessel for at least 30 minutes prior to the start of airgun operations and after an extended shut-down (i.e., greater than approximately 15 minutes for this proposed low-energy seismic survey). When feasible, PSOs will conduct observations during daytime periods when the seismic system is not operating (such as during transits) for comparison of sighting rates and behavior with and without airgun operations and between acquisition periods. Based on PSO observations, the airguns will be shut-down when marine mammals are observed within or about to enter a designated exclusion zone. The exclusion zone is a region in which a possibility exists of adverse effects on animal hearing or other physical effects.

During seismic operations in the Dumont d’Urville Sea of the Southern Ocean, at least two PSOs will be based aboard the *Palmer*. At least one PSO will stand watch at all times while the *Palmer* is operating airguns during the proposed low-energy seismic survey; this procedure will also be followed when the vessel is conducting icebreaking during transit. NSF and ASC will appoint the PSOs with NMFS’s concurrence. The lead PSO would be experienced with marine mammal species in the Southern Ocean, the second PSO would receive additional specialized training from the PSO to ensure that they can identify marine mammal species commonly found in the Southern Ocean. Observations will take place during ongoing daytime operations and nighttime ramp-ups of the airguns.

During the majority of seismic operations, at least one PSO will be on duty from observation platforms (i.e., the best available vantage point on the source vessel) to monitor marine mammals near the seismic vessel. PSO(s) will be on duty in shifts no longer than 4 hours in duration. Other crew will also be instructed to assist in detecting marine mammals and implementing mitigation requirements (if practical). Before the start of the low-energy seismic survey, the crew will be given additional instruction on how to do so. (**Note:** because of the high latitude locations of the study areas, twilight/darkness conditions are expected to be limited to between 3 and 6 hours per day during the proposed action.)

The *Palmer* is a suitable platform for marine mammal observations and will serve as the platform from which PSOs will watch for marine mammals before and during seismic operations. Two locations are likely as observation stations onboard the *Palmer*. Observing stations are located on the bridge level, with the PSO eye level at approximately 16.5 m (54.1 ft) above the waterline and the PSO would have a good view around the entire vessel. In addition, there is an aloft observation tower for the PSO approximately 24.4 m (80.1 ft) above the waterline that is protected from the weather, and affords PSOs an even greater view. Standard equipment for PSOs will be reticle binoculars. Night-vision equipment will not be available or required due to the constant daylight conditions during the Antarctic summer. The PSOs will be in communication with ship's officers on the bridge and scientists in the vessel's operations laboratory, so they can advise promptly of the need for avoidance maneuvers or seismic source shut-down. Observing stations will be at the bridge level and the aloft observation tower. The approximate view around the vessel from the bridge is 270° and 360° from the aloft observation tower. During daytime, the PSO(s) will scan the area around the vessel systematically with reticle binoculars (e.g., 7 × 50 Fujinon FMTRC-SX) and the naked eye. These binoculars will have a built-in daylight compass. Estimating distances is done primarily with the reticles in the binoculars. The PSO(s) will be in direct (radio) wireless communication with ship's officers on the bridge and scientists in the vessel's operations laboratory during seismic operations, so they can advise the vessel operator, science support personnel, and the science party promptly of the need for avoidance maneuvers or a shut-

down of the seismic source. PSOs will monitor for the presence pinnipeds and cetaceans during icebreaking activities, and will be limited to those marine mammal species in proximity to the ice margin habitat. Observations within the buffer zone would also include pinnipeds that may be present on the surface of the sea ice (i.e., hauled-out) and that could potentially dive into the water as the vessel approaches, indicating disturbance from noise generated by icebreaking activities).

When marine mammals are detected within or about to enter the designated exclusion zone, the airguns will immediately be shut-down if necessary. The PSO(s) will continue to maintain watch to determine when the animal(s) are outside the exclusion zone by visual confirmation. Airgun operations will not resume until the animal is confirmed to have left the exclusion zone, or if not observed after 15 minutes for species with shorter dive durations (small odontocetes and pinnipeds) or 30 minutes for species with longer dive durations (mysticetes and large odontocetes, including sperm, killer, and beaked whales).

PSO Data and Documentation

PSOs will record data to estimate the numbers of marine mammals exposed to various received sound levels and to document apparent disturbance reactions or lack thereof. Data will be used to estimate numbers of animals potentially "taken" by harassment (as defined in the MMPA). They will also provide information needed to order a shut-down of the airguns when a marine mammal is within or near the exclusion zone. Observations will also be made during icebreaking activities as well as daytime periods when the *Palmer* is underway without seismic operations (i.e., transits, to, from, and through the study area) to collect baseline biological data.

When a sighting is made, the following information about the sighting will be recorded:

1. Species, group size, age/size/sex categories (if determinable), behavior when first sighted and after initial sighting, heading (if consistent), bearing and distance from seismic vessel, sighting cue, apparent reaction to the seismic source or vessel (e.g., none, avoidance, approach, paralleling, etc.), and behavioral pace.

2. Time, location, heading, speed, activity of the vessel, sea state, wind force, visibility, and sun glare.

The data listed under (2) will also be recorded at the start and end of each observation watch, and during a watch

whenever there is a change in one or more of the variables.

All observations, as well as information regarding ramp-ups or shut-downs will be recorded in a standardized format. Data will be entered into an electronic database. The data accuracy will be verified by computerized data validity checks as the data are entered and by subsequent manual checking of the database by the PSOs at sea. These procedures will allow initial summaries of data to be prepared during and shortly after the field program, and will facilitate transfer of the data to statistical, graphical, and other programs for further processing and archiving.

Results from the vessel-based observations will provide the following information:

1. The basis for real-time mitigation (airgun shut-down).
2. Information needed to estimate the number of marine mammals potentially taken by harassment, which must be reported to NMFS.
3. Data on the occurrence, distribution, and activities of marine mammals in the area where the seismic study is conducted.
4. Information to compare the distance and distribution of marine mammals relative to the source vessel at times with and without seismic activity.
5. Data on the behavior and movement patterns of marine mammals seen at times with and without seismic activity.

NSF and ASC will submit a comprehensive report to NMFS within 90 days after the end of the cruise. The report will describe the operations that were conducted and sightings of marine mammals near the operations. The report submitted to NMFS will provide full documentation of methods, results, and interpretation pertaining to all monitoring. The 90-day report will summarize the dates and locations of seismic operations and all marine mammal sightings (i.e., dates, times, locations, activities, and associated seismic survey activities). The report will minimally include:

- Summaries of monitoring effort—total hours, total distances, and distribution of marine mammals through the study period accounting for Beaufort sea state and other factors affecting visibility and detectability of marine mammals;
- Analyses of the effects of various factors influencing detectability of marine mammals including Beaufort sea state, number of PSOs, and fog/glare;
- Species composition, occurrence, and distribution of marine mammals sightings including date, water depth,

numbers, age/size/gender, and group sizes; and analyses of the effects of seismic operations;

- Sighting rates of marine mammals during periods with and without airgun activities (and other variables that could affect detectability);
- Initial sighting distances versus airgun activity state;
- Closest point of approach versus airgun activity state;
- Observed behaviors and types of movements versus airgun activity state;
- Numbers of sightings/individuals seen versus airgun activity state; and
- Distribution around the source vessel versus airgun activity state.

The report will also include estimates of the number and nature of exposures that could result in “takes” of marine mammals by harassment or in other ways. After the report is considered final, it will be publicly available on the NMFS Web site at: <http://www.nmfs.noaa.gov/pr/permits/incidental.htm#iha>.

In the unanticipated event that the specified activity clearly causes the take of a marine mammal in a manner prohibited by this IHA, such as an injury (Level A harassment), serious injury or mortality (e.g., ship-strike, gear interaction, and/or entanglement), NSF and ASC will immediately cease the specified activities and immediately report the incident to the Chief of the Permits and Conservation Division, Office of Protected Resources, NMFS at 301-427-8401 and/or by email to Jolie.Harrison@noaa.gov and Howard.Goldstein@noaa.gov. The report must include the following information:

- Time, date, and location (latitude/longitude) of the incident;
- Name and type of vessel involved;
- Vessel’s speed during and leading up to the incident;
- Description of the incident;
- Status of all sound source use in the 24 hours preceding the incident;
- Water depth;
- Environmental conditions (e.g., wind speed and direction, Beaufort sea state, cloud cover, and visibility);
- Description of all marine mammal observations in the 24 hours preceding the incident;
- Species identification or description of the animal(s) involved;
- Fate of the animal(s); and
- Photographs or video footage of the animal(s) (if equipment is available).

Activities shall not resume until NMFS is able to review the circumstances of the prohibited take. NMFS shall work with NSF and ASC to determine what is necessary to minimize the likelihood of further prohibited take and ensure MMPA

compliance. NSF and ASC may not resume their activities until notified by NMFS via letter or email, or telephone.

In the event that NSF and ASC discovers an injured or dead marine mammal, and the lead PSO determines that the cause of the injury or death is unknown and the death is relatively recent (i.e., in less than a moderate state of decomposition as described in the next paragraph), NSF and ASC will immediately report the incident to the Chief of the Permits and Conservation Division, Office of Protected Resources, NMFS, at 301-427-8401, and/or by email to Jolie.Harrison@noaa.gov and Howard.Goldstein@noaa.gov. The report must include the same information identified in the paragraph above. Activities may continue while NMFS reviews the circumstances of the incident. NMFS will work with NSF and ASC to determine whether modifications in the activities are appropriate.

In the event that NSF and ASC discovers an injured or dead marine mammal, and the lead PSO determines that the injury or death is not associated with or related to the activities authorized in the IHA (e.g., previously wounded animal, carcass with moderate or advanced decomposition, or scavenger damage), NSF and ASC will report the incident to the Chief of the Permits and Conservation Division, Office of Protected Resources, NMFS, at 301-427-8401, and/or by email to Jolie.Harrison@noaa.gov and Howard.Goldstein@noaa.gov, within 24 hours of discovery. NSF and ASC will provide photographs or video footage (if available) or other documentation of the stranded animal sighting to NMFS. Activities may continue while NMFS reviews the circumstances of the incident.

Estimated Take by Incidental Harassment

Except with respect to certain activities not pertinent here, the MMPA defines “harassment” as: any act of pursuit, torment, or annoyance which (i) has the potential to injure a marine mammal or marine mammal stock in the wild [Level A harassment]; or (ii) has the potential to disturb a marine mammal or marine mammal stock in the wild by causing disruption of behavioral patterns, including, but not limited to, migration, breathing, nursing, breeding, feeding, or sheltering [Level B harassment].

Level B harassment is anticipated and proposed to be authorized as a result of the proposed low-energy marine seismic survey in the Dumont d’Urville Sea off the coast of East Antarctica. Acoustic

stimuli (i.e., increased underwater sound) generated during the operation of the seismic airgun array and icebreaking activities are expected to result in the behavioral disturbance of some marine mammals. There is no evidence that the planned activities could result in injury, serious injury, or mortality for which NSF and ASC seeks the IHA. The required mitigation and monitoring measures will minimize any potential risk for injury, serious injury, or mortality.

The following sections describe NSF and ASC’s methods to estimate take by incidental harassment and present the applicant’s estimates of the numbers of marine mammals that could be affected during the proposed low-energy seismic survey in the Dumont d’Urville Sea off the coast of East Antarctica. The estimates are based on a consideration of the number of marine mammals that could be harassed by approximately 2,800 km (1511.9 nmi) of seismic operations with the two GI airgun array to be used and 1,000 km of icebreaking activities.

During simultaneous operations of the airgun array and the other sound sources, any marine mammals close enough to be affected by the single and multi-beam echosounders, pingers, ADCP, sub-bottom profiler, etc. would already be affected by the airguns. During times when the airguns are not operating, it is unlikely that marine mammals will exhibit more than minor, short-term responses to the echosounders, ADCPs, and sub-bottom profiler given their characteristics (e.g., narrow, downward-directed beam) and other considerations described previously. Therefore, for this activity, take was not authorized specifically for these sound sources beyond that which is already authorized for airguns and icebreaking activities.

There are no stock assessments and very limited population information available for marine mammals in the Dumont d’Urville Sea. Sighting data from the Australian Antarctic Division’s (AAD) BROKE-West surveys (Southwell *et al.*, 2008; 2012) was used to determine and estimate marine mammals densities for mysticetes and odontocetes and AAD data components for pinnipeds, which were not available for the proposed seismic survey’s action area in the Dumont d’Urville Sea. While population density data for cetaceans in the Southern Ocean is sparse to nonexistent, reported sightings data from previous research cruises suggest cetaceans such as those identified in Table 12 of the IHA application span a range greater than 4,000 km (2,159.8 nmi) off the coast of East Antarctica.

The AAD BROKE-West survey was not specifically designed to quantify marine mammals. The data was in terms of animals sighted per time unit, and this sighting data was then converted to an areal density by multiplying the number of animals observed by the estimated area observed during the survey. As such, some marine mammals that were present in the area may not have been observed.

The estimated number of cetaceans and pinnipeds that may be potentially exposed from the proposed seismic airgun operations and icebreaking activities based on sighting data from previous research cruises over a 52-day period and 13-day period. Some of the AAD sighting data was used as the basis for estimating take included "unidentified whale" species, this category was retained and pro-rated to the other species because environmental conditions may be present during the proposed action to limit identification of observed cetaceans. The estimated frequency of sightings data for cetaceans incorporates a correction factor of 5 that assumes only 20% of the animals present were reported due to sea ice and other conditions that may have hindered observation. The 20% factor was intended to conservatively account for this. Conservatively, a 40% correction factor was used for pinnipeds. The expected sightings data incorporates a 40% correction factor to account for seals that may be in the water versus those hauled-out on ice

surface. This correction factor for pinnipeds was conservatively based on Southwell *et al.* (2012), which estimated 20 to 40% of crabeater seals may be in the water in a particular area while the rest are hauled-out. The correction factor takes into consideration some pinnipeds may not be observed due to poor visibility conditions.

Sightings data were collected by the AAD; however, the AAD methodology was not described. Density is generally reported in the number of animals per km or square km. Estimated area observed by observers was calculated by using the average vessel speed (5.6 km/hr) times the estimated hours of the survey to estimate the total distance covered for each of the surveys. This was then converted from the linear distance into an area by assuming a width of 5 km that could be reliably visually surveyed. Therefore, the estimated area was 5,753 km² (1,677.3 nmi²) to obtain mysticete and odontocete densities and the estimated area was 1,419 km² (413.7 nmi²) to obtain pinniped densities.

Of the six species of pinnipeds that may be present in the study area during the proposed action, only four species are expected to be observed and occur mostly near pack ice or coastal areas and not prevalent in open sea areas where the low-energy seismic survey would be conducted. Because density estimates for pinnipeds in that Antarctic regions typically represent individuals that have hauled-out of the water, those

estimates are not representative of individuals that are in the water and could be potentially exposed to underwater sounds during the seismic airgun operations and icebreaking activities; therefore, the pinniped densities have been adjusted to account for this concern. Take was not requested for southern elephant seals and Antarctic fur seals because preferred habitat for these species is not within the proposed action area. Although no sightings of Weddell seals and spectacled porpoises were reported in the BROKE-West sighting data, take was requested for these species based on NMFS recommendation and IWC SOWER data. Although there is some uncertainty about the representatives of the data and the assumptions used in the calculations below, the approach used here is believed to be the best available approach.

Table 5. Estimated densities and possible number of marine mammal species that might be exposed to greater than or equal to 120 dB (icebreaking) and 160 dB (airgun operations) during NSF and ASC's proposed low-energy seismic survey (approximately 1,000 km of tracklines/approximately 3,500 km² ensonified area for icebreaking activities and approximately 2,800 km of tracklines/approximately 5,628 km² ensonified area for airgun operations) in the Dumont d'Urville Sea of the Southern Ocean, February to March 2014.

Species	Reported sightings ^{1,2} *sightings have been pro-rated to include unidentified animals*	Corrected sightings (assume 20% for cetaceans, 40% of pinnipeds in water)	Density in-water [density in-water and/or on-ice] (#/km) ²	Calculated take from seismic airgun operations (i.e., estimated number of individuals exposed to sound levels ≥160 dB re 1 μPa) ³	Calculated take from icebreaking activities (i.e., estimated number of individuals exposed to sound levels ≥120 dB re 1 μPa) ⁴	Approximate percentage of population estimate (calculated total take) ⁵	Total requested take authorization ⁶
Mysticetes:							
Southern right whale	0	0	0	0	0	NA	0.
Humpback whale	238	1,190	0.2068400	1,165	724	5.4	1,165 + 724 = 1,889.
Antarctic minke whale	136	680	0.1181943	666	414	0.4	666 + 414 = 1,080.
Sei whale	4	20	0.0034763	20	13	0.04	20 + 13 = 33.
Fin whale	232	1,160	0.2016255	1,135	706	1.3	1,135 + 706 = 1,841.
Blue whale	2	10	0.0017382	10	7	1.0	10 + 7 = 17.
Odontocetes:							
Sperm whale	32	160	0.0278104	157	98	2.7	157 + 98 = 255.
Arnoux's beaked whale	0	0	0	0	0	NA	0.
Cuvier's beaked whale	0	0	0	0	0	NA	0.
Southern bottlenose beaked whale	0	0	0	0	0	NA	0.
Killer whale	62	310	0.538827	304	189	2.0	304 + 189 = 493.
Long-finned pilot whale	24	120	0.0208578	118	74	0.1	118 + 74 = 192.
Hourglass dolphin	27	135	0.0234650	133	83	0.15	133 + 83 = 216.
Spectacled porpoise	26	130	0.0225690	128	80	NA	128 + 80 = 208.
Pinnipeds:							
Crabeater seal	2,220	888	0.625546 [2.189411]	3,521	7,663	0.2	3,521 + 7,663 = 11,184.
Leopard seal	17	7	0.00479 [0.016766]	27	59	0.04	27 + 59 = 86.
Ross seal	42	17	0.011835 [0.041421]	66	145	0.2	66 + 145 = 211.

Species	Reported sightings ^{1,2} *sightings have been pro-rated to include unidentified animals*	Corrected sightings (assume 20% for cetaceans, 40% of pinnipeds in water)	Density in-water [density in-water and/or on-ice] (#/km) ²	Calculated take from seismic airgun operations (i.e., estimated number of individuals exposed to sound levels ≥160 dB re 1 μPa) ³	Calculated take from icebreaking activities (i.e., estimated number of individuals exposed to sound levels ≥120 dB re 1 μPa) ⁴	Approximate percentage of population estimate (calculated total take) ⁵	Total requested take authorization ⁶
Weddell seal	302	121	0.054 [0.054]	303	189	0.1	303 + 189 = 492.
Southern elephant seal	0	0	0	0	0	NA	0.
Antarctic fur seal	0	0	0	0	0	NA	0.

NA = Not available or not assessed.

¹ Sightings from a 52 day (5,753 km²) period on the AAD BROKE-West survey during January to March 2006.

² Sightings December 3 to 16, 1999 (1,420 km² and 75,564 km²), below 60° South latitude between 110 to 165° East longitude. All sightings were animals hauled-out of the water and on the sea ice.

³ Calculated take is estimated density (reported density times correction factor) multiplied by the area ensonified to 160 dB (rms) around the planned seismic lines, increased by 25% for contingency.

⁴ Calculated take is estimated density (reported density) multiplied by the area ensonified to 120 dB (rms) around the planned transit lines where icebreaking activities may occur.

⁵ Total requested (and calculated) takes expressed as percentages of the species or regional populations.

⁶ Requested Take Authorization includes unidentified animals that were added to the observed and identified species on a pro-rated basis.

Note: Take was not requested for southern elephant seals and Antarctic fur seals because preferred habitat for these species is not within the proposed action area.

Icebreaking in Antarctic waters will occur, as necessary, between the latitudes of approximately 66 to 70° South and between 140 and 165° East. Based on a maximum sea ice extent of 250 km and estimating that the *Palmer* will transit to the innermost shelf and back into open water twice—a round trip transit in each of the potential work regions, it is estimated that the *Palmer* will actively break ice up to a distance of 1,000 km. Based on the ship’s speed of 5 kts under moderate ice conditions, this distance represents approximately 108 hrs of icebreaking operations. This calculation is likely an overestimation because icebreakers often follow leads when they are available and thus do not break ice at all times.

Numbers of marine mammals that might be present and potentially disturbed are estimated based on the available data about marine mammal distribution and densities in the Southern Ocean study are during the austral summer. NSF and ASC estimated the number of different individuals that may be exposed to airgun sounds with received levels greater than or equal to 160 dB re 1 μPa (rms) for seismic airgun operations and greater than or equal to 120 dB re 1 μPa (rms) for icebreaking activities on one or more occasions by considering the total marine area that would be within the 160 dB radius around the operating airgun array and 120 dB radius for the icebreaking activities on at least one occasion and the expected density of marine mammals in the area (in the absence of the a seismic survey and icebreaking activities). The number of possible exposures can be estimated by considering the total marine area that would be within the 160 dB radius (i.e.,

diameter is 1,005 m times 2) around the operating airguns. The ensonified area for icebreaking was estimated by multiplying the distance of the icebreaking activities (1,000 km) by the estimated diameter of the area within the 120 dB radius (i.e., diameter is 1,750 m times 2). The 160 dB radii are based on acoustic modeling data for the airguns that may be used during the proposed action (see Attachment B of the IHA application). As summarized in Table 2 (see Table 11 of the IHA application), the modeling results for the proposed low-energy seismic airgun array indicate the received levels are dependent on water depth. Since the majority of the proposed airgun operations would be conducted in waters 100 to 1,000 m deep, the buffer zone of 1,005 m used for the two 105 in³ GI airguns was used to be more conservative. The expected sighting data for pinnipeds accounts for both pinnipeds that may be in the water and those hauled-out on ice surfaces. While the number of cetaceans that may be encountered within the ice margin habitat would be expected to be less than open water, the estimates utilized expected sightings for the open water and represent conservative estimates. It is unlikely that a particular animal would stay in the area during the entire survey.

The number of different individuals potentially exposed to received levels greater than or equal to 160 dB re 1 μPa (rms) from seismic airgun operations and 120 dB re 1 μPa (rms) for icebreaking activities was calculated by multiplying:

(1) The expected species density (in number/km²), times.

(2) The anticipated area to be ensonified to that level during airgun operations.

Applying the approach described above, approximately 5,628 km² (including the 25% contingency) would be ensonified within the 160 dB isopleth for seismic airgun operations and approximately 3,500 km² would be ensonified within the 120 dB isopleth for icebreaking activities on one or more occasions during the proposed survey. The take calculations within the study sites do not explicitly add animals to account for the fact that new animals (i.e., turnover) are not accounted for in the initial density snapshot and animals could also approach and enter the area ensonified above 160 dB for seismic airgun operations and 120 dB for icebreaking activities; however, studies suggest that many marine mammals will avoid exposing themselves to sounds at this level, which suggests that there would not necessarily be a large number of new animals entering the area once the seismic survey and icebreaking activities started. Because this approach for calculating take estimates does not allow for turnover in the marine mammal populations in the area during the course of the survey, the actual number of individuals exposed may be underestimated, although the conservative (i.e., probably overestimated) line-kilometer distances used to calculate the area may offset this. Also, the approach assumes that no cetaceans or pinnipeds will move away or toward the tracklines as the *Palmer* approaches in response to increasing sound levels before the levels reach 160 dB for seismic airgun operations and 120 dB for icebreaking activities. Another way of interpreting the

estimates that follow is that they represent the number of individuals that are expected (in absence of a seismic and icebreaking program) to occur in the waters that will be exposed to greater than or equal to 160 dB (rms) for seismic airgun operations and greater than or equal to 120 dB (rms) for icebreaking activities.

NSF and ASC's estimates of exposures to various sound levels assume that the proposed surveys will be carried out in full; however, the ensonified areas calculated using the planned number of line-kilometers has been increased by 25% to accommodate lines that may need to be repeated, equipment testing, etc. As is typical during offshore ship surveys, inclement weather and equipment malfunctions are likely to cause delays and may limit the number of useful line-kilometers of seismic operations that can be undertaken. The estimates of the numbers of marine mammals potentially exposed to 120 dB (rms) and 160 dB (rms) received levels are precautionary and probably overestimate the actual numbers of marine mammals that could be involved. These estimates assume that there will be no weather, equipment, or mitigation delays, which is highly unlikely.

Table 5 shows the estimates of the number of different individual marine mammals anticipated to be exposed to greater than or equal to 120 dB re 1 μ Pa (rms) for icebreaking activities and greater than or equal to 160 dB re 1 μ Pa (rms) for seismic airgun operations during the seismic survey if no animals moved away from the survey vessel. The total requested take authorization is given in the far right column of Table 5.

The estimate of the number of individual cetaceans and pinnipeds that could be exposed to seismic sounds with received levels greater than or equal to 160 dB re 1 μ Pa (rms) and sounds from icebreaking activities with received levels greater than or equal to 120 dB re 1 μ Pa (rms) during the proposed survey is (with 25% contingency) in Table 5 of this document. That total (with 25% contingency) includes 1,889 humpback, 1,080 Antarctic minke, 33 sei, 1,841 fin, 17 blue, and 255 sperm whales could be taken by Level B harassment during the proposed seismic survey, which would represent 5.4, 0.4, 0.04, 1.3, 1, and 2.7% of the worldwide or regional populations, respectively. Some of the cetaceans potentially taken by Level B harassment are delphinids and porpoises: killer whales, long-finned pilot whales, hourglass dolphins, and spectacled porpoises are estimated to be

the most common delphinid and porpoise species in the area, with estimates of 493, 192, 216, and 208, which would represent 2, 0.1, and 0.15% (spectacled porpoise population is not available) of the affected worldwide or regional populations, respectively. Most of the pinnipeds potentially taken by Level B harassment are: Crabeater, leopard, Ross, and Weddell seals with estimates of 11,184, 86, 211, and 492, which would represent 0.2, 0.04, 0.2, and 0.1% of the affected worldwide or regional populations, respectively.

Encouraging and Coordinating Research

NSF and ASC will coordinate the planned marine mammal monitoring program associated with the proposed low-energy seismic survey with other parties that express interest in this activity and area. NSF and ASC will coordinate with applicable U.S. agencies (e.g., NMFS), and will comply with their requirements. NSF has already reached out to the Australian Antarctic Division (AAD), who are the proponents of the proposed marine protected area and regularly conduct research expeditions in the marine environment off East Antarctica.

The proposed action would complement fieldwork studying other Antarctic ice shelves, oceanographic studies, and ongoing development of ice sheet and other ocean models. It would facilitate learning at sea and ashore by students, help to fill important spatial and temporal gaps in a lightly sampled region of coastal Antarctica, provide additional data on marine mammals present in the East Antarctic study areas, and communicate its findings via reports, publications and public outreach.

Impact on Availability of Affected Species or Stock for Taking for Subsistence Uses

Section 101(a)(5)(D) of the MMPA also requires NMFS to determine that the authorization will not have an unmitigable adverse effect on the availability of marine mammal species or stocks for subsistence use. There are no relevant subsistence uses of marine mammals in the study area (in the Dumont d'Urville Sea off the coast of East Antarctica) that implicate MMPA section 101(a)(5)(D).

Negligible Impact and Small Numbers Analysis Determination

As a preliminary matter, NMFS typically includes our negligible impact and small numbers analyses and determinations under the same section

heading of our **Federal Register** notices. Despite co-locating these terms, NMFS acknowledges that negligible impact and small numbers are distinct standards under the MMPA and treat them as such. The analyses presented below do not conflate the two standards; instead, each standard has been considered independently and NMFS has applied the relevant factors to inform our negligible impact and small numbers determinations.

NMFS has defined "negligible impact" in 50 CFR 216.103 as "an impact resulting from the specified activity that cannot be reasonably expected to, and is not reasonably likely to, adversely affect the species or stock through effects on annual rates of recruitment or survival." In making a negligible impact determination, NMFS evaluated factors such as:

- (1) The number of anticipated injuries, serious injuries, or mortalities;
- (2) The number, nature, and intensity, and duration of Level B harassment (all relatively limited); and
- (3) The context in which the takes occur (i.e., impacts to areas of significance, impacts to local populations, and cumulative impacts when taking into account successive/contemporaneous actions when added to baseline data);
- (4) The status of stock or species of marine mammals (i.e., depleted, not depleted, decreasing, increasing, stable, impact relative to the size of the population);
- (5) Impacts on habitat affecting rates of recruitment/survival; and
- (6) The effectiveness of monitoring and mitigation measures.

As described above and based on the following factors, the specified activities associated with the marine seismic survey are not likely to cause PTS, or other non-auditory injury, serious injury, or death. The factors include:

- (1) The likelihood that, given sufficient notice through relatively slow ship speed, marine mammals are expected to move away from a noise source that is annoying prior to its becoming potentially injurious;
- (2) The potential for temporary or permanent hearing impairment is relatively low and would likely be avoided through the implementation of the shut-down measures;

No injuries, serious injuries, or mortalities are anticipated to occur as a result of the NSF and ASC's planned low-energy marine seismic survey, and none are proposed to be authorized by NMFS. Table 5 of this document outlines the number of requested Level B harassment takes that are anticipated as a result of these activities. Due to the

nature, degree, and context of Level B (behavioral) harassment anticipated and described (see “Potential Effects on Marine Mammals” section above) in this notice, the activity is not expected to impact rates of annual recruitment or survival for any affected species or stock, particularly given NMFS’s and the applicant’s proposal to implement mitigation, monitoring, and reporting measures to minimize impacts to marine mammals. Additionally, the seismic survey will not adversely impact marine mammal habitat.

For the other marine mammal species that may occur within the proposed action area, there are no known designated or important feeding and/or reproductive areas. Many animals perform vital functions, such as feeding, resting, traveling, and socializing, on a diel cycle (i.e., 24 hr cycle). Behavioral reactions to noise exposure (such as disruption of critical life functions, displacement, or avoidance of important habitat) are more likely to be significant if they last more than one diel cycle or recur on subsequent days (Southall *et al.*, 2007). Additionally, the seismic survey will be increasing sound levels in the marine environment in a relatively small area surrounding the vessel (compared to the range of the animals), which is constantly travelling over distances, and some animals may only be exposed to and harassed by sound for less than day.

Of the 14 marine mammal species under NMFS jurisdiction that may or are known to likely to occur in the study area, five are listed as threatened or endangered under the ESA: southern right, humpback, sei, fin, blue, and sperm whales. These species are also considered depleted under the MMPA. Of these ESA-listed species, incidental take has been requested to be authorized for humpback, sei, fin, blue, and sperm whales. There is generally insufficient data to determine population trends for the other depleted species in the study area. To protect these animals (and other marine mammals in the study area), NSF and ASC must cease or reduce airgun operations if any marine mammal enters designated zones. No injury, serious injury, or mortality is expected to occur and due to the nature, degree, and context of the Level B harassment anticipated, and the activity is not expected to impact rates of recruitment or survival.

As mentioned previously, NMFS estimates that 14 species of marine mammals under its jurisdiction could be potentially affected by Level B harassment over the course of the IHA. The population estimates for the marine mammal species that may be taken by

Level B harassment were provided in Table 4 of this document.

NMFS’s practice has been to apply the 160 dB re 1 μ Pa (rms) received level threshold for underwater impulse sound levels and the 120 dB re 1 μ Pa (rms) received level threshold for icebreaking activities to determine whether take by Level B harassment occurs. Southall *et al.* (2007) provide a severity scale for ranking observed behavioral responses of both free-ranging marine mammals and laboratory subjects to various types of anthropogenic sound (see Table 4 in Southall *et al.* [2007]).

NMFS has preliminarily determined, provided that the aforementioned mitigation and monitoring measures are implemented, the impact of conducting a low-energy marine seismic survey in the Dumont d’Urville Sea off the coast of East Antarctica, February to March 2014, may result, at worst, in a modification in behavior and/or low-level physiological effects (Level B harassment) of certain species of marine mammals.

While behavioral modifications, including temporarily vacating the area during the operation of the airgun(s), may be made by these species to avoid the resultant acoustic disturbance, the availability of alternate areas within these areas for species and the short and sporadic duration of the research activities, have led NMFS to preliminarily determine that the taking by Level B harassment from the specified activity will have a negligible impact on the affected species in the specified geographic region. NMFS believes that the length of the seismic survey, the requirement to implement mitigation measures (e.g., shut-down of seismic operations), and the inclusion of the monitoring and reporting measures, will reduce the amount and severity of the potential impacts from the activity to the degree that it will have a negligible impact on the species or stocks in the action area.

NMFS has preliminary determined, provided that the aforementioned mitigation and monitoring measures are implemented, that the impact of conducting a low-energy marine seismic survey in the Dumont d’Urville Sea off the coast of East Antarctica, January to March 2014, may result, at worst, in a temporary modification in behavior and/or low-level physiological effects (Level B harassment) of small numbers of certain species of marine mammals. See Table 5 for the requested authorized take numbers of marine mammals.

Endangered Species Act

Of the species of marine mammals that may occur in the proposed survey

area, several are listed as endangered under the ESA, including the humpback, sei, fin, blue, and sperm whales. NSF and ASC did not request take of endangered Southern right whales due to the low likelihood of encountering this species during the cruise. Under section 7 of the ESA, NSF, on behalf of ASC and five other research institutions, has initiated formal consultation with the NMFS, Office of Protected Resources, Endangered Species Act Interagency Cooperation Division, on this proposed seismic survey. NMFS’s Office of Protected Resources, Permits and Conservation Division, has initiated formal consultation under section 7 of the ESA with NMFS’s Office of Protected Resources, Endangered Species Act Interagency Cooperation Division, to obtain a Biological Opinion evaluating the effects of issuing the IHA on threatened and endangered marine mammals and, if appropriate, authorizing incidental take. NMFS will conclude formal section 7 consultation prior to making a determination on whether or not to issue the IHA. If the IHA is issued, NSF and ASC, in addition to the mitigation and monitoring requirements included in the IHA, will be required to comply with the Terms and Conditions of the Incidental Take Statement corresponding to NMFS’s Biological Opinion issued to both NSF and ASC, and NMFS’s Office of Protected Resources.

National Environmental Policy Act

With NSF and ASC’s complete application, NSF and ASC provided NMFS a “Draft Initial Environmental Evaluation/Environmental Assessment to Conduct Marine-Based Studies of the Totten Glacier System and Marine Record of Cryosphere—Ocean Dynamics,” (IEE/EA) prepared by AECOM on behalf of NSF and ASC. The IEE/EA analyzes the direct, indirect, and cumulative environmental impacts of the proposed specified activities on marine mammals including those listed as threatened or endangered under the ESA. Prior to making a final decision on the IHA application, NMFS will either prepare an independent EA, or, after review and evaluation of the NSF and ASC IEE/EA for consistency with the regulations published by the Council of Environmental Quality (CEQ) and NOAA Administrative Order 216–6, Environmental Review Procedures for Implementing the National Environmental Policy Act, adopt the NSF and ASC IEE/EA and make a decision of whether or not to issue a

Finding of No Significant Impact (FONSI).

Proposed Authorization

As a result of these preliminary determinations, NMFS propose to issue an IHA to NSF and ASC for conducting the low-energy seismic survey in the tropical western Pacific Ocean, provided the previously mentioned mitigation, monitoring, and reporting requirements are incorporated. The proposed IHA language is provided below:

National Science Foundation, Division of Polar Programs, 4201 Wilson Boulevard, Arlington, Virginia 22230 and Antarctic Support Contract, 7400 South Tucson Way, Centennial, Colorado 80112, is hereby authorized under section 101(a)(5)(D) of the Marine Mammal Protection Act (MMPA) (16 U.S.C. 1371(a)(5)(D)), to harass small numbers of marine mammals incidental to a low-energy marine geophysical (seismic) survey conducted by the RVIB *Nathaniel B. Palmer* (*Palmer*) in the Dumont d'Urville Sea, Antarctica, January to March 2014:

1. This Authorization is valid from January 29 through April 27, 2014.

2. This Authorization is valid only for the *Palmer's* activities associated with low-energy seismic survey operations that shall occur in the following specified geographic area:

In selected regions of the Dumont d'Urville Sea in the Southern Ocean off the coast of East Antarctica and focus on the Totten Glacier and Moscow University Ice Shelf, located on the Sabrina Coast, from greater than approximately 64° South and between approximately 95 to 135° East, and the Mertz Glacier and Cook Ice Shelf systems located on the George V and Oates Coast, from greater than approximately 65° South and between approximately 140 to 165° East. The study sites are characterized by heavy ice cover, with a seasonal break-up in the ice that structures biological patterns. The studies may occur in both areas, or entirely in one or the other, depending on ice conditions. Water depths in the survey area generally range from approximately 100 to 1,000 m, and possibly exceeding 1,000 m in some areas. The low-energy seismic survey will be conducted in International Waters (i.e., high seas), as specified in NSF and ASC's Incidental Harassment Authorization application and the associated NSF and ASC Initial Environmental Evaluation/ Environmental Assessment (IEE/EA).

3. Species Authorized and Level of Takes

(a) The incidental taking of marine mammals, by Level B harassment only, is limited to the following species in the waters of the Southern Ocean off the coast of East Antarctica:

(i) *Mysticetes*—see Table 2 (attached) for authorized species and take numbers.

(ii) *Odontocetes*—see Table 2 (attached) for authorized species and take numbers.

(iii) *Pinnipeds*—see Table 2 (attached) for authorized species and take numbers.

(iv) If any marine mammal species are encountered during seismic activities that are not listed in Table 2 (attached) for authorized taking and are likely to be exposed to sound pressure levels (SPLs) greater than or equal to 160 dB re 1 μ Pa (rms) for seismic airgun operations or greater than or equal to 120 dB re 1 μ Pa (rms) for icebreaking activities, then the Holder of this Authorization must alter speed or course or shut-down the airguns to avoid take.

(b) The taking by injury (Level A harassment), serious injury, or death of any of the species listed in Condition 3(a) above or the taking of any kind of any other species of marine mammal is prohibited and may result in the modification, suspension or revocation of this Authorization.

4. The methods authorized for taking by Level B harassment are limited to the following acoustic sources without an amendment to this Authorization:

(a) A two Generator Injector (GI) airgun array (each with a discharge volume of 45 cubic inches [in³] or 105 in³) with a total volume of 90 in³ or 210 in³ (or smaller);

(b) A multi-beam echosounder;

(c) A single-beam echosounder;

(d) An acoustic Doppler current profiler;

(e) An acoustic locator;

(f) A sub-bottom profiler; and

(g) Icebreaking.

5. The taking of any marine mammal in a manner prohibited under this Authorization must be reported immediately to the Office of Protected Resources, National Marine Fisheries Service (NMFS), at 301-427-8401.

6. Mitigation and Monitoring Requirements

The Holder of this Authorization is required to implement the following mitigation and monitoring requirements when conducting the specified activities to achieve the least practicable adverse impact on affected marine mammal species or stocks:

(a) Utilize one, NMFS-qualified, vessel-based Protected Species Observer (PSO) to visually watch for and monitor marine mammals near the seismic source vessel during daytime airgun operations (from nautical twilight-dawn to nautical twilight-dusk) and before and during ramp-ups of airguns day or night. The *Palmer's* vessel crew shall also assist in detecting marine mammals, when practicable. PSOs shall have access to reticle binoculars (7 × 50 Fujinon). PSO shifts shall last no longer than 4 hours at a time. PSOs shall also make observations during daytime periods when the seismic airguns are not operating for comparison of animal abundance and behavior, when feasible.

(b) PSOs shall conduct monitoring while the airgun array and streamer are being deployed or recovered from the water.

(c) Record the following information when a marine mammal is sighted:

(i) Species, group size, age/size/sex categories (if determinable), behavior when first sighted and after initial sighting, heading (if consistent), bearing and distance from seismic vessel, sighting cue, apparent reaction to the airguns or vessel (e.g., none, avoidance, approach, paralleling, etc., and including responses to ramp-up), and behavioral pace; and

(ii) Time, location, heading, speed, activity of the vessel (including number of airguns operating and whether in state of ramp-up or shut-down), Beaufort sea state and wind force, visibility, and sun glare; and

(iii) The data listed under Condition 6(c)(ii) shall also be recorded at the start and end of each observation watch and during a watch whenever there is a change in one or more of the variables.

(d) Visually observe the entire extent of the exclusion zone (180 dB re 1 μ Pa [rms] for cetaceans and 190 dB re 1 μ Pa [rms] for pinnipeds; see Table 2 [above] for distances) using NMFS-qualified PSOs, for at least 30 minutes prior to starting the airgun array (day or night). If the PSO finds a marine mammal within the exclusion zone, NSF and ASC must delay the seismic survey until the marine mammal(s) has left the area. If the PSO sees a marine mammal that surfaces, then dives below the surface, the PSO shall wait 30 minutes. If the PSO sees no marine mammals during that time, they should assume that the animal has moved beyond the exclusion zone. If for any reason the entire radius cannot be seen for the entire 30 minutes (i.e., rough seas, fog, darkness), or if marine mammals are near, approaching, or in the exclusion zone, the airguns may not be ramped-up. If one airgun is already running at a source level of at

least 180 dB re 1 μ Pa (rms), NSF and ASC may start the second airgun without observing the entire exclusion zone for 30 minutes prior, provided no marine mammals are known to be near the exclusion zone (in accordance with Condition 6[f] below).

(e) Establish a 180 dB re 1 μ Pa (rms) exclusion zone for cetaceans and a 190 dB re 1 μ Pa (rms) exclusion zone for pinnipeds before the two GI airgun array (90 or 210 in³ total volume) is in operation. See Table 2 (above) for distances and exclusion zones.

(f) Implement a "ramp-up" procedure when starting up at the beginning of seismic operations or anytime after the entire array has been shut-down for more than 15 minutes, which means starting with a single GI airgun and adding a second GI airgun after five minutes. During ramp-up, the PSOs shall monitor the exclusion zone, and if marine mammals are sighted, a shut-down shall be implemented as though the full array (both GI airguns) were operational. Therefore, initiation of ramp-up procedures from shut-down requires that the PSOs be able to view the full exclusion zone as described in Condition 6(d) (above).

(g) Alter speed or course during seismic operations if a marine mammal, based on its position and relative motion, appears likely to enter the relevant exclusion zone. If speed or course alteration is not safe or practicable, or if after alteration the marine mammal still appears likely to enter the exclusion zone, further mitigation measures, such as a shut-down, shall be taken.

(h) Shut-down the airgun(s) if a marine mammal is detected within, approaches, or enters the relevant exclusion zone (as defined in Table 2, above). A shut-down means all operating airguns are shut-down (i.e., turned off).

(i) Following a shut-down, the airgun activity shall not resume until the PSO has visually observed the marine mammal(s) exiting the exclusion zone and is not likely to return, or has not been seen within the exclusion zone for 15 minutes for species with shorter dive durations (small odontocetes and pinnipeds) or 30 minutes for species with longer dive durations (mysticetes and large odontocetes, including sperm, killer, and beaked whales).

(j) Following a shut-down and subsequent animal departure, airgun operations may resume following ramp-up procedures described in Condition 6(f).

(k) Marine seismic surveys may continue into night and low-light hours if such segment(s) of the survey is

initiated when the entire relevant exclusion zones are visible and can be effectively monitored.

(l) No initiation of airgun array operations is permitted from a shut-down position at night or during low-light hours (such as in dense fog or heavy rain) when the entire relevant exclusion zone cannot be effectively monitored by the PSO(s) on duty.

7. Reporting Requirements

The Holder of this Authorization is required to:

(a) Submit a draft report on all activities and monitoring results to the Office of Protected Resources, NMFS, within 90 days of the completion of the *Palmer's Dumont d'Urville Sea* off the coast of East Antarctica cruise. This report must contain and summarize the following information:

(i) Dates, times, locations, heading, speed, weather, sea conditions (including Beaufort sea state and wind force), and associated activities during all seismic operations and marine mammal sightings;

(ii) Species, number, location, distance from the vessel, and behavior of any marine mammals, as well as associated seismic activity (e.g., number of shut-downs), observed throughout all monitoring activities.

(iii) An estimate of the number (by species) of marine mammals that: (A) Are known to have been exposed to the seismic activity (based on visual observation) at received levels greater than or equal to 120 dB re 1 μ Pa (rms) (for icebreaking activities), greater than or equal to 160 dB re 1 μ Pa (rms) (for seismic airgun operations), and/or 180 dB re 1 μ Pa (rms) for cetaceans and 190 dB re 1 μ Pa (rms) for pinnipeds with a discussion of any specific behaviors those individuals exhibited; and (B) may have been exposed (based on modeled values for the two GI airgun array) to the seismic activity at received levels greater than or equal to 120 dB re 1 μ Pa (rms) (for icebreaking activities), greater than or equal to 160 dB re 1 μ Pa (rms) (for seismic airgun operations), and/or 180 dB re 1 μ Pa (rms) for cetaceans and 190 dB re 1 μ Pa (rms) for pinnipeds with a discussion of the nature of the probable consequences of that exposure on the individuals that have been exposed.

(iv) A description of the implementation and effectiveness of the: (A) Terms and Conditions of the Biological Opinion's Incidental Take Statement (ITS) (attached); and (B) mitigation measures of the Incidental Harassment Authorization. For the Biological Opinion, the report shall confirm the implementation of each

Term and Condition, as well as any conservation recommendations, and describe their effectiveness, for minimizing the adverse effects of the action on Endangered Species Act-listed marine mammals.

(b) Submit a final report to the Chief, Permits and Conservation Division, Office of Protected Resources, NMFS, within 30 days after receiving comments from NMFS on the draft report. If NMFS decides that the draft report needs no comments, the draft report shall be considered to be the final report.

8. In the unanticipated event that the specified activity clearly causes the take of a marine mammal in a manner prohibited by this Authorization, such as an injury (Level A harassment), serious injury or mortality (e.g., ship-strike, gear interaction, and/or entanglement), NSF and ASC shall immediately cease the specified activities and immediately report the incident to the Chief of the Permits and Conservation Division, Office of Protected Resources, NMFS, at 301-427-8401 and/or by email to Jolie.Harrison@noaa.gov and Howard.Goldstein@noaa.gov. The report must include the following information:

(a) Time, date, and location (latitude/longitude) of the incident; the name and type of vessel involved; the vessel's speed during and leading up to the incident; description of the incident; status of all sound source use in the 24 hours preceding the incident; water depth; environmental conditions (e.g., wind speed and direction, Beaufort sea state, cloud cover, and visibility); description of marine mammal observations in the 24 hours preceding the incident; species identification or description of the animal(s) involved; the fate of the animal(s); and photographs or video footage of the animal (if equipment is available).

Activities shall not resume until NMFS is able to review the circumstances of the prohibited take. NMFS shall work with NSF and ASC to determine what is necessary to minimize the likelihood of further prohibited take and ensure MMPA compliance. NSF and ASC may not resume their activities until notified by NMFS via letter, email, or telephone.

In the event that NSF and ASC discovers an injured or dead marine mammal, and the lead PSO determines that the cause of the injury or death is unknown and the death is relatively recent (i.e., in less than a moderate state of decomposition as described in the next paragraph), NSF and ASC will immediately report the incident to the Chief of the Permits and Conservation Division, Office of Protected Resources,

NMFS, at 301-427-8401, and/or by email to Jolie.Harrison@noaa.gov and Howard.Goldstein@noaa.gov. The report must include the same information identified in Condition 8(a) above. Activities may continue while NMFS reviews the circumstances of the incident. NMFS will work with NSF and ASC to determine whether modifications in the activities are appropriate.

In the event that NSF and ASC discovers an injured or dead marine mammal, and the lead PSO determines that the injury or death is not associated with or related to the activities authorized in Condition 2 of this Authorization (e.g., previously wounded animal, carcass with moderate to advanced decomposition, or scavenger damage), NSF and ASC shall report the incident to the Chief of the Permits and

Conservation Division, Office of Protected Resources, NMFS, at 301-427-8401, and/or by email to Jolie.Harrison@noaa.gov and Howard.Goldstein@noaa.gov, within 24 hours of the discovery. NSF and ASC shall provide photographs or video footage (if available) or other documentation of the stranded animal sighting to NMFS and the Marine Mammal Stranding Network. Activities may continue while NMFS reviews the circumstances of the incident.

9. NSF and ASC is required to comply with the Terms and Conditions of the ITS corresponding to NMFS's Biological Opinion issued to both NSF, ASC, and NMFS's Office of Protected Resources (attached).

10. A copy of this Authorization and the ITS must be in the possession of all contractors and PSOs operating under

the authority of this Incidental Harassment Authorization.

Information Solicited

NMFS requests interested persons to submit comments and information concerning this proposed project and NMFS's preliminary determination of issuing an IHA (see **ADDRESSES**). Concurrent with the publication of this notice in the **Federal Register**, NMFS is forwarding copies of this application to the Marine Mammal Commission and its Committee of Scientific Advisors.

Dated: December 30, 2013.

P. Michael Payne,

*Chief, Permits and Conservation Division,
Office of Protected Resources, National
Marine Fisheries Service.*

[FR Doc. 2013-31471 Filed 12-31-13; 8:45 am]

BILLING CODE 3510-22-P