

A11102 146412

NAT'L INST OF STANDARDS & TECH R.I.C.



A11102146412

Ross, Alberta B/Rate constants for react
QC100 .U573 V65:1979 C.2 NBS-PUB-C 1979



NSRDS-NBS 65

U.S. DEPARTMENT OF COMMERCE / National Bureau of Standards



Rate Constants for Reactions of Inorganic Radicals in Aqueous Solution

QC
100
.U573
NO.65
1979
C.2

NATIONAL BUREAU OF STANDARDS

The National Bureau of Standards¹ was established by an act of Congress March 3, 1901. The Bureau's overall goal is to strengthen and advance the Nation's science and technology and facilitate their effective application for public benefit. To this end, the Bureau conducts research and provides: (1) a basis for the Nation's physical measurement system, (2) scientific and technological services for industry and government, (3) a technical basis for equity in trade, and (4) technical services to promote public safety. The Bureau's technical work is performed by the National Measurement Laboratory, the National Engineering Laboratory, and the Institute for Computer Sciences and Technology.

THE NATIONAL MEASUREMENT LABORATORY provides the national system of physical and chemical and materials measurement; coordinates the system with measurement systems of other nations and furnishes essential services leading to accurate and uniform physical and chemical measurement throughout the Nation's scientific community, industry, and commerce; conducts materials research leading to improved methods of measurement, standards, and data on the properties of materials needed by industry, commerce, educational institutions, and Government; provides advisory and research services to other Government Agencies; develops, produces, and distributes Standard Reference Materials; and provides calibration services. The Laboratory consists of the following centers:

Absolute Physical Quantities² — Radiation Research — Thermodynamics and Molecular Science — Analytical Chemistry — Materials Science.

THE NATIONAL ENGINEERING LABORATORY provides technology and technical services to users in the public and private sectors to address national needs and to solve national problems in the public interest; conducts research in engineering and applied science in support of objectives in these efforts; builds and maintains competence in the necessary disciplines required to carry out this research and technical service; develops engineering data and measurement capabilities; provides engineering measurement traceability services; develops test methods and proposes engineering standards and code changes; develops and proposes new engineering practices; and develops and improves mechanisms to transfer results of its research to the ultimate user. The Laboratory consists of the following centers:

Applied Mathematics — Electronics and Electrical Engineering² — Mechanical Engineering and Process Technology² — Building Technology — Fire Research — Consumer Product Technology — Field Methods.

THE INSTITUTE FOR COMPUTER SCIENCES AND TECHNOLOGY conducts research and provides scientific and technical services to aid Federal Agencies in the selection, acquisition, application, and use of computer technology to improve effectiveness and economy in Government operations in accordance with Public Law 89-306 (40 U.S.C. 759), relevant Executive Orders, and other directives; carries out this mission by managing the Federal Information Processing Standards Program, developing Federal ADP standards guidelines, and managing Federal participation in ADP voluntary standardization activities; provides scientific and technological advisory services and assistance to Federal Agencies; and provides the technical foundation for computer-related policies of the Federal Government. The Institute consists of the following divisions:

Systems and Software — Computer Systems Engineering — Information Technology.

¹Headquarters and Laboratories at Gaithersburg, Maryland, unless otherwise noted; mailing address Washington, D.C. 20234.

²Some divisions within the center are located at Boulder, Colorado, 80303.

The National Bureau of Standards was reorganized, effective April 9, 1978.

JUN 19 1979

not for sale

150100

U573

110 65

1979

5.2

Rate Constants for Reactions of Inorganic Radicals in Aqueous Solution

Alberta B. Ross and Pedatsur Neta

Radiation Chemistry Data Center
Radiation Laboratory
University of Notre Dame
Notre Dame, IN 46556



National Standard Reference Data Series #145

U.S. DEPARTMENT OF COMMERCE, Juanita M. Kreps, Secretary

Jordan J. Baruch, Assistant Secretary for Science and Technology

NATIONAL BUREAU OF STANDARDS, Ernest Ambler, Director

Issued June 1979

Library of Congress Cataloging in Publication Data

Ross, Alberta B

Rate constants for reactions of inorganic radicals in aqueous solution.

(Nat. stand. ref. data ser. ; NSRDS-NBS 65)

"National Standard Reference Data System."

Supt. of Docs. no.: C13.48:65

1. Chemical reaction, Rate of--Tables. 2. Radicals (Chemistry)--Tables.

I. Neta, Pedatsur, joint author. II. Title. III. Series: United States. National Bureau of Standards. National standard reference data series ; NSRDS-NBS 65.

QC100.U573 no. 65 [QD502] 602'.1s [541'.394] 79-607030

NSRDS-NBS 65

Nat. Stand. Ref. Data Ser., Nat. Bur. Stand. (U.S.), 65, 62 pages (June 1979)

CODEN: NSRDAP

© 1979 by the Secretary of Commerce on Behalf of the United States Government

U.S. GOVERNMENT PRINTING OFFICE

WASHINGTON: 1979

For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402

Stock No. 003-003-02072-9 Price \$3.50

(Add 25 percent additional for other than U.S. mailing).

Foreword


The National Standard Reference Data System provides access to the quantitative data of physical science, critically evaluated and compiled for convenience and readily accessible through a variety of distribution channels. The System was established in 1963 by action of the President's Office of Science and Technology and the Federal Council for Science and Technology, and responsibility to administer it was assigned to the National Bureau of Standards.

NSRDS receives advice and planning assistance from a Review Committee of the National Research Council of the National Academy of Sciences-National Academy of Engineering. A number of Advisory Panels, each concerned with a single technical area, meet regularly to examine major portions of the program, assign relative priorities, and identify specific key problems in need of further attention. For selected specific topics, the Advisory Panels sponsor subpanels which make detailed studies of users' needs, the present state of knowledge, and existing data resources as a basis for recommending one or more data compilation activities. This assembly of advisory services contributes greatly to the guidance of NSRDS activities.

The System now includes a complex of data centers and other activities in academic institutions and other laboratories. Components of the NSRDS produce compilations of critically evaluated data, reviews of the state of quantitative knowledge in specialized areas, and computations of useful functions derived from standard reference data. The centers and projects also establish criteria for evaluation and compilation of data and recommend improvements in experimental techniques. They are normally associated with research in the relevant field.

The technical scope of NSRDS is indicated by the categories of projects active or being planned: nuclear properties, atomic and molecular properties, solid state properties, thermodynamic and transport properties, chemical kinetics, and colloid and surface properties.

Reliable data on the properties of matter and materials are a major foundation of scientific and technical progress. Such important activities as basic scientific research, industrial quality control, development of new materials for building and other technologies, measuring and correcting environmental pollution depend on quality reference data. In NSRDS, the Bureau's responsibility to support American science, industry, and commerce is vitally fulfilled.

A handwritten signature in dark ink, reading "E. Ambler." The signature is fluid and cursive, with a large initial "E" and a trailing period.

ERNEST AMBLER, *Director*

Preface

This report is one of a series of data publications on radiation chemistry; the aim of the series is to compile, evaluate, and present the numerical results on processes occurring in systems which have been subjected to ionizing radiation. Various kinds of data are important in radiation chemistry. The quantities which were measured first were the observed radiation yields or G values (molecules formed or destroyed per 100 eV). Various indirect methods based on G values have been used to determine yields of transient species and relative rates of reactions. The spectral properties (optical, electron spin resonance) of transients have provided a direct method for their identification, and rates of the very fast reactions of transients which occur in irradiated systems have been measured directly by spectroscopic methods. Conductivity and luminescence methods have also provided a means of measuring properties of transients and their kinetics. Some reactions which occur in irradiated systems have also been studied by other methods, such as photochemistry, electric discharge, ultrasonics, chemical initiation, electron impact, etc. The emphasis in these publications is on the data of radiation chemistry, but where other pertinent data exist, they are included.

The data of radiation chemistry are voluminous; thousands of systems have been investigated. As a result there are certain collections, *e.g.* rate constants of particular types of reactions or certain properties of transients, for which tabulations of the data are considered essential, but for which critical assessment of each value is impossible. On the other hand, certain systems and properties have been studied so extensively that critical examination of these data is desirable and timely. Authors of this series of data publications have been asked to evaluate the extent to which the data can be critically assessed, to describe their criteria for evaluation, and to designate preferred values whenever possible.

Contents

	Page
Introduction.....	1
Table 1. CO_3^- reactions. (1.1–1.100).....	5
Table 2. SO_4^- reactions. (2.1–2.116).....	14
Table 3. SO_2^- , SO_3^- , S_2O_3^- , S_2O_2^- , SO_5^- and SeO_3^- reactions. (3.1–3.16).....	23
Table 4. NO_3 , NO_3^{2-} , NO_2 , N_2O_4 , N_2O_3 , NO_2^{2-} reactions. (4.1–4.25).....	25
Table 5. PO_4^{2-} reactions. (5.1–5.73).....	28
Table 6. Cl_2^- reactions. (6.1–6.100).....	33
Table 7. Br_2^- reactions. (7.1–7.66).....	38
Table 8. I_2^- reactions. (8.1–8.30).....	42
Table 9. $(\text{SCN})_2^-$ reactions. (9.1–9.41).....	44
Formula Index	46
References	49

Rate Constants for Reactions of Inorganic Radicals in Aqueous Solution

Alberta B. Ross and Pedatsur Neta

Radiation Chemistry Data Center, Radiation Laboratory*,
University of Notre Dame, Notre Dame, IN 46556

Rate constants have been compiled for reactions of various transient inorganic radicals produced by radiolysis or photolysis in aqueous solution. Data are included for the carbonate radical, sulfate radical, phosphate radical, nitrate radical and other nitrogen-, sulfur- and selenium-containing radicals, and the halide and pseudohalide radicals Cl_2^- , Br_2^- , I_2^- and $(\text{SCN})_2^-$. The radicals react with other inorganic ions, as well as aliphatic, aromatic and heterocyclic compounds.



Keywords: Aqueous solution; carbonate radical; chemical kinetics; halogen radical anions; nitrate radical; oxyanion radicals; phosphate radical; photolysis; radiolysis; rates; selenium radicals; sulfate radical; thiocyanate radical anion.

Introduction


The radiolysis of water produces short-lived intermediates, hydrated electrons, hydrogen atoms and hydroxyl radicals, which react rapidly with solutes in the water to form other short-lived reaction intermediates (73-0030, 75-0001, 75-0002, 77-0011)¹. This compilation includes reaction rates for short-lived species derived from a number of inorganic ions, including carbonate, sulfate, sulfite, phosphate, nitrate, thiocyanate, and the halide ions by radiolysis or photolysis. The radicals react by oxidizing a wide variety of solutes in aqueous media, most commonly by electron transfer, but also by hydrogen abstraction, addition, substitution, and disproportionation.

Methods


Carbonate radical. The CO_3^- radical is produced by oxidation of carbonate or bicarbonate ions with OH or SO_4^- radicals,



by photoionization of carbonate



or by photolysis of certain carbonato-metal complexes, e.g.



The CO_3^- radical has a broad optical absorption with λ_{max} 600 nm and ϵ_{max} 1880 $M^{-1} \text{ cm}^{-1}$ (66-0001). The protonation



was suggested to have $\text{p}K_a$ 9.6 (73-7109), but it does not result in shifts in the optical (73-7109) or ESR spectra (75-5244).

*The Radiation Laboratory is operated under Contract EY-76-C-02-0038 with the Department of Energy. The Radiation Chemistry Data Center is supported jointly by the National Bureau of Standards, Office of Standard Reference Data and the Department Energy. This is Radiation Laboratory Document No. NDRL-1933.

¹ Literature references are cited following the tables.

The majority of the rate constants for reactions of CO_3^- or CO_3H were determined by following the decay of these radicals through their optical absorption. The carbonate radical can oxidize certain aromatic and heterocyclic compounds and inorganic ions, probably by direct electron transfer. It can also abstract hydrogen from aliphatic compounds, but this reaction is generally slow. Intermediate or stable products from CO_3^- reactions were identified in only a limited number of cases and the mechanisms are mostly speculative.

Sulfate radical. The SO_4^- radical is produced from peroxodisulfate ions by reaction with e_{aq}^-



Reaction with various other radicals and metal ions can also yield SO_4^- . The other most commonly used method for production of this radical is photolysis of $\text{S}_2\text{O}_8^{2-}$



Oxidation of sulfate ions to yield SO_4^- radicals can be also carried out photolytically or radiolytically, but those processes are relatively inefficient and were not used frequently for obtaining rate constants.

The SO_4^- radical has an optical absorption at λ_{max} 450 nm and ϵ_{max} 1100 $M^{-1} \text{ cm}^{-1}$ (66-0019, 69-0158). This radical may protonate in strongly acidic solutions (67-7274) but no $\text{p}K_a$ value has been determined. The rate constants for reactions of SO_4^- were determined in most cases by following the decay of the 450 nm absorption. In some cases, the kinetics of formation of the product radicals was monitored.






This radical is a strong oxidant which can react with many organic and inorganic compounds by direct electron transfer oxidation. The reaction with benzene and most aromatic and heterocyclic compounds involves this mechanism. However, addition to aliphatic double bonds and abstraction from saturated compounds were also shown to take place. The products and mechanism of reaction of this radical were studied in some detail, many of which are summarized under the corresponding reactions in table 2.

Phosphate radical. The PO_4^{2-} radical and its protonated forms are produced from peroxodiphosphate or phosphate ions by the same methods as described for SO_4^- . The acid-base forms and their optical absorption properties can be summarized as follows (78-1075):

	H_2PO_4	HPO_4^-	PO_4^{2-}
$\text{p}K_a$	5.7	8.9	
λ_{max}	520	510	530 nm
ϵ_{max}	1850	1550	2150 $M^{-1} \text{ cm}^{-1}$


The rate constants for reactions of these radicals were determined in most cases by following the decay of their broad absorption in the 500-540 nm region. These radicals can abstract hydrogen from saturated organic compounds, add to olefins, and oxidize many aromatic compounds. H_2PO_4 is somewhat similar to SO_4^- in its reactivity, but HPO_4^- and PO_4^{2-} are less reactive, both in H abstraction reactions and in electron transfer oxidations.

Other sulfur-, nitrogen- and selenium radicals. Various radicals have been produced from sulfite, selenite, nitrite, nitrate, and related anions, by one-electron oxidation or reduction, carried out either radiolytically or photolytically. Some of the details for the production and observation of these radicals are given in tables 3 and 4. Representative examples are:




Only a few rate constants have been reported for these various radicals, most of them are for the radical-radical reaction. The reactions of these radicals with organic and inorganic compounds were studied in only a very limited number of cases so that the rates and mechanisms are largely unknown.

Halide and pseudohalide radicals. Radicals of the type X_2^- can be produced from halides and some pseudohalides. Tables 6 – 9 list the rates for Cl_2^- , Br_2^- , I_2^- and $(SCN)_2^-$, respectively. The radicals are produced by oxidation of the halide ion followed by the complex formation. Oxidation of X^- can be achieved in most cases by reaction with OH




although complex and acid-base equilibria are involved, which cause the Cl^- reaction to be efficient only in acid solution (73–1039). Oxidation of Cl^- in neutral and alkaline solutions can be carried out by SO_4^- (e.g. 75–5244).



Photolysis of halides can also result in their oxidation.

The X_2^- radicals are formed in the equilibrium



which is shifted predominantly to the right ($K \cong 10^5 M^{-1}$) (65–0383, 68–0375, 73–1039). Their reaction kinetics are conveniently studied by observing their intense absorptions near 340 for Cl_2^- , 360 for Br_2^- , 380 for I_2^- and 480 for $(SCN)_2^-$, all of the radicals having ϵ of the order of $10^4 M^{-1} cm^{-1}$.

The reactivities of the X_2^- radicals generally decrease in this order: Cl_2^- , Br_2^- , $(SCN)_2^-$, I_2^- . Cl_2^- can abstract hydrogen from aliphatic compounds but the equivalent reactions of the other X_2^- radicals are barely detectible in the pulse experiments. The reaction of Cl_2^- with some unsaturated compounds produces Cl adducts, and presumably the other X_2^- radicals may react in a similar fashion, although little information is available on them. The main reaction of the X_2^- radicals is the one-electron oxidation, which takes place with many aromatic and heterocyclic compounds and with certain inorganic ions. However, X_2^- radicals are weaker oxidants than SO_4^- and H_2PO_4 . The latter can oxidize benzene and most of its derivatives, while X_2^- can only oxidize phenols, anilines, and anisoles.

Arrangement of the Tables

Entries in each table have been numbered in order to provide for the assembly of an index. Within the tables reactions of the transient with the same or other transient species are listed first. Next are reactions of inorganic solutes which have been ordered alphabetically by main element; the organic reactants follow in alphabetical order by name. The reaction is given whenever evidence for the nature of the products is available; otherwise, only the reactants are given in column 2.

Rate Constants (k): Most of the second order rate constants are derived from pseudo-first order rates for the decay of the transient or the build-up of a product. The error limits given are those reported in the original papers; lacking such a statement the uncertainty in the rate constant should be assumed to be $\pm 25\%$. Second order rates for reaction of two radicals have usually been derived from measurements of $2k/\epsilon$; the value of ϵ used to calculate $2k$ has been listed in the *Comments* column. Possibilities for error in the determination of ϵ increase the error limits for $2k$, which should be assumed to be $\pm 50\%$, lacking a specific error statement.

In some cases the rates are designated (rel.) and have been calculated with reference to the rate of a competing reaction, given in the *Comments* column. Ratios of rates are listed for a few reactions in the form k/k_x in which k_x is the rate of reaction of the transient with competing solute, X.

Ionic Strength (I): The ionic strength has been quoted from the original paper or calculated from the given solute concentrations, if that information was available. When the reported rate constants were corrected for ionic strength the designation $\rightarrow 0$ is used.

Comments: The source of the transient and method of measurement are included in this column, along with other information on product identification, or related values for equilibrium constants, pK , or isotope effects. Temperature is assumed to be room temperature unless otherwise specified.

References: The references are designated by the serial number of the paper assigned by the Radiation Chemistry Data Center; the first two digits of the number represent the year in which the work was published.

Abbreviations and Symbols

A	frequency factor	ident.	identification
abs.	absorption	K	equilibrium constant
alk.	alkaline	k_f	specific rate of the forward reaction
bipy	2,2'-bipyridine	k_r	specific rate of the reverse reaction
BuOH	butanol	L	ligand
calcd.	calculated	M	mol dm^{-3}
c.k.	competition kinetics	meas.	measured
concn.	concentration	MeOH	methanol
contg.	containing	obs.	observed
cor.	corrected	ϕ	quantum yield
detd.	determined	p.b.k.	product-buildup kinetics
dien	diethylenetriamine	phot.	photolysis
d.k.	decay kinetics	polymn.	polymerization
e-r.	electron radiolysis	p.r.	pulse radiolysis
ϵ	extinction coefficient (molar absorptivity)	2-PrOH	2-propanol
E_a	activation energy	rel.	relative
esr	electron spin resonance	RNO	<i>N,N</i> -dimethyl- <i>p</i> -nitrosoaniline
estd.	estimated	$\Delta S^\ddagger$	activation entropy
f.phot.	flash photolysis	satd.	saturated
formn.	formation	soln.	solution
G	radiation yield (molecules per 100 eV)	T	temperature
ΔG°	free energy	$t_{0.5}$	half-life
γ -r.	gamma radiolysis	therm.	thermal
gly	glycinato	unpub.	unpublished
$\Delta H^\ddagger$	activation enthalpy		

TABLE 1. Rates of reaction of CO_3^- in aqueous solution*

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
1.1	$\text{CO}_3^- + \text{CO}_3^{2-} \rightarrow \text{CO}_2 + \text{CO}_4^{2-}$ See 78-1506 for mechanism	$2k = 1.2_3 \times 10^7$	8.4-13.5	$\rightarrow 0$	p.r.	D.k. at 600 nm in N_2O -satd. soln.; $\epsilon = 1860 M^{-1} \text{cm}^{-1}$; $k_{\text{obs}} \approx 5 \times 10^7$ at pH 13-13.5	66-0139
		$2k = 4 \times 10^7$	7-9	0.1	f.phot.	D.k. at 600 nm in air-satd. $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.; $\epsilon = 1830 M^{-1} \text{cm}^{-1}$.	73-7109
		$2k = 1.9 \times 10^7$ $2k = 2.9 \times 10^7$	10.2 12.7	-	p.r.	D.k. at 600 nm in N_2O -satd. soln.; $\epsilon = 1830 M^{-1} \text{cm}^{-1}$; 0.1 M K_2CO_3 .	73-7109
		$2k = 6.9 \times 10^7$	11.8	-	p.r.	D.k. in 0.1 M Na_2CO_3 soln.; See above.	73-7109
1.2	$\text{CO}_3^- + \text{O}_2^- \rightarrow \text{CO}_3^{2-} + \text{O}_2$ (or $\rightarrow \text{CO}_5^{2-}$) See 70-0247)	1.5×10^9	11.6	0.3	p.r.	D.k. of CO_3^- at 600 nm or O_2^- at 260 nm in O_2 -satd. soln.; $\epsilon(260 \text{ nm})$ for $\text{O}_2^- = 1220 M^{-1} \text{cm}^{-1}$ assuming $\epsilon(600 \text{ nm})$ for $\text{CO}_3^- = 1800 M^{-1} \text{cm}^{-1}$.	66-0001
		$(4 \pm 1) \times 10^8$	~ 11.8	0.6	f.phot.	D.k. at 260 nm (O_2^-), $\epsilon = 1850 M^{-1} \text{cm}^{-1}$ and 600 nm (CO_3^-) in O_2 -satd. soln., $\epsilon = 1860 M^{-1} \text{cm}^{-1}$; product (CO_5^{2-} ?) has $\epsilon(260 \text{ nm}) = 410 M^{-1} \text{cm}^{-1}$.	70-0247
		7.5×10^8	8-13	-	f.phot.	D.k. at 600 nm in air-satd. soln. contg. NaHCO_3 and 2,6-anthraquinonesulfonate.	72-7335, 72-7464
1.2a	$\text{CO}_3^- + \text{Br}^-$	$< 5 \times 10^5$	~ 11	-	p.r.	No reaction.	78-1506
1.3	$\text{CO}_3^- + \text{BrO}^- \rightarrow \text{CO}_3^{2-} + \text{BrO}$	$(4.3 \pm 0.4) \times 10^7$	13	0.4	p.r.	D.k.	68-0153
1.4	$\text{CO}_3^- + \text{BrO}_2^- \rightarrow \text{CO}_3^{2-} + \text{BrO}_2$	$(1.1 \pm 0.1) \times 10^8$	13	0.4	p.r.	D.k.	68-0153
<i>Co(II) ions</i>							
1.5	$\text{CO}_3^- + \text{Co}_{\text{aq}}^{2+}$	$(4.4 \pm 0.4) \times 10^6$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7109
		2.8×10^6	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.5a	$\text{CO}_3^- + \text{CoL}(\text{H}_2\text{O})_2^{2+}$ L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-1,4,8,11-tetraene	6.9×10^8	4.7	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.5b	$\text{CO}_3^- + \text{CoL}(\text{H}_2\text{O})_2^{2+}$ L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene	7.3×10^8	4.7	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
<i>Co(III) complexes</i>							
1.6	$\text{CO}_3^- + \text{Co}(\text{NH}_3)_4\text{CO}_3^+$	$< 10^6$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7109
1.7	$\text{CO}_3^- + \text{Co}(\text{NH}_3)_4(\text{OH})_2^{3+}$	$(1.4 \pm 0.1) \times 10^7$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7109
1.7a	$\text{CO}_3^- + \text{Co}(\text{NH}_3)_6^{3+}$	$< 5 \times 10^4$	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.7b	$\text{CO}_3^- + \text{Co}(\text{en})_3^{3+}$	$< 1 \times 10^5$	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.7c	$\text{CO}_3^- + \text{Co}(\text{NH}_3)_5\text{OPO}_3\text{H}^+$	$< 4 \times 10^5$	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.7d	$\text{CO}_3^- + \text{Co}(\text{NH}_3)_5\text{Cl}^{2+}$	2.0×10^6	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501

TABLE 1. Rates of reaction of CO_3^- in aqueous solution^a—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
1.7e	$\text{CO}_3^- + \text{Co}(\text{NH}_3)_5\text{Br}^{2+}$	5.7×10^6	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.7f	$\text{CO}_3^- + \text{Co}(\text{NH}_3)_5\text{OSO}_3^+$	1.5×10^6	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.7g	$\text{CO}_3^- + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CC}_6\text{H}_5^{2+}$	7×10^5	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.7h	$\text{CO}_3^- + \text{Co}(\text{NH}_3)_5\text{O}_2\text{CCH}_3^{2+}$	1.1×10^6	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.7i	$\text{CO}_3^- + \text{Co}(\text{NH}_3)_5\text{NO}_2^{2+}$	1.0×10^8	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.7j	$\text{CO}_3^- + \text{Co}(\text{NH}_3)_5\text{OSO}_2^+$	1.1×10^6	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.8	$\text{CO}_3^- + \text{Co}(\text{NH}_3)_5(\text{OH}_2)^{3+}$	$(2.9 \pm 0.3) \times 10^6$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7109
		1.7×10^6	>7.6	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
		$\sim 3 \times 10^4$	<5.6	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.; pK_a for complex = 6.6.	78-1501
1.8a	$\text{CO}_3^- + \text{Cr}(\text{NH}_3)_5\text{Cl}^{2+}$	$< 1 \times 10^5$	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.8b	$\text{CO}_3^- + \text{Cr}(\text{NH}_3)_5\text{OH}_2^{3+}$	7×10^5	>6.3	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
		4×10^5	<4.3	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.; pK_a for complex = 5.3.	78-1501
1.8c	$\text{CO}_3^- + \text{Cu}_{aq}^{2+}$	$\sim 10^4$	4.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.9	$\text{CO}_3^- + \text{Fe}(\text{CN})_6^{4-} \rightarrow \text{CO}_3^{2-} + \text{Fe}(\text{CN})_6^{3-}$	2.7×10^8	11.6	—	p.r.	D.k. at 600 nm as well as p.b.k. at 420 nm for ferricyanide ion in soln. contg. 0.072 M Na_2CO_3 , 1.22×10^{-4} M ferrocyanide, 0.03% oxygen and 0.1 atm. N_2O ; at pH 13 $k \cong (3.5-4.0) \times 10^8$.	66-0139
1.9a	$\text{CO}_3^- + \text{H}_2\text{O}_2 \rightarrow \text{CO}_3^{2-} + \text{HO}_2^-$	8×10^5 5.6×10^7	8-9 13-14	3	f.phot.	D.k. at 600 nm in N_2 -satd. soln.; pH-dependent (9-13).	70-0247
1.10	$\text{CO}_3^- + \text{I}^- \rightarrow \text{CO}_3^{2-} + \text{I}$	$(1.3 \pm 0.3) \times 10^8$	~ 11	$\rightarrow 0$	p.r.	D.k. at 600 nm.	78-1506
1.10a	$\text{CO}_3^- + \text{Ir}(\text{NH}_3)_5\text{Cl}^{2+}$	2.4×10^7	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.10b	$\text{CO}_3^- + \text{Mn}_{aq}^{2+}$	1.5×10^7	6.0	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.10c	$\text{CO}_3^- + \text{NO}_2^- \rightarrow \text{CO}_3^{2-} + \text{NO}_2$	4.0×10^5	~ 11	$\rightarrow 0$	p.r.	D.k. at 600 nm.	78-1506
1.10d	$\text{CO}_3^- + \text{NO}_2 \rightarrow \text{CO}_2 + \text{NO}_3^-$	1.0×10^9	~ 11	—	p.r.	Est. from opt. and condy. d.k.	78-1506
1.10e	$\text{CO}_3^- + \text{Ni}_{aq}^{2+}$	$< 10^4$	5.8	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.11	$\text{CO}_3^- + \text{cis-Pt}^{\text{II}}(\text{gly})_2$	$(4.4 \pm 0.8) \times 10^9$	8.5	0.05	p.r.	D.k. in 0.05 M NaHCO_3 .	77-1053
1.12	$\text{CO}_3^- + \text{trans-Pt}^{\text{II}}(\text{gly})_2$	$(3.4 \pm 0.6) \times 10^9$	8.5	0.05	p.r.	D.k. in 0.05 M NaHCO_3 .	77-1053
1.12a	$\text{CO}_3^- + \text{Rh}(\text{NH}_3)_5\text{Cl}^{2+}$	$< 10^4$	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.12b	$\text{CO}_3^- + \text{Rh}(\text{NH}_3)_5\text{OH}_2^{3+}$	1×10^5	>6.9	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
		$< 5 \times 10^4$	<4.9	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.; pK_a for complex = 5.9.	78-1501

TABLE 1. Rates of reaction of CO_3^- in aqueous solution^a—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
1.13	$\text{CO}_3^- + \text{Ru}(\text{bipy})_3^{2+}$	$(4.0 \pm 0.4) \times 10^8$	—	0.02	p.r.	D.k. of the Ru complex in N_2O -satd. soln. contg. 0.005 M each of HCO_3^- and CO_3^{2-} .	77-1093
1.13a	$\text{CO}_3^- + \text{Ru}(\text{NH}_3)_6^{3+}$	6.0×10^6	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.13b	$\text{CO}_3^- + \text{Ru}(\text{NH}_3)_5\text{Cl}^{2+}$	7.7×10^6	6.5	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.13c	$\text{CO}_3^- + \text{Ru}(\text{NH}_3)_5\text{OH}_2^{3+}$	1.4×10^9	>5.2	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
		1.8×10^8	<3.2	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.; pK_a for complex = 4.2.	78-1501
1.13d	$\text{CO}_3^- + \text{SCN}^- \rightarrow \text{CO}_3^{2-} + \text{SCN}$	$(8 \pm 1.5) \times 10^5$	~ 11	—	p.r.	D.k. at 600 nm.	78-1506
1.13e	$\text{CO}_3^{2-} + \text{SO}_3^{2-} \rightarrow \text{CO}_3^{2-} + \text{SO}_3^{2-}$	1.0×10^7	~ 11	$\rightarrow 0$	p.r.	D.k. at 600 nm.	78-1506
1.13f	$\text{CO}_3^- + \text{SO}_3^{2-} \rightarrow \text{CO}_2 + \text{SO}_4^{2-}$	5.0×10^8	9.6	—	p.r.	D.k. at 260 nm; also condy. study	78-1506
1.14	$\text{CO}_3^- + \text{UO}_2^{2+}$	$\sim 1.5 \times 10^5$	—	0.16	f.phot.	D.k. at 580 nm; results somewhat irreproducible.	76-7279
1.14a	$\text{CO}_3^- + \text{Zn}_{aq}^{2+}$	$< 10^4$	4.7	0.04	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	78-1501
1.15	Acetanilide	3.2×10^5	7.0	0.06	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	75-7313
1.16	Acetate ion	6×10^2	12.1–12.7	—	f.phot.	D.k. at 600 nm in air-satd. soln. contg. 0.05 M $\text{Na}_2\text{S}_2\text{O}_8$, 0.5 M Na_2CO_3 ; CO_3^- from reaction of $\text{SO}_4^{2-} + \text{CO}_3^{2-}$.	72-7383
1.17	Acetone	1.6×10^2	12.1–12.7	—	f.phot.	D.k. at 600 nm in air-satd. soln. contg. 0.05 M $\text{Na}_2\text{S}_2\text{O}_8$, 0.5 M Na_2CO_3 ; CO_3^- from reaction of $\text{SO}_4^{2-} + \text{CO}_3^{2-}$.	72-7383
1.18	Acetonitrile	3.2×10^3	12.1–12.7	—	f.phot.	D.k. at 600 nm in air-satd. soln. contg. 0.05 M $\text{Na}_2\text{S}_2\text{O}_8$, 0.5 M Na_2CO_3 ; CO_3^- from reaction of $\text{SO}_4^{2-} + \text{CO}_3^{2-}$.	72-7383
1.19	Acetophenone	1.0×10^7	~ 12.5	—	f.phot.	D.k. at 550 nm in deoxy-generated soln. contg. 0.3 M Na_2CO_3 .	71-7574
		$(3 \pm 1) \times 10^5$	—	—	p.r.	D.k. at 600 nm; CO_3^- also reacts with acetophenone-OH adduct, $k = (1.5 \pm 0.5) \times 10^9$.	78-1506
1.20	N-Acetylcysteine	$(1.8 \pm 0.4) \times 10^8$	12.0	0.003	f.phot.	D.k. at 600 nm; rate pH dependent; value from graph; CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$.	75-7110
1.21	N-Acetylglycine	$< 10^4$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.22	N-Acetylglycylglycine	$< 10^4$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.23	N-Acetyltryptophan	$(4.2 \pm 0.4) \times 10^8$ $(6.2 \pm 1.5) \times 10^8$	7 11.8	0.1 0.1	f.phot. f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7109
1.24	Alanine	$< 10^3$	7.0	0.06	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352

TABLE 1. Rates of reaction of CO_3^- in aqueous solution^a—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
1.25	Aniline	5.4×10^8	7.0	0.06	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	75-7313
1.25a	Anisole	6.0×10^8	—	—	p.r.	D.k. at 600 nm.	78-1506
		2.8×10^5	—	—	p.r.	D.k. at 600 nm; CO_3^-	78-1506
						also reacts with anisole-OH adduct, $k \approx 3 \times 10^9$	
1.26	Anthrasemiquinone-1-sulfonate radical ion $\cdot\text{CO}_3^- + \text{AQ}^- \rightarrow \text{CO}_3^{2-} + \text{AQ}$	4.6×10^9	—	—	f.phot.	D.k. at 600 nm in air-satd. soln. contg. 0.05 M $\text{Na}_2\text{S}_2\text{O}_8$ and 0.5 M Na_2CO_3 ; CO_3^- from reaction of $\text{SO}_4^- + \text{CO}_3^{2-}$; semiquinone formed from $\text{CO}_3^{2-} +$ triplet AQ.	72-7383
1.27	Anthrasemiquinone-2-sulfonate radical ion	2.2×10^9	—	—	f.phot.	See 1.26.	72-7383
1.28	Anthrasemiquinone-2,6-disulfonate radical ion	$(5.4 \pm 0.6) \times 10^8$	7.5	—	f.phot.	D.k. in Na_2CO_3 soln.	69-7297
		2×10^9	8.0	—	f.phot.	D.k. in O_2 -free carbonate soln.	73-7569
		1.2×10^9	6.5	—	f.phot.	D.k. in NaHCO_3 soln.	72-7335
		1.4×10^9	8-13	—	f.phot.	D.k. in air-satd. 0.05 M NaHCO_3 soln.	72-7464
1.29	Anthrasemiquinone-2,7-disulfonate radical ion	1.9×10^9	—	—	f.phot.	See 1.26.	72-7383
1.30	Arginine	$(9 \pm 0.9) \times 10^4$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.31	Ascorbate ion $\text{CO}_3^- + \text{AH}^- \rightarrow \cdot\text{A}^- + \text{CO}_3^{2-} + \text{H}^+$	$(1.1 \pm 0.1) \times 10^9$	11	—	p.r.	D.k. at 600 nm in N_2O -satd. 0.5 M carbonate soln.	73-3006
1.32	Aspartate ion	$< 10^4$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.33	Benzene	3×10^3	7.0	0.06	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln; uncertainty 50-100%.	75-7313
		$< 5 \times 10^4$	11.7	—	p.r.	Benzene ($1.4 \times 10^{-3} M$) had no effect on decay of CO_3^- ; CO_3^- reacts with benzene-OH adduct, $k = 2 \times 10^9$	78-1506
1.34	Benzophenone	1.5×10^6	—	—	f.phot.	D.k. at 550 nm in O_2 -free carbonate soln.	71-7574
1.35	<i>p</i> -Bromophenoxide ion <i>tert</i> -Butanol See 2-Methyl-2-propanol.	1.8×10^8	12.2	0.3	p.r.	D.k. at 570 nm in N_2O -satd. carbonate soln.	77-1098
1.36	Carbon tetrachloride	—	12.0-13.5	—	p.r.	Decay of CO_3^- faster in CCl_4 than in N_2O soln.	66-0139
1.37	Chloroacetate ion	$< 2.0 \times 10^3$	12.1-12.7	1.65	f.phot.	D.k. at 600 nm in air-satd. $\text{S}_2\text{O}_8^{2-}$ soln.; CO_3^- generated by $\text{SO}_4^- + \text{CO}_3^{2-} \rightarrow \text{SO}_4^{2-} + \text{CO}_3^-$.	72-7383
1.38	<i>p</i> -Chlorophenoxide ion	1.9×10^8	12.2	0.3	p.r.	D.k. at 570 nm in N_2O -satd. carbonate soln.	77-1098
1.39	α -Chymotrypsin	1.0×10^9	11.3	0.12	p.r.	D.k. in N_2O -satd. soln.; mol wt. 20,000; concn. effect.	74-1096
		$(1.2 \pm 0.1) \times 10^9$	7-11	0.03	f.phot.	D.k. at 600 nm; mol. wt. 25,000; HCO_3^- or CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; k independent of pH.	73-7352, 75-7110

TABLE 1. Rates of reaction of CO_3^- in aqueous solution^a—Continued

No.	Reaction	$k(\text{M}^{-1} \text{s}^{-1})$	pH	I	Method	Comment	Ref.
1.40	Cysteine	$(2.7 \pm 0.3) \times 10^8$	11.2 ± 0.3	0.3	p.r.	D.k. in N_2O -satd. soln.	72-0036
		$(4.6 \pm 0.5) \times 10^7$	7.0	0.03	f.phot.	D.k. at 600 nm; HCO_3^-	73-7352,
		3.5×10^8	9-10	0.003	f.phot.	or CO_3^- generated from	75-7110
		2.5×10^8	12	0.003	f.phot.	$\text{Co}(\text{NH}_3)_4\text{CO}_3^+$.	
1.41	Cysteine methyl ester	$\sim 1 \times 10^7$	4-10	0.003	f.phot.	D.k. at 600 nm; value from graph; HCO_3^- or CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$.	75-7110
1.42	Cystine dimethyl ester	$(7.2 \pm 0.7) \times 10^6$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.43	Diethyl disulfide	4.5×10^7 6.6×10^7	~ 8 ~ 11	—	p.r.	D.k. at 600 nm in N_2O -satd. soln.; 0.1-1.0 M HCO_3^- or CO_3^{2-} .	76-1143
1.44	<i>N,N</i> -Dimethylaniline	1.8×10^9	7.0	0.06	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	75-7313
		$(1.4 \pm 0.2) \times 10^9$	—	—	p.r.	D.k. at 600 nm, as well as condy.	78-1506
1.45	Dimethyl disulfide	1.0×10^8 8.0×10^7	~ 8 ~ 11	—	p.r.	D.k. at 600 nm in N_2O -satd. soln.; 0.1-1.0 M HCO_3^- or CO_3^{2-} .	76-1143
1.46	3,3'-Dithiobis(propionate ion)	$(1.3 \pm 0.1) \times 10^7$ $(3.0 \pm 0.3) \times 10^7$	6.8 11.5	0.1	f.phot.	D.k. at 600 nm; CO_3H and CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$.	73-7109
		$(1.0 \pm 0.1) \times 10^7$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
		1.3×10^7	7-12	0.003	f.phot.	D.k. at 600 nm; CO_3H and CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; value from graph.	75-7110
1.47	Dithiothreitol (<i>threo</i> -2,3-Dihydroxy-1,4-dithiobutane)	$(4.1 \pm 0.4) \times 10^8$	11	0.3	p.r.	D.k. at 600 nm in N_2O -satd. carbonate soln.	73-1020
1.48	Duroquinone	$< 10^6$	12	—	p.r.	No reaction; previously reported (76-7587) $k = 2 \times 10^9$ suggested to be for different reaction.	78-1506
1.49	Ethanol $\text{CO}_3^- + \text{CH}_3\text{CH}_2\text{OH} (+\text{OH}^-) \rightarrow \text{CO}_3^{2-} + \text{CH}_3\text{CHOH} + \text{H}_2\text{O}$	$1.5-1.7 \times 10^4$	12.5	—	f.phot.	D.k. in soln. contg. 0.3 M Na_2CO_3 and $2 \times 10^{-4} M$ acetophenone or benzophenone.	71-7574
		1.5×10^4	12.5	—	f.phot.	D.k. at 600 nm in air-satd. soln. contg. 0.05 M $\text{Na}_2\text{S}_2\text{O}_8$ and 0.5 M Na_2CO_3 ; CO_3^- generated from $\text{SO}_4^{2-} + \text{CO}_3^{2-} \rightarrow \text{SO}_4^{2-} + \text{CO}_3^-$.	72-7383
1.50	Ethoxybenzene	4.1×10^5	7.0	0.06	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	75-7313
1.50a	Ethylenediaminetetraacetate ion	1.1×10^6	—	$\rightarrow 0$	p.r.	D.k. at 600 nm.	78-1506
1.51	Formate ion	$(1.1 \pm 0.1) \times 10^5$	6.4	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7109
		1.6×10^5	—	$\rightarrow 0$	p.r.	D.k. at 600 nm.	78-1506
1.52	Glucose	$(7 \pm 0.7) \times 10^4$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.53	Glutathione	$(5.3 \pm 0.5) \times 10^6$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.54	Glutathione disulfide	$(1.3 \pm 0.1) \times 10^6$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352

TABLE 1. Rates of reaction of CO_3^- in aqueous solution^a—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	l	Method	Comment	Ref.
1.55	Glycine	$< 10^3$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.56	Glycylglycine	$(2 \pm 0.2) \times 10^4$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.57	Glycylglycylglycine	$(4 \pm 0.4) \times 10^4$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.58	Glycylglycyltryptophan	$(7 \pm 0.7) \times 10^8$	7.0	0.003	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln; k is pH dependent, $k = 4 \times 10^8$ at pH 10.	74-7296
1.59	Glycylhistidine	$(4.3 \pm 0.4) \times 10^6$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.60	Glycyltryptophan	$(7.2 \pm 0.7) \times 10^8$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
		$(7.8 \pm 0.8) \times 10^8$	7	0.003	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; k is pH dependent, $k = 4.5 \times 10^8$ at pH 10.	74-7296
1.61	Glycyltyrosine	$(3.0 \pm 0.3) \times 10^7$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.62	Histidine	$(7 \pm 0.7) \times 10^6$	11.2 ± 0.3	0.3	p.r.	D.k. in N_2O -satd. soln.	72-0036
		$(5.6 \pm 0.6) \times 10^6$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
		8.5×10^6	10	0.003	f.phot.	D.k. at 600 nm; k is pH dependent; $k \sim 1 \times 10^6$ at pH 5 (values from graph); CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$.	75-7110
1.63	<i>p</i> -Hydroxybenzoate ion	7.9×10^7	12.2	0.3	p.r.	D.k. at 570 nm in N_2O -satd. carbonate soln.	77-1098
1.64	Imidazole	$(5.5 \pm 0.6) \times 10^5$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.65	Indole	3.2×10^8	7.0, 12.0	0.1	f.phot.	D.k. at 600 nm; CO_3H or CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$.	73-7109
		4.1×10^8	5-13	0.003	f.phot.	D.k. at 600 nm; CO_3H or CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; value from graph.	74-7296
1.66	Indole-3-propionate ion	$(4.1 \pm 0.4) \times 10^8$	7.0	0.1	f.phot.	D.k. at 600 nm; CO_3H and CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$.	73-7109
		$(6.8 \pm 1.7) \times 10^8$	12.0	0.1			
		4.2×10^8	7-11	0.003	f.phot.	D.k. at 600 nm; CO_3H and CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; at $\mu = 0.1$ k increases with pH; values from graph.	74-7296
	Isopropanol	See 2-Propanol.					
1.67	Lysozyme	$(5.5 \pm 0.6) \times 10^8$	7.0-12	0.03	f.phot.	D.k. at 600 nm; CO_3H and CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$.	73-7352 75-7110
1.68	3-Mercaptopropionate ion	$(2.4 \pm 0.6) \times 10^8$	12.0	0.003	f.phot.	D.k. at 600 nm; CO_3H and CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; k is dependent on pH, $k \sim 3 \times 10^7$ at pH 7, values from graph.	75-7110

TABLE 1. Rates of reaction of CO_3^- in aqueous solution^a—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
1.69	Methanol	5×10^3	12.5	—	f.phot.	D.k. at 600 nm in air-satd. soln. contg. 0.5 M Na_2CO_3 ; CO_3^- generated from $\text{SO}_4^{2-} + \text{CO}_3^{2-} \rightarrow \text{SO}_4^{2-} + \text{CO}_3^-$.	72-7383
		$\leq 2.6 \times 10^3$	6.4	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7109
1.70	Methionine	$(1.2 \pm 0.1) \times 10^8$	11.2 ± 0.3	0.3	p.r.	D.k. in N_2O -satd. soln.	72-0036
		$(3.6 \pm 0.4) \times 10^7$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
		$(4.8 \pm 0.5) \times 10^7$	11.0	0.003	f.phot.	D.k. at 600 nm; CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; slight pH effect.	75-7110
1.71	<i>p</i> -Methoxyphenoxide ion	5.2×10^8	12.2	0.3	p.r.	D.k. at 570 nm in N_2O -satd. carbonate soln.	77-1098
1.71a	Methyl radical $\text{CO}_3^- + \text{CH}_3 \rightarrow \text{CH}_3\text{OCO}_2^-$	$(3 \pm 1) \times 10^9$	—	—	p.r.	D.k. at 600 nm in Na_2CO_3 - CH_3Cl soln.	78-1506
1.71b	<i>N</i> -Methylaniline	1.8×10^9	—	—	p.r.	D.k. at 600 nm.	78-1506
1.72	<i>S</i> -Methylcysteine	$(2.5 \pm 0.2) \times 10^7$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
		$(5 \pm 0.5) \times 10^7$	11.0	0.003	f.phot.	D.k. at 600 nm; CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; slight pH effect.	75-7110
1.73	<i>p</i> -Methylphenoxide ion	4.8×10^8	12.2	0.3	p.r.	D.k. at 570 nm in N_2O -satd. carbonate soln.	77-1098
1.74	2-Methyl-2-propanol (<i>tert</i> -Butanol)	$\leq 1.6 \times 10^2$	6.4	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7109
1.75	<i>N</i> -Methyltryptophan	$(4.3 \pm 0.4) \times 10^8$	7-11	0.003	f.phot.	D.k. at 600 nm; CO_3H and CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$.	74-7296
1.76	Nitromethane	$(1 \pm 0.1) \times 10^6$	7	0.003	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	75-7110
1.77	<i>aci</i> -Nitromethane anion $\text{CO}_3^- + \text{CH}_2=\text{NO}_2^- \rightarrow \text{O}_2\text{COCH}_2\text{NO}_2^-$	$(1.5 \pm 0.4) \times 10^7$	12	0.003	f.phot.	D.k. at 600 nm; CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; product obs. by esr (72-5050).	75-7110
1.78	<i>p</i> -Nitrophenoxide ion	4.8×10^7	12.2	0.3	p.r.	D.k. at 570 nm in N_2O -satd. carbonate soln.	77-1098
1.79	Norpseudopelletierine- <i>N</i> -oxyl (9-Azabicyclo[3,3,1]nonan-3-one-9-oxyl)	$(1.1 \pm 0.1) \times 10^9$	—	0.015	p.r.	D.k. at 600 nm in N_2O -satd. carbonate soln.	71-0061
1.80	Penicillamine (3-Mercaptovaline)	2×10^7 2.4×10^8 1.2×10^8	4 10 12	0.003	f.phot.	D.k. at 600 nm; CO_3H or CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; values from graph.	75-7110
1.81	Phenol	$(2.2 \pm 0.2) \times 10^7$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
		4.9×10^6	7.0	0.06	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.; k increases with pH to 5×10^7 at pH 12.	75-7313
1.82	Phenoxide ion	2.4×10^8	12.2	0.3	p.r.	D.k. at 570 nm in N_2O -satd. carbonate soln.	77-1098
		3.3×10^8	—	—	p.r.	D.k. at 600 nm.	78-1506

TABLE 1. Rates of reaction of CO_3^- in aqueous solution^a—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
1.83	Phenylalanine	$< 1 \times 10^6$	11.2±0.3	0.3	p.r.	D.k. in N_2O -satd. soln.	72-0036
		$(5 \pm 0.5) \times 10^4$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.84	Phenylalanylglycine	$(4.0 \pm 0.4) \times 10^5$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.85	1-Propanol	1.9×10^4	12.1–12.7	–	f.phot.	D.k. at 600 nm in air-satd. soln. contg. 0.05 M $\text{Na}_2\text{S}_2\text{O}_8$ and 0.5 M Na_2CO_3 ; CO_3^- generated from $\text{SO}_4^- + \text{CO}_3^{2-}$.	72-7383
1.86	2-Propanol $\text{CO}_3^- + (\text{CH}_3)_2\text{CHOH} \rightarrow \text{HCO}_3^- + (\text{CH}_3)_2\text{COH}$	3.9×10^4	12.1–12.7	–	f.phot.	D.k. at 600 nm in air-satd. soln. contg. 0.05 M $\text{Na}_2\text{S}_2\text{O}_8$ and 0.5 M Na_2CO_3 ; CO_3^- generated from $\text{SO}_4^- + \text{CO}_3^{2-}$.	72-7383
		$\leq 4 \times 10^4$	6.4	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7109
		5×10^4	–	–	p.r.	D.k. at 600 nm.	78-1506
1.87	Ribonuclease A	$(5.0 \pm 0.5) \times 10^7$	7.0	0.03	f.phot.	D.k. at 600 nm; CO_3H or CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; k is pH dependent; value from graph; mol. wt. 13,700.	73-7352,
		1.4×10^8	11.0	0.03	f.phot.		75-7110
1.88	2,2,6,6-Tetra-methyl-4-oxo-1-piperidinyloxy (TAN)	$(4.8 \pm 0.5) \times 10^8$	–	0.015	p.r.	D.k. at 600 nm in N_2O -satd. carbonate soln.	71-0061
		4.6×10^8	10–11	–	p.r.	D.k. at 600 nm in air-satd. soln.	71-0618
1.89	Thymine	$< 10^4$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.90	Toluene	4.3×10^4	7.0	0.06	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	75-7313
1.91	Trichloroacetate ion	$< 1.5 \times 10^2$	12.1–12.7	–	f.phot.	D.k. in air-satd. soln. contg. 0.5 M Na_2CO_3 and 0.05 M $\text{Na}_2\text{S}_2\text{O}_8$; CO_3^- generated from $\text{SO}_4^- + \text{CO}_3^{2-}$.	72-7383
1.92	Trypsin	$(6.8 \pm 0.7) \times 10^8$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; mol. wt. 23,800.	73-7352
1.93	Tryptamine	$(1.25 \pm 0.1) \times 10^9$	8	0.003	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; k is pH dependent, $k \approx 9 \times 10^8$ at pH 12; values from graph.	74-7296
1.94	Tryptophan	$(4.4 \pm 0.4) \times 10^8$	11.2±0.3	0.3	p.r.	D.k. in N_2O -satd. soln.	72-0036
		4.3×10^8	12	0.003	f.phot.	D.k. at 600 nm; CO_3H or CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; k is pH dependent, $k \approx 6.2 \times 10^8$ at pH 9 and 7×10^8 at pH 7; values from graph.	74-7296 73-7352
1.95	Tryptophan methyl ester	$(9.5 \pm 1) \times 10^8$	7	0.003	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.; k pH-dependent.	74-7296
1.96	Tryptophanamide	$(1.3 \pm 0.1) \times 10^9$	7	0.003	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.; k pH-dependent.	74-7296

TABLE 1. Rates of reaction of CO_3^- in aqueous solution^a—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
1.97	Tryptophanylglycine	$(7 \pm 0.7) \times 10^8$	6	0.003	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.; k pH-dependent.	74-7296
1.98	Tyrosine	$(2.9 \pm 0.3) \times 10^8$	11.2 ± 0.3	0.3	p.r.	D.k. in N_2O -satd. soln.	72-0036
		$(4.5 \pm 0.4) \times 10^7$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
		$(1.4 \pm 0.1) \times 10^8$	11	0.03	f.phot.	D.k. at 600 nm; CO_3^- generated from $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$; k pH dependent.	75-7110
1.99	Uracil	$< 10^4$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352
1.100	Urea	$< 10^3$	7.0	0.03	f.phot.	D.k. at 600 nm in $\text{Co}(\text{NH}_3)_4\text{CO}_3^+$ soln.	73-7352

^aThe transient is represented in the anionic form; $\text{p}K_a$ has been reported to be 9.6.

TABLE 2. Rates of reactions of SO_4^- in aqueous solution

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
2.1	$\text{SO}_4^- + \text{SO}_4^{2-} \rightarrow \text{S}_2\text{O}_8^{2-}$	$2k = 1.0 \times 10^9$	5.5	0.06	f.phot.	D.k. at 455 nm in aerated sulfate soln.; recalcd. for $\epsilon_{455} = 1100 M^{-1} \text{cm}^{-1}$.	67-7012
		$2k = (8.8 \pm 0.8) \times 10^8$	0.1	1	f.phot.	D.k. at 455 nm in aerated $\text{S}_2\text{O}_8^{2-}$ soln.; recalcd. for $\epsilon_{455} = 1100 M^{-1} \text{cm}^{-1}$.	67-7058
		$(7.5 \pm 0.25) \times 10^8$	1.0	0.13			
		$(9.5 \pm 1) \times 10^8$	4.8	0.03			
		$2k = (1.6 \pm 0.3) \times 10^9$	—	1 M H_2SO_4	f.phot.	D.k. in aerated $2.5 \times 10^{-4} M$ ceric sulfate; recalcd. for $\epsilon_{455} = 1100 M^{-1} \text{cm}^{-1}$.	67-7274
		$2k = 3.6 \times 10^9$	—	4 M H_2SO_4	p.r.	D.k.; used $\epsilon_{450} \cong 1000 M^{-1} \text{cm}^{-1}$ (66-0019).	73-1030
2.2	$\text{SO}_4^- + \text{AsO}_2^- \rightarrow \text{SO}_4^{2-} + \cdot\text{AsO}_2$	8.0×10^8 (rel.)	7-8	—	phot.	C.k. with fumarate ion in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^- adduct obs. by esr; rel. to $k(\text{SO}_4^- + \text{OH}^-) = 7.3 \times 10^7$.	73-5403
2.3	$\text{SO}_4^- + \text{Br}^- \rightarrow \text{SO}_4^{2-} + \text{Br}$	$(3.5 \pm 0.4) \times 10^9$	7	0.03	p.r.	D.k. at 450 nm in air-satd. $\text{S}_2\text{O}_8^{2-}$ soln. contg. 0.05 M <i>tert</i> -BuOH.	75-1069
		1.6×10^9 (rel.)	7-8	—	phot.	C.k. with fumarate ion in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^- adduct obs. by esr; rel. to $k(\text{SO}_4^- + \text{OH}^-) = 7.3 \times 10^7$.	73-5403
2.4	$\text{SO}_4^- + \text{HCO}_3^- \rightarrow \text{SO}_4^{2-} + \text{CO}_3^{2-} + \text{H}^+$	$(9.1 \pm 0.4) \times 10^6$	7.5-8.5	0.03	f.phot.	D.k. at 330 nm (SO_4^-) as well as p.b.k. at 600 nm (CO_3^{2-}) in aerated $10^{-2} M \text{S}_2\text{O}_8^{2-}$ soln.	67-7058
		9.3×10^6 (rel.)	7-8	—	phot.	C.k. with fumarate in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^- adduct obs. by esr; rel. to $k(\text{SO}_4^- + \text{OH}^-) = 7.3 \times 10^7$.	73-5403
2.5	$\text{SO}_4^- + \text{CN}^-$	8.0×10^7 (rel.)	7-8	—	phot.	C.k. with fumarate in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^- adduct obs. by esr; rel. to $k(\text{SO}_4^- + \text{OH}^-) = 7.3 \times 10^7$.	73-5403
2.6	$\text{SO}_4^- + \text{OCN}^-$	5.0×10^8 (rel.)	7-8	—	phot.	C.k. with fumarate in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^- adduct obs. by esr; rel. to $k(\text{SO}_4^- + \text{OH}^-) = 7.3 \times 10^7$.	73-5403
2.7	$\text{SO}_4^- + \text{SCN}^- \rightarrow \text{SO}_4^{2-} + \text{SCN}$	$(5.2 \pm 0.5) \times 10^9$	7	0.03	p.r.	D.k. at 450 nm in air-satd. $\text{S}_2\text{O}_8^{2-}$ soln. contg. 0.05 M <i>tert</i> -BuOH.	75-1069

TABLE 2. Rates of reactions of SO_4^- in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
2.8	$\text{SO}_4^- + \text{Ce}^{\text{III}} \rightarrow \text{SO}_4^{2-} + \text{Ce}^{\text{IV}}$	$(1.43 \pm 0.3) \times 10^8$	—	1 M H_2SO_4	f.phot.	D.k. at 455 nm; Ce^{III} produced from ceric sulfate.	67-7274
		1.3×10^8 (rel.)	—	4 M H_2SO_4	γ -r.	Air-satd. soln. contains Ce^{IV} , Ce^{III} and formic acid; ratios calcd. from assumed mechanism: $k(\text{SO}_4^- + \text{Ce}^{\text{III}})/k(\text{SO}_4^- + \text{HCOOH}) \cong 160$; $k(\text{SO}_4^- + \text{Ce}^{\text{III}})/k(\text{SO}_4^- + 2\text{-PrOH}) = 1.6$; rel. to $k(\text{SO}_4^- + 2\text{-PrOH}) = 8 \times 10^7$.	72-0094
2.9	$\text{SO}_4^- + \text{Cl}^- \rightarrow \text{SO}_4^{2-} + \text{Cl}$	1.9×10^8 (rel.)	7-8	—	phot.	C.k. with fumarate ion in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^- adduct obs. by esr; rel. to $k(\text{SO}_4^- + \text{OH}^-) = 7.3 \times 10^7$.	73-5403
		3.1×10^8	6.8	—	p.r.	D.k. at 480 nm; soln. contains $\text{S}_2\text{O}_8^{2-}$, <i>tert</i> -BuOH and phosphate buffer.	75-5244
		1.3×10^8	—	—	p.r.	P.b.k. (Cl_2^-) in 0.002 M $\text{S}_2\text{O}_8^{2-}$; $k = 4.1 \times 10^8$ in 2 M SO_4^{2-} soln.	76-1141
2.10	$\text{SO}_4^- + \text{Cr}^{2+}$	$> 10^9$	1.0	—	therm.	Estimated.	68-9084
2.11	$\text{SO}_4^- + \text{Fe}^{2+} \rightarrow \text{SO}_4^{2-} + \text{Fe}^{3+}$	9.9×10^8	—	1	p.r.	D.k. at 450 nm.	66-0019
2.12	$\text{SO}_4^- + \text{NH}_2\text{OH}$	8.5×10^8	8.2	—	p.r.	—	78-1075
2.13	$\text{SO}_4^- + \text{NH}_2\text{OH}^+$	1.5×10^7	4.1	—	p.r.	—	78-1075
2.14	$\text{SO}_4^- + \text{N}_2\text{H}_4$	8.1×10^8	9.5	—	p.r.	—	78-1075
2.15	$\text{SO}_4^- + \text{N}_2\text{H}_5^+$	2.1×10^8	4	—	p.r.	—	78-1075
2.16	$\text{SO}_4^- + \text{N}_3^- \rightarrow \text{SO}_4^{2-} + \cdot\text{N}_3$	7.4×10^8 (rel.)	7-8	—	phot.	C.k. with fumarate in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^- adduct obs. by esr; rel. to $k(\text{SO}_4^- + \text{OH}^-) = 7.3 \times 10^7$.	73-5403
2.17	$\text{SO}_4^- + \text{NO}_2^- \rightarrow \text{SO}_4^{2-} + \text{NO}_2$	$\sim 3 \times 10^9$	7	—	p.r.	—	78-1075
		8.8×10^8	7	—	p.r.	—	78-1075
2.18	$\text{SO}_4^- + \text{NO}_3^-$	1.4×10^6 (rel.)	9	—	phot.	C.k. with RNO in air-satd. $\text{S}_2\text{O}_8^{2-}$ soln.; rel. to $k(\text{SO}_4^- + \text{EtOH}) = 6.2 \times 10^7$.	70-7234
		2.1×10^6 (rel.)	7-8	—	phot.	C.k. with fumarate in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^- adduct obs. by esr; rel. to $k(\text{SO}_4^- + \text{OH}^-) = 7.3 \times 10^7$.	73-5403
2.19	$\text{SO}_4^- + \text{OH}^- \rightarrow \text{SO}_4^{2-} + \text{OH}$	4.6×10^7	> 11	—	p.r.	D.k. at 460 nm; soln. contains $\text{S}_2\text{O}_8^{2-}$.	69-0158
		$(6.5 \pm 1.0) \times 10^7$	alk.	—	p.r.	D.k. vs. OH^- concn. in O_2 -free $\text{S}_2\text{O}_8^{2-}$ soln.	72-7008
		$(8.3 \pm 0.8) \times 10^7$ (rel.)	> 11	0.06-0.08	p.r.	C.k. with tyrosine; effect of pH on formn. of tyrosine transient at 410 nm; rel. to $k(\text{SO}_4^- + \text{tyr}) = 3.2 \times 10^9$ at pH 7.	75-1069
		7.3×10^7	—	—	p.r.	D.k. at 450 nm; soln. contains $\text{S}_2\text{O}_8^{2-}$ and <i>tert</i> -BuOH.	75-5244

TABLE 2. Rates of reactions of SO_4^- in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
2.20	$\text{SO}_4^- + \text{H}_2\text{O} \rightarrow \text{HSO}_4^- + \text{OH}^-$	$10^3\text{--}10^4 s^{-1}$ $< 3 \times 10^3 s^{-1}$	— 7	— —	therm. p.r.	Estimated. Extrapolated from d.k. vs. OH^- concn. (see 2.19).	68-9084 72-7008
2.21	$\text{SO}_4^- + \text{H}_2\text{O}_2$	1.2×10^7	7	—	p.r.	—	78-1075
2.22	$\text{SO}_4^- + \text{H}_2\text{PO}_2^- \rightarrow \text{SO}_4^{2-} + \text{H}^+ + \text{HPO}_2^-$	3.6×10^6 (rel.)	7-8	—	phot.	C.k. with fumarate ion in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^- adduct obs. by esr; rel. to $k(\text{SO}_4^- + \text{OH}^-) = 7.3 \times 10^7$.	73-5403
2.23	$\text{SO}_4^- + \text{HPO}_3^{2-} \rightarrow \text{SO}_4^{2-} + \text{H}^+ + \text{PO}_3^{2-}$	1.8×10^8 4.3×10^7 (rel.)	7 7-8	— —	p.r. phot.	— C.k. with fumarate ion in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^- adduct obs. by esr; rel. to $k(\text{SO}_4^- + \text{OH}^-) = 7.3 \times 10^7$.	78-1075 73-5403
2.24	$\text{SO}_4^- + \text{H}_2\text{PO}_3^-$	6.2×10^7 1.6×10^7	8.3 4	— —	p.r. p.r.	— —	78-1075 78-1075
2.25	$\text{SO}_4^- + \text{H}_2\text{PO}_4^-$	$\leq 7 \times 10^4$	7	—	p.r.	—	78-1075
2.26	$\text{SO}_4^- + \text{HPO}_4^{2-}$	$(1.2 \pm 0.3) \times 10^6$	9	—	p.r.	—	78-1075
2.27	$\text{SO}_4^- + \text{SO}_3^{2-} \rightarrow \text{SO}_4^{2-} + \text{SO}_3^-$	$> 5.3 \times 10^8$	—	—	p.r.	Rate with HSO_3^- higher by at least a factor of 2.5.	72-7008
		2.6×10^8 (rel.)	7-8	—	phot.	C.k. with fumarate ion in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^- adduct obs. by esr; rel. to $k(\text{SO}_4^- + \text{OH}^-) = 7.3 \times 10^7$.	73-5403
2.28	$\text{SO}_4^- + \text{HSO}_5^- \rightarrow \text{HSO}_4^- + \text{SO}_5^-$	$< 10^5$	—	—	p.r.	D.k. at 450 nm.	77-1047
2.29	$\text{SO}_4^- + \text{SiO}_3^{2-} \rightarrow \text{SO}_4^{2-} + \text{SiO}_3^-$	2×10^7	—	—	f. phot.	—	70-7262
2.30	$\text{SO}_4^- + \text{Ti}^+$	$(1.70 \pm 0.2) \times 10^9$	—	1 M H_2SO_4	f.phot.	D.k. at 455 nm; soln. contains ceric sulfate- H_2SO_4 .	67-7274
2.31	Acetanilide $\text{SO}_4^- + \text{C}_6\text{H}_5\text{NHCOCH}_3 \rightarrow \text{SO}_4^{2-} + \text{C}_6\text{H}_5^+\text{NHCOCH}_3$	3.6×10^9	7	—	p.r.	D.k. at 450 nm; soln. contains 0.01-0.05 M $\text{S}_2\text{O}_8^{2-}$.	77-1001
2.32	Acetate ion $\text{SO}_4^- + \text{CH}_3\text{COO}^- \rightarrow \text{SO}_4^{2-} + \cdot\text{CH}_3 + \text{CO}_2$ (1) $\text{SO}_4^- + \text{CH}_3\text{COO}^- \rightarrow \text{HSO}_4^- + \cdot\text{CH}_2\text{COO}^-$ (2)	5.0×10^6	6.8	—	p.r.	D.k. at 450 nm; soln. contains $\text{S}_2\text{O}_8^{2-}$, <i>tert</i> -BuOH, and phosphate buffer; predominantly (1); CO_2 yield meas. by γ -r. (78-1505).	75-5244
		8.7×10^6 (rel.)	7-8	—	phot.	C.k. with fumarate ion in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^- adduct obs. by esr; rel. to $k(\text{SO}_4^- + \text{OH}^-) = 7.3 \times 10^7$; product ident. by esr.	73-5403
2.33	Acetic acid	$(8.8 \pm 0.2) \times 10^4$	—	1 M H_2SO_4	p.r.	D.k. at 455 nm; soln. contains ceric sulfate- H_2SO_4 .	67-7274
2.34	Acetophenone $\text{SO}_4^- + \text{C}_6\text{H}_5\text{COCH}_3 \rightarrow \text{SO}_4^{2-} + \text{C}_6\text{H}_5^+\text{COCH}_3$	3.1×10^8	7	0.03	p.r.	D.k. at 450 nm in $\sim 10^{-2}$ M $\text{S}_2\text{O}_8^{2-}$ soln.	77-1001
2.35	<i>p</i> -Acetylbenzoate ion $\text{SO}_4^- + \text{CH}_3\text{COC}_6\text{H}_4\text{COO}^- \rightarrow \text{SO}_4^{2-} + \text{CH}_3\text{COC}_6\text{H}_4^+\text{COO}^-$	2.0×10^8	7	0.03	p.r.	D.k. at 450 nm in $\sim 10^{-2}$ M $\text{S}_2\text{O}_8^{2-}$ soln.	77-1001

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
2.36	Acrylamide $SO_4^- + CH_2 = CHCONH_2$	1.6×10^8	7.3	-	p.r.	-	77-1503
2.37	Acrylate ion $SO_4^- + CH_2 = CHCOO^-$	1.1×10^8	6.5	-	p.r.	-	77-1503
2.38	Acrylonitrile $SO_4^- + CH_2 = CHCN(+H_2O) \rightarrow$ $CH_2OHCHCN + HSO_4^-$	1.7×10^8	~ 7	-	p.r.	Soln. contains $S_2O_8^{2-}$; product obs. by absorption spectrum.	69-0158
2.39	Alanine $SO_4^- + CH_3CHNH_3^+ COO^-$	8.1×10^7	7.4	-	p.r.	-	77-1503
		$(1.0 \pm 0.1) \times 10^7$	7	0.03	p.r.	D.k.	75-1069
2.40	Allyl alcohol $SO_4^- + CH_2 = CHCH_2OH$	1.5×10^9	6.8	-	p.r.	-	77-1503
2.41	Allyl cyanide $SO_4^- + CH_2 = CHCH_2CN$	1.1×10^9	7	-	p.r.	-	77-1503
2.42	Anisole $SO_4^- + C_6H_5OCH_3 \rightarrow$ $\cdot C_6H_5^+ OCH_3 + SO_4^{2-}$	$(4.9 \pm 0.5) \times 10^9$	-	-	p.r.	P.b.k.; product ident. by esr.	75-1171
2.43	Benzamide $SO_4^- + C_6H_5CONH_2 \rightarrow$ $SO_4^{2-} + \cdot C_6H_5^+ CONH_2$	1.9×10^8	7	0.03	p.r.	D.k. at 450 nm in $\sim 10^{-2} M S_2O_8^{2-}$ soln.	77-1001
2.44	Benzene $SO_4^- + C_6H_6 \rightarrow$ $\cdot C_6H_6^+ + SO_4^{2-}$ $\cdot C_6H_6^+ (+ H_2O) \rightarrow$ $\cdot C_6H_6OH + H^+$	8.0×10^8	~ 7	-	p.r.	Soln. contains $S_2O_8^{2-}$; final product ident. by absorption spectrum.	69-0158
		6.7×10^8 (rel.)	9	-	phot.	C.k. with RNO in air- satd. $S_2O_8^{2-}$ soln.; rel. to $k(SO_4^- + EtOH) =$ 6.2×10^7 .	70-7234
		$\sim 3 \times 10^9$	7	0.03	p.r.	D.k. at 450 nm in $S_2O_8^{2-}$ soln. contg. 0.1 M tert-BuOH; also p.b.k. at 315 nm (76-1187).	77-1001
		1,3,5-Benzenetricarboxylate ion 1,2,4,5-Benzenetetracarboxylate ion	See Trimesate ion. See Pyromellitate ion.				
2.45	Benzoate ion $SO_4^- + C_6H_5COO^- \rightarrow$ $SO_4^{2-} + \cdot C_6H_5^+ COO^-$ $C_6H_5^+ COO^- (+ H_2O) \rightarrow$ $HOC_6H_5COO^- + H^+(1)$ $C_6H_5^+ COO^- \rightarrow$ $\cdot C_6H_5 + CO_2(2)$	1.2×10^9	7	0.03	p.r.	D.k. at 450 nm in $\sim 10^{-2} M S_2O_8^{2-}$ soln.; CO_2 yield in γ -r. suggests reaction (2) contributes 56% (78-1505). Absorption of OH adduct in p.r. suggests reaction (1) contributes 20% (78-1504).	77-1001
2.46	Benzonitrile $SO_4^- + C_6H_5CN \rightarrow$ $SO_4^{2-} + \cdot C_6H_5^+ CN$	1.2×10^8	7	0.03	p.r.	D.k. at 450 nm in $\sim 10^{-2} M S_2O_8^{2-}$ soln.	77-1001
2.47	Benzyl methyl ether $SO_4^- + C_6H_5CH_2OCH_3$	$k/k_2-PrOH = 18$	1.8	-	therm.	$T = 75^\circ C$; c.k. in $S_2O_8^{2-}$ -alcohol soln.; obs. C_6H_5CHO yield.	74-9006
2.48	o-Bromobenzoate ion $SO_4^- + BrC_6H_4COO^- \rightarrow$ $SO_4^{2-} + BrC_6H_4^+ COO^-$ $BrC_6H_4^+ COO^- (+ H_2O) \rightarrow$ $\cdot OC_6H_4COO^- + 2H^+ +$ $Br^- (1)$ $BrC_6H_4^+ COO^- (+ H_2O) \rightarrow$ $BrC_6H_4(OH)COO^- + H^+ (2)$ $BrC_6H_4^+ COO^- \rightarrow$ $\cdot C_6H_4Br + CO_2(3)$	8.7×10^8	7	0.03	p.r.	D.k. at 450 nm in $\sim 10^{-2} M S_2O_8^{2-}$ soln.; Br^- and CO_2 yields meas. on γ -r. indicate reaction (1) contributes $\sim 16\%$ and reaction (3) $\sim 10\%$ (78-1505).	77-1001

TABLE 2. Rates of reactions of SO_4^- in aqueous solution—Continued

No.	Reaction	$k(M^{-1} \text{ s}^{-1})$	pH	I	Method	Comment	Ref.
2.49	<i>p</i> -Bromobenzoate ion <i>tert</i> -Butanol See 2-Methyl-2-propanol.	1.0×10^9	7	0.03	p.r.	D.k. at 450 nm in $\sim 10^{-2} M \text{ S}_2\text{O}_8^{2-}$ soln.; Br^- and CO_2 yields meas. on γ -r. indicate $\sim 30\%$ debromination and $\sim 40\%$ decarboxylation; see reactions under <i>o</i> -bromobenzoate (78-1505).	77-1001
2.50	<i>p</i> -Chlorobenzoate ion	3.6×10^8	7	0.03	p.r.	D.k. at 450 nm in $\sim 10^{-2} M \text{ S}_2\text{O}_8^{2-}$ soln.; CO_2 yield meas. on γ -r. indicates $\sim 40\%$ decarboxylation (78-1505).	77-1001
2.51	Crotonic acid Crotononitrile See Allyl cyanide	7.7×10^8	4.8	—	p.r.	—	77-1503
2.52	<i>p</i> -Cyanobenzoate ion	3.3×10^7	4.8	0.03	p.r.	D.k. at 450 nm in $\sim 10^{-2} M \text{ S}_2\text{O}_8^{2-}$ soln.; CO_2 yield meas. on γ -r. indicates $\sim 30\%$ decarboxylation (78-1505).	77-1001
2.53	Cycloheptanol	$k/k_{2-\text{proH}} = 2.4$	1.8	—	therm.	$T = 75^\circ\text{C}$; c.k. in $\text{S}_2\text{O}_8^{2-}$ -alcohol soln.; obs. ketone formn.; for 1- <i>d</i> -cycloheptanol ratio = 1.5.	74-9006
2.54	Cyclohexene $\text{SO}_4^- + \text{C}_6\text{H}_{10} (+\text{H}_2\text{O}) \rightarrow \text{C}_6\text{H}_{10}\text{OH} + \text{HSO}_4^-$	4.1×10^8	~ 7	—	p.r.	Soln. contains $\text{S}_2\text{O}_8^{2-}$; product obs. by absorption spectrum.	69-0158
2.55	Diethyl disulfide $\text{SO}_4^- + \text{RSSR} \rightarrow \text{SO}_4^{2-} + (\text{RSSR})^+$	2.6×10^8	~ 4.5	0.3	p.r.	D.k. at 300 nm in Ar-satd. 0.1 $M \text{ S}_2\text{O}_8^{2-}$ soln.	76-1143
2.56	1,2-Dimethoxybenzene $\text{SO}_4^- + \text{DMB} \rightarrow \text{SO}_4^{2-} + (\text{DMB})^+$	$(5.0 \pm 0.5) \times 10^9$	—	0.003	p.r.	P.b.k.; product ident. by esr and absorption spectra.	75-1171
2.57	1,3-Dimethoxybenzene $\text{SO}_4^- + \text{DMB} \rightarrow \text{SO}_4^{2-} + (\text{DMB})^+$	$(7.0 \pm 0.7) \times 10^9$	—	0.003	p.r.	P.b.k.; product ident. by esr and absorption spectra.	75-1171
2.58	1,4-Dimethoxybenzene $\text{SO}_4^- + \text{DMB} \rightarrow \text{SO}_4^{2-} + (\text{DMB})^+$	$(7.2 \pm 0.7) \times 10^9$	—	0.003	p.r.	P.b.k.; product ident. by esr and absorption spectra.	75-1171
2.59	2,3-Dimethoxybenzoate ion	8.5×10^9	> 3	0.03	p.r.	P.b.k. in Ar-satd. $\text{S}_2\text{O}_8^{2-}$ soln.	77-1006
2.60	2,4-Dimethoxybenzoate ion	3.8×10^9	> 3	0.03	p.r.	P.b.k. in Ar-satd. $\text{S}_2\text{O}_8^{2-}$ soln.	77-1006
2.61	3,4-Dimethoxybenzoate ion	4.5×10^9	> 3	0.03	p.r.	P.b.k. in Ar-satd. $\text{S}_2\text{O}_8^{2-}$ soln.	77-1006
2.62	2,6-Dimethoxybenzoate ion	2.5×10^9	> 3	0.03	p.r.	P.b.k. in Ar-satd. $\text{S}_2\text{O}_8^{2-}$ soln.	77-1006
2.63	3,5-Dimethoxybenzoate ion	4.4×10^9	> 3	0.03	p.r.	P.b.k. in Ar-satd. $\text{S}_2\text{O}_8^{2-}$ soln.	77-1006
2.64	Dimethyl disulfide $\text{SO}_4^- + \text{RSSR} \rightarrow \text{SO}_4^{2-} + (\text{RSSR})^+$	3.8×10^8	~ 4.5	0.3	p.r.	D.k. at 300 nm in Ar-satd. 0.1 $M \text{ S}_2\text{O}_8^{2-}$ soln.	76-1143

TABLE 2. Rates of reactions of $\text{SO}_4^{\cdot -}$ in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
2.64a	Dioxane $\text{SO}_4^{\cdot -} + \text{C}_4\text{H}_8\text{O}_2 \rightarrow$ $\text{HSO}_4^- + \text{C}_4\text{H}_7\text{O}_2$	1.6×10^7	7–8	0.003	p.r.	C.k.; rel. to $k(\text{SO}_4^{\cdot -} + \text{TMB}) = 2.4 \times 10^9$; obs. buildup of TMB^+ (see 2.108); $k_H/k_D = 1.7$.	78–1076
2.65	Ethanol $\text{SO}_4^{\cdot -} + \text{CH}_3\text{CH}_2\text{OH} \rightarrow$ $\text{HSO}_4^- + \text{CH}_3\text{CHOH}$	3×10^7 $(7.7 \pm 2.2) \times 10^7$ $(6.2 \pm 1.4) \times 10^7$ $(3.4 \pm 0.3) \times 10^7$	– 4.8 1.0 –	1 0.03 1 $1 M$ H_2SO_4	p.r. f.phot. f.phot. p.r.	D.k. at 450 nm; HSO_4^- soln. D.k. in aerated $10^{-2} M \text{K}_2\text{S}_2\text{O}_8$ soln. D.k. at 455 nm in ceric sulfate– H_2SO_4 soln. C.k., rel. to $k(\text{SO}_4^{\cdot -} + \text{TMB}) = 2.4 \times 10^9$; obs. buildup of TMB^+ (see 2.108); $k_H/k_D = 2.4$.	66–0019 67–7058 67–7274 78–1076
2.66	1-(<i>p</i> -Ethyl- phenyl)ethanol	$k/k_{2-\text{PrOH}} = 26$	1.8	–	therm.	$T = 75^\circ\text{C}$; c.k. in $\text{S}_2\text{O}_8^{2-}$ alcohol soln.; obs. ketone formation.	74–9006
2.67	Formate ion $\text{SO}_4^{\cdot -} + \text{HCOO}^- \rightarrow$ $\text{SO}_4^{2-} + \text{CO}_2^-$	$(1.7 \pm 0.2) \times 10^8$ 2.7×10^8	7 7–8	0.03 –	p.r. phot.	D.k.; $\text{S}_2\text{O}_8^{2-}$ soln. C.k. with fumarate ion in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate– $\text{SO}_4^{\cdot -}$ adduct obs. by esr; rel. to $k(\text{SO}_4^{\cdot -} + \text{OH}^-) = 7.3 \times 10^7$.	75–1069 73–5403
2.68	Formic acid	$(1.35 \pm 0.2) \times 10^6$	–	$1 M$ H_2SO_4	f.phot.	D.k. at 455 nm in ceric sulfate– H_2SO_4 soln.	67–7274
2.69	Fumarate ion $\text{SO}_4^{\cdot -} + ^-\text{OOCCH}=\text{CHCOO}^- \rightarrow$ $^-\text{OOCCHCH}(\text{OSO}_3^-)\text{COO}^- (1)$ $\text{SO}_4^{\cdot -} + ^-\text{OOCCH}=\text{CHCOO}^- \rightarrow$ $\text{SO}_4^{2-} + ^-\text{OOCCH}=\text{CH}\cdot + \text{CO}_2 (2)$	1.6×10^7	7	–	p.r.	D.k. at 450 nm; $k_1/k_2 \cong 1$ from $G(\text{CO}_2)$ by γ -r. (78–1505); adduct obs. by esr (75–5244).	77–1106
2.70	Glycine	$(9 \pm 0.9) \times 10^6$	7	–	p.r.	D.k. in $\text{S}_2\text{O}_8^{2-}$ soln.	75–1069
2.71	Histidine	$\sim 2.5 \times 10^9$	7	0.03	p.r.	D.k. in $\text{S}_2\text{O}_8^{2-}$ soln.	75–1069
2.72	Homophthalate ion $\text{SO}_4^{\cdot -} + \text{ArCH}_2\text{COO}^- \rightarrow$ $\text{SO}_4^{2-} + \text{ArCH}_2\cdot + \text{CO}_2$	$(1.1 \pm 0.2) \times 10^9$	7	–	p.r.	D.k. at 450 nm as well as p.b.k. at 328 nm ($\text{ArCH}_2\cdot$) in $\text{S}_2\text{O}_8^{2-}$ soln.; CO_2 yield meas. by γ -r. (78–1505) and absorption of $\text{ArCH}_2\cdot$ radical by p.r. (78–1504) indicate predominant decarboxylation from side chain.	76–1187
2.73	<i>p</i> -Hydroxybenzoate ion $\text{SO}_4^{\cdot -} + \text{HOC}_6\text{H}_4\text{COO}^- \rightarrow$ $\text{SO}_4^{2-} + ^\cdot\text{OC}_6\text{H}_4\text{COO}^- + \text{H}^+$	2.5×10^9	7	–	p.r.	D.k. in $\text{S}_2\text{O}_8^{2-}$ soln.	77–1001
2.74	Lysozyme	1.8×10^{10}	7	0.03	p.r.	D.k. in $\text{S}_2\text{O}_8^{2-}$ – <i>tert</i> -BuOH soln.	75–1069
2.75	Malonate ion $\text{SO}_4^{\cdot -} + ^-\text{OOCCH}_2\text{COO}^- \rightarrow$ $\text{SO}_4^{2-} + ^-\text{OOCCH}_2\cdot + \text{CO}_2$	5.5×10^6	7	–	p.r.	D.k. at 450 nm; radical obs. by esr; CO_2 yield meas. by γ -r. (78–1505).	77–1106
2.76	Methacrylic acid	1.1×10^9 7×10^8	2.9 6.9	– –	p.r. p.r.	–	77–1503
2.77	Methacrylonitrile	3.8×10^8	7	–	p.r.	–	77–1503

TABLE 2. Rates of reactions of SO_4^- in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
2.78	Methanol	2×10^7	—	1	p.r.	D.k. at 450 nm.	66-0019
	$\text{SO}_4^- + \text{CH}_3\text{OH} \rightarrow$	$(2.5 \pm 0.4) \times 10^7$	4.8	0.03	f.phot.	D.k. in aerated $10^{-2} M$	67-7058
	$\text{HSO}_4^- + \text{CH}_2\text{OH}$	$(2.5 \pm 0.6) \times 10^7$	1.0	—		$\text{K}_2\text{S}_2\text{O}_8$ soln.	
		$(1.1 \pm 0.2) \times 10^7$	—	1 M	f.phot.	D.k. at 455 nm in ceric sulfate- H_2SO_4 soln.	67-7274
		7.6×10^6	9	—	phot.	C.k. with RNO in air-satd. $\text{S}_2\text{O}_8^{2-}$ soln.; rel. to $k(\text{SO}_4^- + \text{EtOH}) = 6.2 \times 10^7$.	70-7234
		(rel.)					
		1.3×10^7	7-8	—	phot.	C.k. with fumarate in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^- adduct obs. by esr; rel. to $k(\text{SO}_4^- + \text{OH}^-) = 7.3 \times 10^7$.	73-5403
		(rel.)					
		$(1.1 \pm 0.1) \times 10^7$	7	0.03	p.r.	D.k. in $\text{S}_2\text{O}_8^{2-}$ soln.	75-1069
		3.2×10^6	7-8	0.003	p.r.	C.k.; rel. to $k(\text{SO}_4^- + \text{TMB}) = 2.4 \times 10^9$; obs. buildup of TMB^+ (see 2.108); $k_H/k_D = 2.7$. D.k. in $\text{S}_2\text{O}_8^{2-}$ soln.	78-1076
2.79	Methionine	1.1×10^9	7	0.03	p.r.		75-1069
	Methoxybenzene	See Anisole					
2.80	2-Methoxybenzoate ion	7.0×10^9	> 3	0.03	p.r.	P.b.k. in Ar-satd. $\text{S}_2\text{O}_8^{2-}$ soln.	77-1006
2.81	3-Methoxybenzoate ion	7.6×10^9	> 3	0.03	p.r.	P.b.k. in Ar-satd. $\text{S}_2\text{O}_8^{2-}$ soln.	77-1006
2.82	4-Methoxybenzoate ion	3.5×10^9	7	—	p.r.	P.b.k. at 560 nm (cation radical) in $\text{S}_2\text{O}_8^{2-}$ soln.	77-1001
	$\text{SO}_4^- + \text{CH}_3\text{OC}_6\text{H}_4\text{COO}^- \rightarrow$						
	$\text{SO}_4^{2-} + \text{CH}_3\text{OC}_6\text{H}_4^+\text{COO}^-$	7.6×10^9	> 3	0.03	p.r.	P.b.k. in Ar-satd. $\text{S}_2\text{O}_8^{2-}$ soln.	77-1006
2.83	1-Methoxy-2-methyl-1-phenylpropane	$k/k_{2-\text{PrOH}} = 14$	1.8	—	therm.	$T = 75^\circ\text{C}$; c.k. in $\text{S}_2\text{O}_8^{2-}$ -alcohol soln.; obs. benzaldehyde and isopropyl phenyl ketone formn.	74-9006
	$\text{SO}_4^- + \text{C}_6\text{H}_5\text{CH}(\text{OCH}_3)\text{CH}(\text{CH}_3)_2$						
2.84	Methyl methacrylate	1.0×10^9	7.3	—	p.r.	—	77-1503
	$\text{SO}_4^- + \text{CH}_2=\text{C}(\text{CH}_3)\text{COOCH}_3$						
2.85	2-Methyl-2-propanol	$(9.1 \pm 1.0) \times 10^5$	—	—	p.r.	D.k. in $\text{S}_2\text{O}_8^{2-}$ soln.	72-7008
	$\text{SO}_4^- + (\text{CH}_3)_3\text{COH} \rightarrow$	$k/k_{2-\text{PrOH}} = 0.05$	1.8	—	therm.	$T = 75^\circ\text{C}$; c.k. in $\text{S}_2\text{O}_8^{2-}$ -alcohol soln.; obs. acetone formn.	74-9006
	$\text{HSO}_4^- + \text{CH}_2\text{C}(\text{CH}_3)_2\text{OH}$						
		$(8.0 \pm 0.8) \times 10^5$	7	0.03	p.r.	D.k. in $\text{S}_2\text{O}_8^{2-}$ soln.	75-1069
		4.0×10^5	7-8	0.003	p.r.	C.k.; rel. to $k(\text{SO}_4^- + \text{TMB}) = 2.4 \times 10^9$; obs. buildup of TMB^+ (see 2.108).	78-1076
2.86	<i>p</i> -Nitroso- <i>N,N</i> -dimethylaniline	2.3×10^9	7	—	phot.	Air-satd. $\text{S}_2\text{O}_8^{2-}$ soln.; rel. to $k(\text{SO}_4^- + \text{EtOH}) = 6.2 \times 10^7$.	70-7234
	$\text{SO}_4^- + \text{NOC}_6\text{H}_4\text{N}(\text{CH}_3)_2$	(rel.)					
2.87	3-Pentanol	$k/k_{2-\text{PrOH}} = 1.3$	1.8	—	therm.	$T = 75^\circ\text{C}$; c.k. in $\text{S}_2\text{O}_8^{2-}$ -alcohol soln.; obs. ketone formn.	74-9006
	$\text{SO}_4^- + (\text{C}_2\text{H}_5)_2\text{CHOH}$						
2.88	1-Phenyl-3-butanol	$k/k_{2-\text{PrOH}} = 25$	1.8	—	therm.	$T = 75^\circ\text{C}$; c.k. in $\text{S}_2\text{O}_8^{2-}$ -alcohol soln.; obs. ketone and acetaldehyde formn.	74-9006
	$\text{SO}_4^- + \text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{CHOHCH}_3$						

TABLE 2. Rates of reactions of SO_4^{2-} in aqueous solution—Continued

No.	Reaction	$k(M^{-1} \text{ s}^{-1})$	pH	I	Method	Comment	Ref.
2.89	1-Phenylethanol $\text{SO}_4^{2-} + \text{C}_6\text{H}_5\text{CHOHCH}_3$	$k/k_{2-\text{PrOH}} = 22$	1.8	—	therm.	$T = 75^\circ\text{C}$; c.k. in $\text{S}_2\text{O}_8^{2-}$ -alcohol soln.; obs. ketone formn.	74-9006
2.90	1-Phenyl-2-propanol $\text{SO}_4^{2-} + \text{C}_6\text{H}_5\text{CH}_2\text{CHOHCH}_3$	$k/k_{2-\text{PrOH}} = 42$	1.8	—	therm.	$T = 75^\circ\text{C}$; c.k. in $\text{S}_2\text{O}_8^{2-}$ -alcohol soln.; obs. acetaldehyde formn.	74-9006
2.91	2-Phenyl-2-propanol $\text{SO}_4^{2-} + \text{C}_6\text{H}_5\text{COH}(\text{CH}_3)_2$	$k/k_{2-\text{PrOH}} = 8.4$	1.8	—	therm.	$T = 75^\circ\text{C}$; c.k. in $\text{S}_2\text{O}_8^{2-}$ -alcohol soln.; obs. ketone formn.	74-9006
2.92	1-Propanol	4.8×10^7 (rel.)	9	—	phot.	C.k. with RNO in air-satd. $\text{S}_2\text{O}_8^{2-}$ soln.; rel. to $k(\text{SO}_4^{2-} + \text{EtOH}) = 6.2 \times 10^7$.	70-7234
		5.8×10^7	7-8	—	phot.	C.k. with fumarate in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^{2-} adduct obs. by esr; rel. to $k(\text{SO}_4^{2-} + \text{OH}^-) = 7.3 \times 10^7$.	73-5403
2.93	2-Propanol	$(8.5 \pm 3.0) \times 10^7$	4.4	—	f.phot.	D.k. in aerated $10^{-2} M$ $\text{K}_2\text{S}_2\text{O}_8$ soln.	67-7058
		$(9.1 \pm 2.8) \times 10^7$	1.0	—	f.phot.	D.k. at 455 nm in ceric sulfate- H_2SO_4 soln.	67-7274
		$(4.6 \pm 0.2) \times 10^7$	—	$1 M$ H_2SO_4	f.phot.	D.k. at 455 nm in ceric sulfate- H_2SO_4 soln.	67-7274
		8.5×10^7 (rel.)	9	—	phot.	C.k. with RNO in air-satd. $\text{S}_2\text{O}_8^{2-}$ soln.; rel. to $k(\text{SO}_4^{2-} + \text{EtOH}) = 6.2 \times 10^7$.	70-7234
		4.0×10^7	7-8	—	phot.	C.k. with fumarate in $\text{S}_2\text{O}_8^{2-}$ soln.; effect of solute on fumarate- SO_4^{2-} adduct obs. by esr; rel. to $k(\text{SO}_4^{2-} + \text{OH}^-) = 7.3 \times 10^7$.	73-5403
		8.0×10^7	7	0.03	p.r.	D.k. at 450 nm; $\text{S}_2\text{O}_8^{2-}$ - <i>tert</i> -BuOH soln.	75-1069
2.94	Propionate ion $\text{SO}_4^{2-} + \text{CH}_3\text{CH}_2\text{COO}^- \rightarrow$ $\text{SO}_4^{2-} + \text{CH}_3\text{CH}_2 + \text{CO}_2$	4.6×10^6	7	—	p.r.	D.k. at 450 nm; CO_2 yield obs. by γ -r. (78-1505).	77-1106
			7	—	p.r.	D.k. at 450 nm; $\text{S}_2\text{O}_8^{2-}$ soln.	76-1187
			3.9	—	p.r.	D.k. at 450 nm; $\text{S}_2\text{O}_8^{2-}$ soln.	76-1187
2.95	Pyridine $\text{SO}_4^{2-} + \text{C}_5\text{H}_5\text{N}$	$(2.2 \pm 0.2) \times 10^8$	7	—	p.r.	D.k. at 450 nm; $\text{S}_2\text{O}_8^{2-}$ soln.	76-1187
2.96	Pyridinium ion $\text{SO}_4^{2-} + \text{C}_5\text{H}_5\text{NH}^+$	$(2.0 \pm 0.4) \times 10^7$	3.9	—	p.r.	D.k. at 450 nm; $\text{S}_2\text{O}_8^{2-}$ soln.	76-1187
2.97	Pyromellitate ion (1,2,4,5-Benzenetetracarboxylate ion)	$(1.7 \pm 0.2) \times 10^7$	9	—	p.r.	D.k. at 450 nm; $\text{S}_2\text{O}_8^{2-}$ soln.	76-1187
2.98	Ribonuclease	1.2×10^{10}	7	0.03	p.r.	D.k. at 450 nm; $\text{S}_2\text{O}_8^{2-}$ - <i>tert</i> -BuOH soln.	75-1069
2.99	Serine $\text{SO}_4^{2-} + \text{HOCH}_2\text{CHNH}_3^+\text{COO}^-$	2.3×10^7	7	0.03	p.r.	D.k. at 450 nm; $\text{S}_2\text{O}_8^{2-}$ - <i>tert</i> -BuOH soln.	75-1069
2.100	Succinate ion $\text{SO}_4^{2-} + ^-\text{OOCCH}_2\text{CH}_2\text{COO}^- \rightarrow$ $\text{SO}_4^{2-} + ^-\text{OOCCH}_2\text{CH}_2 + \text{CO}_2$	7.1×10^6	7	—	p.r.	D.k. at 450 nm; CO_2 yield obs. by γ -r. (78-1505).	77-1106

TABLE 2. Rates of reactions of $\text{SO}_4^{\cdot -}$ in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
2.101	Terephthalate ion $\text{SO}_4^{\cdot -} + \text{C}_6\text{H}_4(\text{COO}^-)_2 \rightarrow$ $\text{SO}_4^{2-} + \text{C}_6\text{H}_4\text{COO}^- + \text{CO}_2$ (1) $\text{SO}_4^{\cdot -} + \text{C}_6\text{H}_4(\text{COO}^-)_2 (+\text{OH}^-) \rightarrow$ $\text{SO}_4^{2-} + \text{HOC}_6\text{H}_4(\text{COO}^-)_2$ (2)	1.7×10^8	9	—	p.r.	D.k. at 450 nm; $\sim 60\%$ (1) and $\sim 40\%$ (2) based on CO_2 yields by γ -r. (78-1505).	77-1001
2.101a	Tetrahydrofuran $\text{SO}_4^{\cdot -} + \text{C}_4\text{H}_8\text{O} \rightarrow$ $\text{HSO}_4^- + \text{C}_4\text{H}_7\text{O}$	1.0×10^8	7-8	0.003	p.r.	C.k.; rel. to $k(\text{SO}_4^{\cdot -} + \text{TMB}) = 2.4 \times 10^9$; obs. buildup of TMB^+ (see 2.108); $k_H/k_D = 2.0$.	78-1076
2.102	<i>o</i> -Toluate ion $\text{SO}_4^{\cdot -} + \text{CH}_3\text{C}_6\text{H}_4\text{COO}^- \rightarrow$ $\text{HSO}_4^- + \cdot\text{CH}_2\text{C}_6\text{H}_4\text{COO}^-$	1.4×10^9	7	—	p.r.	D.k. at 450 nm; product ident. by abs. spectrum by p.r. (78-1504); CO_2 yield by γ -r. $< 10\%$ (78-1505).	77-1001
2.103	<i>m</i> -Toluate ion	2.0×10^9	7	—	p.r.	D.k. at 450 nm; abs. spectrum by p.r. indicates predominant formation of OH adducts (78-1504).	77-1001
2.104	<i>p</i> -Toluate ion $\text{SO}_4^{\cdot -} + \text{CH}_3\text{C}_6\text{H}_4\text{COO}^- \rightarrow$ $\text{SO}_4^{2-} + \text{CH}_3\text{C}_6\text{H}_4 + \text{CO}_2$ (1) $\text{SO}_4^{\cdot -} + \text{CH}_3\text{C}_6\text{H}_4\text{COO}^- \rightarrow$ $\text{SO}_4^{2-} + \text{CH}_2\text{C}_6\text{H}_4\text{COO}^-$ (2) $\text{SO}_4^{\cdot -} + \text{CH}_3\text{C}_6\text{H}_4\text{COO}^- (+\text{OH}^-) \rightarrow$ $\text{SO}_4^{2-} + \text{CH}_3\text{C}_6\text{H}_4(\text{OH})\text{COO}^-$ (3)	1.8×10^9	7	—	p.r.	D.k. at 450 nm; $\sim 30\%$ (1), 30% (2) and 40% (3) based on CO_2 yields by γ -r. (78-1505) and abs. spectra by p.r. (78-1504).	77-1001
2.105	Trimesate ion (1,3,5-Benzenetri-carboxylate ion)	$(8.3 \pm 0.8) \times 10^7$	9	—	p.r.	D.k. at 450 nm.	76-1187
2.106	1,2,3-Trimethoxybenzene	$\sim (6 \text{ to } 8) \times 10^9$	—	0.003	p.r.	P.b.k.; product ident. by optical absorption and esr spectra.	75-1171
2.107	1,2,4-Trimethoxybenzene	$(7.8 \pm 0.8) \times 10^8$	—	0.003	p.r.	P.b.k.; product ident. by optical absorption and esr spectra (see 2.108).	75-1171
2.108	1,3,5-Trimethoxybenzene (TMB) $\text{SO}_4^{\cdot -} + \text{TMB} \rightarrow$ $\text{SO}_4^{2-} + (\text{TMB})^{\cdot +}$	$(1.8 \pm 0.2) \times 10^9$ $(2.4 \pm 0.5) \times 10^9$	— 7-8	0.003 0.003	p.r. p.r.	P.b.k.; product ident. by optical absorption and esr spectra.	75-1171 78-1076
2.109	2,3,4-Trimethoxybenzoate ion	2.5×10^9	> 3	0.03	p.r.	P.b.k. in Ar-satd. $\text{S}_2\text{O}_8^{2-}$ soln.	77-1006
2.110	3,4,5-Trimethoxybenzoate ion	5.0×10^9	> 3	0.03	p.r.	P.b.k. in Ar-satd. $\text{S}_2\text{O}_8^{2-}$ soln.	77-1006
2.111	2,4,5-Trimethoxybenzoate ion	4.4×10^9	> 3	0.03	p.r.	P.b.k. in Ar-satd. $\text{S}_2\text{O}_8^{2-}$ soln.	77-1006
2.112	2,4,6-Trimethoxybenzoate ion	2.6×10^9	> 3	0.03	p.r.	P.b.k. in Ar-satd. $\text{S}_2\text{O}_8^{2-}$ soln.	77-1006
2.113	Trimethylanilinium ion $\text{SO}_4^{\cdot -} + \text{C}_6\text{H}_5\text{N}(\text{CH}_3)_3^+ \rightarrow$ $\text{SO}_4^{2-} + \cdot\text{C}_6\text{H}_5 + \text{N}(\text{CH}_3)_3^+$	1.5×10^8	7	—	p.r.	D.k. at 450 nm; $\text{S}_2\text{O}_8^{2-}$ soln.	77-1001
2.114	Tryptophan	$\sim 2.3 \times 10^9$	7	0.03	p.r.	D.k. at 450 nm; $\text{S}_2\text{O}_8^{2-}$ - <i>tert</i> -BuOH soln.	75-1069
2.115	Tyrosine $\text{SO}_4^{\cdot -} + \text{ArOH} \rightarrow$ $\text{SO}_4^{2-} + \text{ArOH}^{\cdot +}$ $\text{ArOH}^{\cdot +} \rightarrow \text{ArO}^{\cdot} + \text{H}^+$	3.2×10^9 $(3.0 \pm 0.2) \times 10^9$	7 6.8	0.03 —	p.r. p.r.	D.k. at 450 nm; $\text{S}_2\text{O}_8^{2-}$ - <i>tert</i> -BuOH soln. D.k. at 450 nm as well as p.b.k. at 407 nm (aryloxy radical).	75-1069 76-1112
2.116	Vinyl acetate $\text{SO}_4^{\cdot -} + \text{AcOCHCH}_2 (+\text{H}_2\text{O}) \rightarrow$ $\text{AcOCHCH}_2\text{OH} + \text{HSO}_4^-$	1.1×10^9	~ 7	—	p.r.	Soln. contains $\text{S}_2\text{O}_8^{2-}$; product obs. by absorption spectrum.	69-0158

TABLE 3. Rates of reaction of SO_2^- , SO_3^- , S_2O_3^- , S_2O_2^- , SO_5^- , SeO_2^- and SeO_3^- in aqueous solution

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
3.1	$2\text{SO}_2^- \rightarrow \text{S}_2\text{O}_4^{2-}$	$2k = (1.0 \pm 0.1) \times 10^{10}$	~ 4	$\rightarrow 0$	f.phot.	D.k. at 370 nm, as well as p.b.k. at 330 nm in 0.06 M HSO_3^- soln.; assumed $\epsilon(330 \text{ nm}) = 5700 M^{-1} \text{ cm}^{-1}$; $2k \leq 1.3 \times 10^{10}$ based on $2k/\epsilon(370) \cong 2.5 \times 10^7$.	72-7008
		$2k = (1.3 \pm 0.4) \times 10^9$	acid	-	p.r.	D.k. at 360 nm ($\epsilon = 600 M^{-1} \text{ cm}^{-1}$) as well as p.b.k. at 320 nm in $\text{SO}_2 + \text{HSO}_3^-$ soln.	74-1033
		$2k = (2.2 \pm 0.3) \times 10^9$	3.1	~ 1	p.r.	P.b.k. at 320 nm in 10^{-3} M $\text{HSO}_3^-/\text{SO}_2$ soln. contg. 1 M $\text{HCOO}^-/\text{HCOOH}$; $\epsilon(320) \cong 8500 M^{-1} \text{ cm}^{-1}$.	75-1118
3.2	$2\text{SO}_3^- \rightarrow \text{SO}_3 + \text{SO}_3^{2-}$ $2\text{SO}_3^- \rightarrow \text{S}_2\text{O}_6^{2-}$	-	-	0.03	f.phot.	D.k. at 270-290 nm in N_2 or air-satd. SO_3^{2-} soln.; $2k/\epsilon = 1.3$ to 1.9×10^6 .	68-7072
		-	-	0.03	f.phot.	D.k. at 275 nm in air-free 0.1 M $\text{Na}_2\text{S}_2\text{O}_6$ soln.; $2k/\epsilon = (3.5 \pm 2.0) \times 10^6$.	68-7281
		$2k = 1.9 \times 10^9$ (rel.)	11.8	0.03	e-r.	Obs. steady-state SO_3^- concn. by esr in N_2O -satd. SO_3^{2-} soln.; rel. to $2k(\text{CH}_2\text{CO}_2^- + \cdot\text{CH}_2\text{CO}_2^-) = 1.0 \times 10^9$.	72-5049
		$2k = (1.1 \pm 0.2) \times 10^9$	3.7-9.8	$\rightarrow 0$	f.phot.	D.k. at 255-320 nm in $\text{SO}_3^{2-}/\text{HSO}_3^-$ soln.; $2k/\epsilon = 8.3 \times 10^5 - 3.6 \times 10^6$; same in $\text{S}_2\text{O}_6^{2-}$ soln.	72-7008
		$2k = 8.5 \times 10^8$	5.4, 10.2	$\rightarrow 0$	p.r.	D.k. in $\text{SO}_3^{2-}/\text{HSO}_3^-$ soln.	72-7008
		$2k = 5.4 \times 10^8$ (rel.)	-	-	phot.	C.k. in soln. contg. 1.6×10^{-3} M SO_3^{2-} , 0.68 M acetone and 10^{-3} M $\text{Na}_4\text{B}_2\text{O}_7$; rel. to $2k = 1.4 \times 10^9$ for second-order decay of $(\text{CH}_3)_2\text{COH}$.	73-5022
		$2k = (1.4 \pm 0.2) \times 10^9$ $2k = (0.85 \pm 0.2) \times 10^9$	10 5	-	p.r.	D.k. at 255 nm ($\epsilon = 1200 \pm 50 M^{-1} \text{ cm}^{-1}$) in $\text{SO}_2/\text{HSO}_3^-$ soln.; $k = 1.4 \times 10^9$ by pulse conductivity at pH 9.5; sulfate is formed by hydrolysis of SO_3 ; rel. amounts of sulfate and dithionate formed depend on pH.	74-1033
3.3	$\text{SO}_3^- + \text{N}_2\text{O} \rightarrow \text{SO}_3 + \text{O}^- + \text{N}_2$	$(3.5 \pm 1.0) \times 10^7$	13	1	p.r.	D.k. at 260 nm in N_2O -satd. soln. contg. 0.002 M SO_3^{2-} ; best fit with $k(\text{O}^- + \text{SO}_3^{2-}) = 2.5 \times 10^8$ and $\epsilon(260 \text{ nm})$ for $\text{SO}_3^- = 1300 M^{-1} \text{ cm}^{-1}$.	71-0461
3.4	$\text{SO}_3^- + \text{O}_2 \rightarrow \text{SO}_5^-$	$< 10^9$	-	-	f.phot.	Estd.	72-7008
3.5	$\text{SO}_3^- + \text{CH}_3\text{CH}_2\text{OH}$	$< 2 \times 10^3$	-	0.1	f.phot.	D.k. at 270 nm in N_2 -satd. 0.1 M $\text{S}_2\text{O}_6^{2-}$ soln.	72-7008
3.6	$\text{SO}_3^- + (\text{CH}_3)_2\text{CHOH}$	$< 10^3$	-	0.1	f.phot.	D.k. in N_2 -satd. 0.1 M $\text{S}_2\text{O}_6^{2-}$ soln.	72-7008

TABLE 3. Rates of reaction of SO_2^- , SO_3^- , S_2O_3^- , S_2O_2^- , SO_5^- , SeO_2^- and SeO_3^- in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	l	Method	Comment	Ref.
3.7	$\text{S}_2\text{O}_3^- + \text{S}_2\text{O}_3^{2-}$	$2k = (7.1 \pm 1.8) \times 10^9$	7	0.003	f.phot.	D.k. at 380 nm in N_2 -satd. $\text{S}_2\text{O}_3^{2-}$ soln.; $2k/\epsilon = 4.1 \times 10^6$; same at pH 10.9; addn. of ethanol, carbonate ion or allyl alcohol did not affect the decay rate.	68-7072
3.8	$\text{S}_2\text{O}_3^- + \text{S}_2\text{O}_3^{2-} \rightarrow \text{S}_4\text{O}_6^{3-}$	8×10^4	4.5, 9.6	<0.002	p.r.	P.b.k. at 370 nm or d.k. at 320 nm in $\text{S}_2\text{O}_3^{2-}$ soln.; $k = 1.8 \times 10^9$ at $l = 0.1$.	73-1027
3.9	$\text{S}_2\text{O}_2^- + \text{S}_2\text{O}_2^-$	$2k/\epsilon \approx 5 \times 10^4$	7, 10.9	0.003	f.phot.	D.k. at 270 nm in $\text{S}_2\text{O}_3^{2-}$ soln.; addn. of O_2 , methanol, ethanol or allyl alcohol did not affect the decay rate.	68-7072
3.10	$\text{SO}_5^- + \text{SO}_5^-$	$2k \approx 4 \times 10^8$	—	1	p.r.	D.k. in N_2O -satd. HSO_5^- soln.; $\epsilon(265 \text{ nm}) > 560 M^{-1} \text{ cm}^{-1}$; not simple order.	72-7008
3.11	$\text{SO}_5^- + \text{CH}_3\text{CH}_2\text{OH}$	$< 10^3$	9	—	f.phot.	D.k. in air-contg. $\text{S}_2\text{O}_6^{2-}$ soln.	72-7008
3.12	$2 \text{SeO}_2^- \rightarrow \text{Se}_2\text{O}_4^{2-}$	$2k/\epsilon = 8.8 \times 10^5$ $2k/\epsilon = 9.2 \times 10^5$	6.2 4.5–7.5	—	p.r.	D.k. in soln. of SeO_2 contg. 2-methyl-2-propanol; $\lambda_{\text{max}} = 330 \text{ nm}$; $\epsilon_{\text{max}} = 1450 M^{-1} \text{ cm}^{-1}$.	77-1173
3.13	$2 \text{SeO}_3^- \rightarrow \text{Se}_2\text{O}_6^{2-}$ or $\rightarrow \text{SeO}_3 + \text{SeO}_3^{2-}$	$2k/\epsilon = 4.0 \times 10^5$	9–11	—	p.r.	D.k. in soln. of SeO_2 contg. N_2O ; $\lambda_{\text{max}} = 430 \text{ nm}$; $\epsilon_{\text{max}} = 1350 M^{-1} \text{ cm}^{-1}$.	77-1173
3.14	$2 \text{HSeO}_3 \rightarrow \text{products}$	$2k/\epsilon = 7.5 \times 10^5$	5–6	—	p.r.	D.k. in soln. of SeO_2 contg. N_2O ; $\lambda_{\text{max}} = 430 \text{ nm}$; $\epsilon_{\text{max}} = 600 M^{-1} \text{ cm}^{-1}$.	77-1173
3.15	$2 \text{H}_2\text{SeO}_3^+ \rightarrow \text{products}$	$2k/\epsilon = 2.1 \times 10^6$	1	—	p.r.	D.k. in soln. of SeO_2 contg. HClO_4 ; $\lambda_{\text{max}} = 430 \text{ nm}$; $\epsilon = 930 M^{-1} \text{ cm}^{-1}$.	77-1173
3.16	$\text{SeO}_3^- + \text{tryptophan}$	$(3.4 \pm 0.4) \times 10^9$	7.4	0.006	p.r.	D.k. in N_2O -satd. SeO_3^{2-} soln.	76-1151

TABLE 4. Rates of reaction of NO_3 , NO_2 , N_2O_4 , NO_3^{2-} , N_2O_3 , NO_2^{2-} in aqueous solution

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
4.1	$\text{NO}_3 \rightarrow$ first order decay	$(7.59 \pm 0.21) \times 10^3 s^{-1}$	acid	—	p.r.	D.k. at 635 nm in 0.1–4 $M \text{HNO}_3$.	67-0002
		$9.5 \times 10^2 s^{-1}$	acid	—	f.phot.	D.k. in 0.1–6 $M \text{HNO}_3$ contg. $\text{K}_2\text{Ce}(\text{NO}_3)_6$.	67-7274
		$9.7 \times 10^3 s^{-1}$	2.8	—	p.r.	D.k. at 670 nm in 4 $M \text{NaNO}_3$; k varies with dose, concn. and pH, $k \sim 3 \times 10^4 s^{-1}$ at pH 7.	69-0417
		$8 \times 10^3 s^{-1}$	acid	—	p.r.	D.k. at 600 nm in 0.2–15 $M \text{HNO}_3$; 1st and 2nd order rates detd. by graphical method; dependent on dose and concn.	74-1140
		$(1.6 \pm 0.2) \times 10^5 s^{-1}$	nat	—	p.r.	D.k. at 635 nm in 5.0 and 7.35 $M \text{NaNO}_3$ soln.; discrepancy in k values attributed to dose rate effects.	76-1171
4.2	$\text{NO}_3 + \text{NO}_3 \rightarrow \text{N}_2\text{O}_6$	$2k = 8.4 \times 10^8$	acid	—	p.r.	D.k. at 670 nm in 1–4.0 $M \text{NaNO}_3$, $\geq 0.12 M \text{HNO}_3$ soln.; $2k/\epsilon = 2.8 \times 10^6$.	69-0417
		$2k = (0.79 \pm 0.04) \times 10^6$	acid	—	f.phot.	D.k. at 635 nm in 3.5–15 $M \text{HNO}_3$ and $10^{-3} M (\text{NH}_4)_2\text{Ce}(\text{NO}_3)_6$; $\epsilon(635 \text{ nm}) = 250 \pm 90 M^{-1} \text{cm}^{-1}$.	70-7728
		$2k = 3.7 \times 10^8$	acid	—	p.r.	D.k. at 600 nm in 0.2–15 $M \text{HNO}_3$; $\epsilon(600 \text{ nm}) = 285 M^{-1} \text{cm}^{-1}$; $2k/\epsilon = 1.3 \times 10^6 (\pm 20\%)$.	74-1140
4.3	$\text{NO}_3 + \text{Ce}^{\text{III}} \rightarrow$ $\text{Ce}^{\text{IV}} + \text{NO}_3^-$	$(1.70 \pm 0.04) \times 10^6$	acid	—	f.phot.	D.k. at 635 nm in 6 $M \text{HNO}_3$ contg. $10^{-3} M (\text{NH}_4)_2\text{Ce}(\text{NO}_3)_6$; $E_s = 1.0 \pm 0.2 \text{ kcal mol}^{-1}$ ($4.2 \pm 0.8 \text{ kJ mol}^{-1}$) based on 6 runs at 50°C ; k in 1–15 $M \text{HNO}_3$ varied from 0.64 to 4.1×10^6 indicating presence of several Ce^{III} complexes.	64-7009 70-7727
		$(3.70 \pm 0.1) \times 10^5$	nat	—	f.phot.	D.k. at 600 nm in 0.1 $M \text{K}_2\text{Ce}(\text{NO}_3)_6$ contg. $\text{Ce}(\text{NO}_3)_3$.	67-7274
		$(1.3 \pm 0.3) \times 10^6$	acid	—	p.r.	D.k. at 600 nm in 2 $M \text{HNO}_3$ contg. $10^{-2} M \text{Ce}^{\text{III}}$; cor. for $\text{NO}_3 + \text{NO}_3$.	74-1140
		1.0×10^8	—	—	p.r.	P.b.k. at 345 nm (Cl_2^-) in 2 $M \text{NO}_3^-$ soln.	76-1141
4.4	$\text{NO}_3 + \text{Cl}^-$	1.0×10^8	—	—	p.r.	D.k. at 600 nm in 2 $M \text{HNO}_3$ contg. $10^{-3} M \text{Fe}^{2+}$; cor. for $\text{NO}_3 + \text{NO}_3$.	74-1140
4.5	$\text{NO}_3 + \text{Fe}^{2+}$	$(8.0 \pm 1.6) \times 10^6$	acid	—	p.r.	D.k.	69-0417
4.6	$\text{NO}_3 + \text{NO}_2^-$	1.2×10^9	7	—	p.r.	D.k. at 600 nm.	67-7274
4.7	$\text{NO}_3 + \text{Ti}^+$	$(3.46 \pm 0.1) \times 10^7$	—	—	f.phot.	D.k. at 600 nm.	67-7274
4.8	Acetic acid $\text{NO}_3 + \text{CH}_3\text{COOH}$	$(4.6 \pm 0.4) \times 10^4$	—	—	f.phot.	D.k.	73-7569
4.9	2,6-Disulfoanthra- quinone radical ion (A^-) $\text{NO}_3 + \text{A}^- \rightarrow \text{NO}_3^- + \text{A}$	1.0×10^8	8.0	—	f.phot.	D.k.	67-7274
4.10	Ethanol $\text{NO}_3 + \text{CH}_3\text{CH}_2\text{OH}$	$(3.9 \pm 0.4) \times 10^6$	2	—	f.phot.	D.k. at 600 nm in 5 $M \text{HNO}_3$; cor. for $\text{NO}_3 + \text{NO}_3$.	74-1140
		$(2.2 \pm 0.4) \times 10^6$	acid	—	p.r.		

TABLE 4. Rates of reaction of NO_3^- , NO_2 , N_2O_4 , NO_3^{2-} , N_2O_3 , NO_2^{2-} in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
4.11	Ethylene glycol $\text{NO}_3^- + \text{CH}_2\text{OHCH}_2\text{OH}$	$(1.6 \pm 0.3) \times 10^6$	acid	—	p.r.	D.k. at 600 nm in 5 M HNO_3 ; cor. for $\text{NO}_3^- + \text{NO}_3^-$.	74-1140
4.12	Formic acid $\text{NO}_3^- + \text{HCOOH}$	$(2.06 \pm 0.1) \times 10^5$	—	—	f.phot.	D.k. at 600 nm.	67-7274
4.13	Glycerol $\text{NO}_3^- + \text{CH}_2\text{OHCHOHCH}_2\text{OH}$	$(1.8 \pm 0.4) \times 10^6$	acid	—	p.r.	D.k. at 600 nm in 5 M HNO_3 ; cor. for $\text{NO}_3^- + \text{NO}_3^-$.	74-1140
4.14	Methanol $\text{NO}_3^- + \text{CH}_3\text{OH}$	$(1.0 \pm 0.1) \times 10^6$ $(1.2 \pm 0.2) \times 10^6$	— acid	—	f.phot. p.r.	D.k. at 600 nm. D.k. at 600 nm in 5 M HNO_3 ; cor. for $\text{NO}_3^- + \text{NO}_3^-$.	67-7274 74-1140
4.15	2-Propanol $\text{NO}_3^- + (\text{CH}_3)_2\text{CHOH}$	$(3.6 \pm 0.2) \times 10^6$ $(2.3 \pm 0.5) \times 10^6$	— acid	—	f.phot. p.r.	D.k. at 600 nm. D.k. at 600 nm in 5 M HNO_3 ; cor. for $\text{NO}_3^- + \text{NO}_3^-$.	67-7274 74-1140
4.16	$\text{NO}_2 + \text{NO}_2 \rightarrow \text{N}_2\text{O}_4$	$2k = (7.6 \pm 2.7) \times 10^7$	—	—	f.phot.	C.k. with ferrocyanide ion in NO_3^- soln.; obs. effect of intensity on buildup of ferricyanide at 420 nm; assumed $\epsilon_{\text{Fe}}/\epsilon_{\text{NO}_2} \cong 10$ and $k(\text{NO}_2 + \text{Fe}(\text{CN})_6^{4-}) = 4.3 \times 10^6$.	68-7024
		$2k = 3 \times 10^7$	7	—	p.r.	D.k. at 400 nm in NaNO_3 ($>0.5 M$) soln.; $2k/\epsilon = 1.44 \times 10^5$; assumed $\epsilon_{400} \cong 208 M^{-1} \text{cm}^{-1}$.	69-0417
		$2k = (9.0 \pm 2.0) \times 10^8$	—	—	p.r.	D.k. at 400 nm in NO_2^- soln. saturated with N_2O ; $K = 1.53 \times 10^{-5} M$ for $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2$.	69-0439
		$2k = (9.2 \pm 3.0) \times 10^8$	—	—	p.r.	D.k. at 410 nm in NO_2^- soln. saturated with N_2O .	76-1171
4.17	$\text{NO}_2 + \text{Fe}(\text{CN})_6^{4-} \rightarrow \text{NO}_2^- + \text{Fe}(\text{CN})_6^{3-}$	$(4.3 \pm 1.0) \times 10^6$	7	—	f.phot.	P.b.k. at 420 nm in NO_3^- soln.	68-7024
4.18	$\text{NO}_2 + \text{NO} \rightarrow \text{N}_2\text{O}_3$	1.1×10^9	7	—	p.r.	D.k. at 400 nm in $\text{NO}-\text{N}_2\text{O}$ soln.; $k_r = 8.0 \times 10^4 s^{-1}$.	70-0228
4.19	$\text{N}_2\text{O}_4 (+ \text{H}_2\text{O}) \rightarrow \text{NO}_2^- + \text{NO}_3^- + 2\text{H}^+$	$(1.0 \pm 0.1) \times 10^3 s^{-1}$ $(6.0 \pm 1.0) \times 10^4 s^{-1}$	7 —	—	p.r. p.r.	— D.k. at 350 nm.	69-0439 76-1171
4.20	$\text{NO}_3^{2-} + \text{H}_2\text{O} \rightarrow \text{NO}_2 + 2\text{OH}^-$	1.0×10^3	alk.	—	p.r.	D.k. at 290 nm; also condy. change; 400 nm absorption obs. (NO_2); $pK = 7.8, 4.8$ for $(\text{H}^+ +) \text{NO}_3^{2-} \rightleftharpoons \text{HNO}_3^- (+ \text{H}^+) \rightleftharpoons \text{H}_2\text{NO}_3$.	70-0151
		1.3×10^3	—	—	p.r.	Conductivity change in NO_3^- soln. contg. 0.2 M MeOH.	70-0254
		1.6×10^3	—	—	p.r.	D.k. at 260 nm.	76-1171
4.21	$\text{HNO}_3^- \rightarrow \text{NO}_2 + \text{OH}^-$	—	—	—	p.r.	$t_{0.5} = 3\mu s$.	70-0151
4.22	$\text{NO}_3^{2-} + \text{O}_2 \rightarrow \text{NO}_3^- + \text{O}_2^-$	2×10^8 (rel.)	12	—	γ -r.	Evaluated $k/k(\text{NO}_3^{2-} + \text{H}_2\text{O}) = 2 \times 10^5$ from dependence of $G(\text{NO}_2^-)$ on $[\text{O}_2]$; assume $k(\text{NO}_3^{2-} + \text{H}_2\text{O}) = 1 \times 10^3$.	67-0032
4.23	$\text{NO}_3^{2-} + \text{H}_2\text{PO}_4^- \rightarrow \text{HNO}_3^- + \text{HPO}_4^{2-}$	5×10^8	7	—	p.r.	D.k. at 290 nm.	70-0151

TABLE 4. Rates of reaction of NO_3 , NO_2 , N_2O_4 , NO_3^{2-} , N_2O_3 , NO_2^{2-} in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
4.24	$\text{N}_2\text{O}_3 (+ \text{H}_2\text{O}) \rightarrow 2\text{NO}_2^- + 2\text{H}^+$	$5.3 \times 10^2 s^{-1}$	5	—	p.r.	D.k. at 260 nm as well as condy. change in NO_2^- -NO- N_2O soln.; $K(\text{N}_2\text{O}_3 \rightleftharpoons \text{NO} + \text{NO}_2) = 7.3 \times 10^{-5} M$. (See also 70-7264).	70-0228
4.25	$\text{NO}_2^{2-} + \text{H}_2\text{O} \rightarrow \text{NO} + 2\text{OH}^-$	1.0×10^3	9.0	—	p.r.	D.k. at 270 nm; soln. contains 0.25 <i>M tert</i> -BuOH; $pK = 7.7, 5.7$ for $(\text{H}^+ +) \text{NO}_2^{2-} \rightleftharpoons \text{HNO}_2^- (+ \text{H}^+) \rightleftharpoons \text{H}_2\text{NO}_2$. (See also 70-7264).	69-0439
		$(7.7 \pm 1.7) \times 10^2$	—	—	p.r.	D.k. at 270 nm.	76-1181

TABLE 5. Rates of reaction of PO_4^{2-} in aqueous solution

No.	Reaction	$k(\text{M}^{-1} \text{s}^{-1})$	pH	I	Method	Comment	Ref.
5.1	$\text{H}_2\text{PO}_3 + \text{H}_2\text{PO}_4 \rightarrow \text{H}_4\text{P}_2\text{O}_8$	$2k \cong 1 \times 10^9$	3.5, 4.1	~ 1	p.r.	D.k. at 500 nm in 1 <i>M</i> H_2PO_4^- contg. N_2O ; $2k/\epsilon = 2.5 \times 10^6$; $\epsilon \cong 400 \text{ M}^{-1} \text{cm}^{-1}$.	70-0302
		$2k = (2.5 \pm 0.5) \times 10^8$	3.8	-	p.r.	D.k. at 500 nm in 1-3 <i>M</i> phosphate satd. with N_2O ; interpreted from $2k/\epsilon = (11 \pm 1) \times 10^3 \text{ cm s}^{-1}$ and $\epsilon = 230 \text{ M}^{-1} \text{cm}^{-1}$.	73-1050
5.2	$\text{HPO}_3^- + \text{HPO}_4^- \rightarrow \text{H}_2\text{P}_2\text{O}_8^{2-}$	$2k = 4.7 \times 10^8$	6.6- 8.2	~ 3	p.r.	D.k. at 500 nm in 1 <i>M</i> HPO_4^{2-} contg. N_2O ; $2k/\epsilon = 5.5 \times 10^5 - 6.1 \times 10^5$; $\epsilon \cong 800 \text{ M}^{-1} \text{cm}^{-1}$.	70-0302
		$2k \cong 1.4 \times 10^8$	~ 9	$\rightarrow 0$	p.r.	D.k. at 500 nm in N_2O -satd. 0.3 <i>M</i> phosphate; interpreted from $2k/\epsilon \cong 3-7 \times 10^3 \text{ cm s}^{-1}$ and $\epsilon = 720 \text{ M}^{-1} \text{cm}^{-1}$.	73-1050
5.3	$\text{PO}_4^{2-} + \text{PO}_4^{2-} \rightarrow \text{P}_2\text{O}_8^{4-}$	$2k \cong 1 \times 10^9$	11.2- 11.7	~ 6	p.r.	D.k. at 500 nm in 1 <i>M</i> PO_4^{3-} contg. N_2O ; $2k/\epsilon = 1.9 \times 10^5 - 2.4 \times 10^5$; $\epsilon = 4800 \text{ M}^{-1} \text{cm}^{-1}$.	70-0302
		$2k = (1.0 \pm 0.2) \times 10^8$	~ 12	$\rightarrow 0$	p.r.	D.k. at 500 nm in N_2O -satd. 0.3 <i>M</i> phosphate; interpreted from $2k/\epsilon \cong 10^3 \text{ cm s}^{-1}$ and $\epsilon = 2700 \text{ M}^{-1} \text{cm}^{-1}$.	73-1050
5.4	$\text{HPO}_4^- + \text{Br}^- \rightarrow \text{HPO}_3^{2-} + \text{Br}$	6.5×10^6	9	~ 0.2	p.r.	D.k. at 520 nm in 0.02 <i>M</i> $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.5	$\text{H}_2\text{PO}_4 + \text{Br}^-$	8×10^8	4	~ 0.06	p.r.	D.k. at 520 nm in 0.02 <i>M</i> $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.6	$\text{HPO}_4^- + \text{Cl}^-$	$\leq 10^4$	7	~ 0.12	p.r.	D.k. at 520 nm in 0.02 <i>M</i> $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.7	$\text{H}_2\text{PO}_3 + \text{Cl}^- \rightarrow \text{H}_2\text{PO}_4^- + \text{Cl}$	2.2×10^6	4	~ 0.06	p.r.	D.k. at 520 nm in 0.02 <i>M</i> $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.8	$\text{PO}_4^{2-} + \text{I}^- \rightarrow \text{PO}_4^{3-} + \text{I}$	3×10^8	12	~ 0.2	p.r.	D.k. at 520 nm in 0.02 <i>M</i> $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.9	$\text{HPO}_4^- + \text{N}_2\text{H}_4$	4.9×10^8	9.4	~ 0.2	p.r.	D.k. at 520 nm in 0.02 <i>M</i> $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.10	$\text{HPO}_4^- + \text{N}_2\text{H}_5^+$	1.4×10^8	7	~ 0.12	p.r.	D.k. at 520 nm in 0.02 <i>M</i> $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.11	$\text{H}_2\text{PO}_3 + \text{N}_2\text{H}_5^+$	1.9×10^8	4	~ 0.06	p.r.	D.k. at 520 nm in 0.02 <i>M</i> $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.12	$\text{HPO}_4^- + \text{N}_3^-$	1.1×10^8	7	~ 0.12	p.r.	D.k. at 520 nm in 0.02 <i>M</i> $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.13	$\text{HPO}_4^- + \text{NH}_2\text{OH}$	4.9×10^8	9	~ 0.2	p.r.	D.k. at 520 nm in 0.02 <i>M</i> $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.14	$\text{H}_2\text{PO}_3 + \text{NH}_3\text{OH}^+$	1.2×10^7	4	~ 0.06	p.r.	D.k. at 520 nm in 0.02 <i>M</i> $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.15	$\text{HPO}_4^- + \text{NO}_2^- \rightarrow \text{HPO}_3^{2-} + \text{NO}_2$	1.4×10^7	7	~ 0.12	p.r.	D.k. at 520 nm in 0.02 <i>M</i> $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.16	$\text{PO}_4^{2-} + \text{OH}^- \rightarrow \text{PO}_4^{3-} + \text{OH}$	$(2.0 \pm 0.2) \times 10^5$	> 12	~ 1	p.r.	D.k. at 500 nm in 0.3 <i>M</i> HPO_4^{2-} soln.; first order decay rate 9.15×10^3 at pH 12.65.	73-1050
		5×10^5	> 12	~ 0.2	p.r.	D.k. at 520 nm in 0.02 <i>M</i> $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.17	$\text{HPO}_4^- + \text{H}_2\text{O}_2 \rightarrow \text{HPO}_3^{2-} + 2\text{H}^+ + \text{O}_2^-$	$(2.7 \pm 1.0) \times 10^7$	9.0	0.1	f.phot.	D.k. at 500 nm in 0.03 <i>M</i> Na_2HPO_4 soln.	70-0326
5.18	$\text{H}_2\text{PO}_3 + \text{H}_2\text{O}_2$	$(5.5 \pm 1.0) \times 10^7$	4.5	0.1	f.phot.	D.k. at 500 nm in 0.1 <i>M</i> NaH_2PO_4 soln.	70-0326

TABLE 5. Rates of reaction of PO_4^{2-} in aqueous solution — Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
5.19	$\text{PO}_4^{2-} + \text{HPO}_3^{2-} \rightarrow \text{PO}_4^{3-} + \text{H}^+ + \text{PO}_3^{2-}$	5.5×10^5	12	~ 0.2	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.20	$\text{HPO}_4^- + \text{HPO}_3^{2-}$	5.9×10^9	9	~ 0.2	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.21	$\text{H}_2\text{PO}_4 + \text{H}_2\text{PO}_3^-$	4×10^7	4	~ 0.06	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.22	$\text{PO}_4^{2-} + \text{H}_2\text{PO}_2^- \rightarrow \text{PO}_4^{3-} + \text{H}^+ + \text{HPO}_2^-$	7.9×10^7	12	~ 0.2	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.23	$\text{HPO}_4^- + \text{H}_2\text{PO}_2^-$	5.9×10^7	7	~ 0.12	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.24	$\text{H}_2\text{PO}_4 + \text{H}_2\text{PO}_2^-$	3.9×10^8	4	~ 0.06	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.25	$\text{HPO}_4^- + \text{HPO}_3^{2-} \rightarrow \text{H}_2\text{PO}_4^- + \text{PO}_5^{2-}$	$< 10^5$	-	-	p.r.	D.k. at 520 nm.	77-1047
5.26	$\text{PO}_4^{2-} + \text{SO}_3^{2-} \rightarrow \text{PO}_4^{3-} + \text{SO}_3^-$	4.1×10^7	12	~ 0.2	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.27	$\text{HPO}_4^- + \text{SO}_3^{2-}$	2.7×10^7	9	~ 0.2	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.28	$\text{H}_2\text{PO}_4 + \text{HSO}_3^-$	2.7×10^8	4	~ 0.06	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.29	$\text{HPO}_4^- + \text{S}_2\text{O}_3^{2-}$	1.0×10^8	7	~ 0.12	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.30	$\text{HPO}_4^- + \text{S}_2\text{O}_5^{2-}$	1.9×10^8	7	~ 0.12	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	78-1075
5.31	Acetate ion $\text{HPO}_4^- + \text{CH}_3\text{COO}^- \rightarrow \text{H}_2\text{PO}_4^- + \text{CH}_2\text{COO}^-$	8.5×10^4	7.1	~ 0.12	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
5.32	Acetic acid $\text{H}_2\text{PO}_4 + \text{CH}_3\text{COOH} \rightarrow \text{H}_3\text{PO}_4 + \text{CH}_2\text{COOH} (1)$ $\text{H}_2\text{PO}_4 + \text{CH}_3\text{COOH} \rightarrow \text{H}_3\text{PO}_4 + \text{CO}_2 + \text{CH}_3 (2)$	3.4×10^5	3.6	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.; $k_1/k_2 \cong 0.5$ detd. from CO_2 yields by γ -r. (78-1505).	77-1106
5.33	Acetone $\text{H}_2\text{PO}_4 + \text{CH}_3\text{COCH}_3 \rightarrow \text{H}_3\text{PO}_4 + \text{CH}_2\text{COCH}_3$	3.3×10^5	3.2-4.6	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
5.34	Acrylamide	$\leq 10^6$	7.4	~ 0.12	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1503
		2.2×10^8	4	~ 0.06	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1503
5.35	Acrylic acid $\text{HPO}_4^- + \text{CH}_2=\text{CHCOO}^- \rightarrow ^-\text{HO}_3\text{POCH}_2\text{CHOO}^-$ $\text{H}_2\text{PO}_4 + \text{CH}_2=\text{CHCOOH} \rightarrow \text{H}_2\text{O}_3\text{POCH}_2\text{CHCOOH}$	6.2×10^6	7.5	~ 0.12	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.; product radical obs. by esr (77-5209).	77-1503
		1.6×10^8	4	~ 0.06	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1503
5.36	Acrylonitrile	4.4×10^7	7.3	~ 0.12	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1503
		6×10^7	3.8	~ 0.06	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1503
5.37	Alanine $\text{PO}_4^{2-} + \text{CH}_3\text{CH}(\text{NH}_2)\text{COO}^-$	1.6×10^7	12	~ 0.2	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
5.38	Allyl alcohol	2.1×10^8	7	~ 0.12	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1503
		1.4×10^9	4	~ 0.06	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1503
5.39	Allyl cyanide	3.4×10^7	7.2	~ 0.12	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1503
		8.8×10^8	4	~ 0.06	p.r.	D.k. at 520 nm in 0.02 M $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1503

TABLE 5. Rates of reaction of PO_4^{2-} in aqueous solution — Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
5.40	<i>p</i> -Aminobenzoic acid $\text{H}_2\text{PO}_4 + \text{NH}_2\text{C}_6\text{H}_4\text{COOH}$	1.5×10^9	3.3	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
5.41	Benzoate ion $\text{HPO}_4^{2-} + \text{C}_6\text{H}_5\text{COO}^-$	$< 10^7$	7.1	~ 0.12	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
5.42	Benzoic acid $\text{H}_2\text{PO}_4 + \text{C}_6\text{H}_5\text{COOH} \rightarrow$ $(\text{C}_6\text{H}_5\text{COOH})^+ + \text{H}_2\text{PO}_4^-$ $(\text{C}_6\text{H}_5\text{COOH})^+ \rightarrow \text{C}_6\text{H}_5 + \text{CO}_2 + \text{H}^+ (1)$ $(\text{C}_6\text{H}_5\text{COOH})^+ + \text{H}_2\text{O} \rightarrow \text{HOC}_6\text{H}_5\text{COOH} (2)$	2.4×10^8	3.2	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.; $k_1/k_2 \approx 1$ detd. from CO_2 yield by γ -r. (78-1505).	77-1106
5.43	<i>p</i> -Chlorobenzoate ion $\text{H}_2\text{PO}_4 + \text{ClC}_6\text{H}_4\text{COO}^-$	4.8×10^7	5.0	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
5.44	<i>p</i> -Chlorobenzoic acid $\text{H}_2\text{PO}_4 + \text{ClC}_6\text{H}_4\text{COOH}$	1.0×10^8	3.3	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
5.45	Crotonate ion $\text{HPO}_4^{2-} + \text{CH}_3\text{CH}=\text{CHCOO}^-$	3.5×10^6	7.4	~ 0.12	p.r.	—	77-1503
5.46	Crotonic acid $\text{H}_2\text{PO}_4 + \text{CH}_3\text{CH}=\text{CHCOOH}$	4.5×10^8	4	~ 0.06	p.r.	—	77-1503
5.47	<i>p</i> -Cyanobenzoate ion $\text{H}_2\text{PO}_4 + \text{CNC}_6\text{H}_4\text{COO}^-$	1.0×10^7	4.6	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
5.47a	<i>p</i> -Cyanophenoxide ion	1.9×10^8	11.2	~ 0.2	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1503
5.48	Deoxyribose $\text{HPO}_4^{2-} + \text{C}_5\text{H}_{10}\text{O}_5 \rightarrow$ $\text{H}_2\text{PO}_4^- + \text{C}_5\text{H}_8\text{O}_5$	$(7.5 \pm 1.2) \times 10^7$	9.0	—	f.phot.	D.k. at 500 nm in 0.03 <i>M</i> HPO_4^{2-} .	70-0326
5.49	Dihydrouracil	$< (2.9 \pm 0.6) \times 10^7$	9.0	—	f.phot.	D.k. at 500 nm in 0.03 <i>M</i> HPO_4^{2-} .	70-0326
5.50	Ethanol $\text{PO}_4^{2-} + \text{CH}_3\text{CH}_2\text{OH} \rightarrow$ $\text{HPO}_4^{2-} + \text{CH}_3\text{CHOH}$ $\text{HPO}_4^{2-} + \text{CH}_3\text{CH}_2\text{OH} \rightarrow$ $\text{H}_2\text{PO}_4^- + \text{CH}_3\text{CHOH}$	1.9×10^7 $(4.0 \pm 0.6) \times 10^7$	12.0 9.0	~ 0.2 —	p.r. f.phot.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln. D.k. at 500 nm in 0.03 <i>M</i> HPO_4^{2-} .	77-1106 70-0326
		2.0×10^7	7	~ 0.12	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
	$\text{H}_2\text{PO}_4 + \text{CH}_3\text{CH}_2\text{OH} \rightarrow$ $\text{H}_3\text{PO}_4 + \text{CH}_3\text{CHOH}$	7.7×10^7	4	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
5.51	Formate ion $\text{PO}_4^{2-} + \text{HCOO}^- \rightarrow$ $\text{HPO}_4^{2-} + \text{COO}^-$ $\text{HPO}_4^{2-} + \text{HCOO}^- \rightarrow$ $\text{H}_2\text{PO}_4^- + \text{COO}^-$	2.2×10^7 $(2.9 \pm 0.7) \times 10^7$	12 9.0	~ 0.2 —	p.r. f.phot.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln. D.k. at 500 nm in 0.03 <i>M</i> HPO_4^{2-} soln.	77-1106 70-0326
		2.5×10^7	7	~ 0.12	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
	$\text{H}_2\text{PO}_4 + \text{HCOO}^- \rightarrow$ $\text{H}_3\text{PO}_4 + \text{COO}^-$	1.5×10^8	4.5	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
5.52	Fumaric acid $\text{H}_2\text{PO}_4 + \text{HOOCCH}=\text{CHCOOH} \rightarrow$ $\text{HOOCCHCH}(\text{COOH})\text{OPO}_3\text{H}_2$	1.5×10^7	3.2– 4.6	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.; product radical obs. by esr (77-5209).	77-1106
5.53	D-Glucose $\text{HPO}_4^{2-} + \text{C}_6\text{H}_{12}\text{O}_6 \rightarrow$ $\text{H}_2\text{PO}_4^- + \text{C}_6\text{H}_{11}\text{O}_6$ $\text{H}_2\text{PO}_4 + \text{C}_6\text{H}_{12}\text{O}_6 \rightarrow$ $\text{H}_3\text{PO}_4 + \text{C}_6\text{H}_{11}\text{O}_6$	$(8.0 \pm 0.2) \times 10^7$ $(1.1 \pm 0.3) \times 10^8$	9.0 4.5	— —	f.phot. f.phot.	D.k. at 500 nm in 0.03 <i>M</i> HPO_4^{2-} soln. D.k. at 500 nm in 0.1 <i>M</i> H_2PO_4^- soln.	70-0326 70-0326
5.54	Glycine $\text{PO}_4^{2-} + \text{NH}_2\text{CH}_2\text{COO}^- \rightarrow$ $\text{HPO}_4^{2-} + \text{NH}_2\text{CHCOO}^-$ $\text{H}_2\text{PO}_4 + \text{NH}_3^+\text{CH}_2\text{COO}^- \rightarrow$ $\text{H}_3\text{PO}_4 + \text{NH}_3^+\text{CHCOO}^-$	2.6×10^7 $< 10^5$	12 4.5	~ 0.2 ~ 0.06	p.r. p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln. D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106 77-1106

TABLE 5. Rates of reaction of PO_4^{2-} in aqueous solution — Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
5.55	<i>p</i> -Hydroxybenzoate ion $\text{HPO}_4^{2-} + \text{OHC}_6\text{H}_4\text{COO}^- \rightarrow$ $\text{H}_2\text{PO}_4^- + \text{OC}_6\text{H}_4\text{COO}^-$	1.7×10^8	7.2	~ 0.12	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
	$\text{PO}_4^{2-} + \text{OHC}_6\text{H}_4\text{COO}^- \rightarrow$ $\text{HPO}_4^{2-} + \text{OC}_6\text{H}_4\text{COO}^-$	5×10^7	11.4	~ 0.2	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1503
5.56	<i>p</i> -Hydroxybenzoic acid $\text{H}_2\text{PO}_4^- + \text{OHC}_6\text{H}_4\text{COOH} \rightarrow$	1.3×10^9	3.3	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106
5.57	Maleic acid $\text{H}_2\text{PO}_4^- + \text{HOOCCH}=\text{CHCOOH} \rightarrow$ $\text{HOOCCHCH}(\text{COOH})\text{OPO}_3\text{H}_2$	3.1×10^7	3.2– 4.6	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln. Product radical obs. by esr (77-5209).	77-1106
5.58	Malonic acid $\text{H}_2\text{PO}_4^- + \text{CH}_2(\text{COOH})_2 \rightarrow$ $\text{H}_3\text{PO}_4 + \text{CH}(\text{COOH})_2$ (1) $\text{H}_2\text{PO}_4^- + \text{CH}_2(\text{COOH})_2 \rightarrow$ $\text{H}_3\text{PO}_4 + \text{CH}_2\text{COOH} + \text{CO}_2$ (2)	1.8×10^5	3.2– 4.6	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln. Product radical obs. by esr (77-5209); $k_1/k_2 \cong 1$ detd. from CO_2 yield by γ -r. (78-1505).	77-1106
5.59	Methacrylate ion $\text{HPO}_4^{2-} + \text{CH}_2=\text{C}(\text{CH}_3)\text{COO}^-$	2.3×10^7	7.3	~ 0.12	p.r.	—	77-1503
5.60	Methacrylic acid $\text{H}_2\text{PO}_4^- + \text{CH}_2=\text{C}(\text{CH}_3)\text{COOH}$	7.3×10^8	4	~ 0.06	p.r.	—	77-1503
5.61	Methacrylonitrile $\text{HPO}_4^{2-} + \text{CH}_2=\text{C}(\text{CH}_3)\text{CN}$ $\text{H}_2\text{PO}_4^- + \text{CH}_2=\text{C}(\text{CH}_3)\text{CN}$	2.8×10^7 3.9×10^8	7.4 3.9	~ 0.12 ~ 0.06	p.r. p.r.	— —	77-1503 77-1503
5.62	Methanol $\text{PO}_4^{2-} + \text{CH}_3\text{OH} \rightarrow$ $\text{HPO}_4^{2-} + \text{CH}_2\text{OH}$ $\text{HPO}_4^{2-} + \text{CH}_3\text{OH} \rightarrow$ $\text{H}_2\text{PO}_4^- + \text{CH}_2\text{OH}$ $\text{H}_2\text{PO}_4^- + \text{CH}_3\text{OH} \rightarrow$ $\text{H}_3\text{PO}_4 + \text{CH}_2\text{OH}$	1.0×10^7 1.0×10^7 4.1×10^7	12 7 4	~ 0.2 ~ 0.12 ~ 0.06	p.r. p.r. p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln. D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln. D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106 77-1106 77-1106
5.62a	<i>p</i> -Methoxyphenoxide ion $\text{H}_2\text{PO}_4^- + \text{CH}_2\text{C}(\text{CH}_3)_2\text{OH}$	8.2×10^8	11.3	~ 0.2	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1503
5.63	Methyl methacrylate $\text{HPO}_4^{2-} + \text{CH}_2=\text{C}(\text{CH}_3)\text{COOCH}_3$ $\text{H}_2\text{PO}_4^- + \text{CH}_2=\text{C}(\text{CH}_3)\text{COOCH}_3$	3.9×10^7 6.2×10^8	7.3 4	~ 0.12 ~ 0.06	p.r. p.r.	— —	77-1503 77-1503
5.64	2-Methyl-2-propanol $\text{PO}_4^{2-} + (\text{CH}_3)_3\text{COH} \rightarrow$ $\text{HPO}_4^{2-} + \text{CH}_2\text{C}(\text{CH}_3)_2\text{OH}$ $\text{HPO}_4^{2-} + (\text{CH}_3)_3\text{COH} \rightarrow$ $\text{H}_2\text{PO}_4^- + \text{CH}_2\text{C}(\text{CH}_3)_2\text{OH}$ $\text{H}_2\text{PO}_4^- + (\text{CH}_3)_3\text{COH} \rightarrow$ $\text{H}_3\text{PO}_4 + \text{CH}_2\text{C}(\text{CH}_3)_2\text{OH}$	4.2×10^5 4.5×10^5 3.9×10^6	12 7 4	~ 0.2 ~ 0.12 ~ 0.06	p.r. p.r. p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln. D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln. D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106 77-1106 77-1106
5.64a	Phenoxide ion $\text{H}_2\text{PO}_4^- + (\text{CH}_3)_2\text{CHOH} \rightarrow$ $\text{HPO}_4^{2-} + (\text{CH}_3)_2\text{COH}$ $\text{HPO}_4^{2-} + (\text{CH}_3)_2\text{CHOH} \rightarrow$ $\text{H}_2\text{PO}_4^- + (\text{CH}_3)_2\text{COH}$	5.9×10^8 $(4.0 \pm 1.0) \times 10^7$ 2.5×10^7	11.6 9.0 7	~ 0.2 — ~ 0.12	p.r. f.phot. p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln. D.k. at 500 nm in 0.03 <i>M</i> HPO_4^{2-} soln. D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1503 70-0326 77-1106
5.65	2-Propanol $\text{H}_2\text{PO}_4^- + (\text{CH}_3)_2\text{CHOH} \rightarrow$ $\text{H}_3\text{PO}_4 + (\text{CH}_3)_2\text{COH}$	1.8×10^7 $(1.6 \pm 0.3) \times 10^8$ 1.4×10^8	12 4.5 4	~ 0.2 — ~ 0.06	p.r. f.phot. p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln. D.k. at 500 nm in 0.1 <i>M</i> H_2PO_4^- soln. D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77-1106 70-0326 77-1106

TABLE 5. Rates of reaction of PO_4^{2-} in aqueous solution — Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
5.66	Propionic acid $\text{H}_2\text{PO}_4 + \text{CH}_3\text{CH}_2\text{COOH} \rightarrow$ $\text{H}_3\text{PO}_4 + \text{CH}_3\text{CHCOOH}$ (1) $\text{H}_2\text{PO}_4 + \text{CH}_3\text{CH}_2\text{COOH} \rightarrow$ $\text{H}_3\text{PO}_4 + \text{CH}_3\text{CH}_2 + \text{CO}_2$ (2)	4.2×10^6	3.2– 4.6	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.; $k_1/k_2 \cong 4$ detd. from CO_2 yield by γ -r. (78–1505).	77–1106
5.67	Ribose $\text{HPO}_4^{2-} + \text{C}_5\text{H}_{10}\text{O}_5 \rightarrow$ $\text{H}_2\text{PO}_4^- + \text{C}_5\text{H}_9\text{O}_5$	$(9.0 \pm 2.0) \times 10^7$	9.0	–	f.phot.	D.k. at 500 nm in 0.03 M HPO_4^{2-} soln.	70–0326
5.68	Succinic acid $\text{H}_2\text{PO}_4 + (\text{CH}_2\text{COOH})_2 \rightarrow$ $\text{H}_3\text{PO}_4 + \text{HOOCCH}_2\text{CHCOOH}$	1.6×10^6	3.2– 4.6	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77–1106
5.69	Terephthalate ion $\text{H}_2\text{PO}_4 + \text{HOOC}_6\text{H}_4\text{COO}^-$	3.5×10^7	5.2	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77–1106
5.70	Terephthalic acid $\text{H}_2\text{PO}_4 + \text{HOOC}_6\text{H}_4\text{COOH}$	$\sim 6 \times 10^7$	3.4	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77–1106
5.71	Thymine $(9.6 \pm 1.0) \times 10^7$	$(9.6 \pm 1.0) \times 10^7$	9.0	~ 0.2	f.phot.	D.k. at 500 nm in 0.03 M HPO_4^{2-} soln.	70–0326
5.72	Toluic acid $\text{H}_2\text{PO}_4 + \text{CH}_3\text{C}_6\text{H}_4\text{COOH}$	5.4×10^8	3.3	~ 0.06	p.r.	D.k. at 520 nm in $\text{Li}_4\text{P}_2\text{O}_8$ soln.	77–1106
5.73	Uracil $(9.7 \pm 2.0) \times 10^7$	$(9.7 \pm 2.0) \times 10^7$	9.0	–	f.phot.	D.k. at 500 nm in 0.03 M HPO_4^{2-} soln.	70–0326
	$(6.0 \pm 2.0) \times 10^8$	$(6.0 \pm 2.0) \times 10^8$	4.5	–	f.phot.	D.k. at 500 nm in 0.1 M H_2PO_4^- soln.	70–0326

TABLE 6. Rates of reaction of Cl_2^- in aqueous solution

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
6.1	$\text{Cl}_2^- + \text{Cl}_2^- \rightarrow \text{Cl}_3^- + \text{Cl}^-$	$2k = (1.4 \pm 0.2) \times 10^{10}$	1.1, 6	0.5	f.phot.	D.k. in NaCl-HClO_4 soln.; assumed $\epsilon(350 \text{ nm}) = 1.25 \times 10^4 M^{-1} \text{ cm}^{-1}$.	67-7171
		$2k = (1.4 \pm 0.3) \times 10^{10}$	3.1	-	p.r.	D.k. in $\text{N}_2\text{O-satd. Cl}^-$ soln. assuming $\epsilon(360 \text{ nm}) = 10^4 M^{-1} \text{ cm}^{-1}$; independent of pH 0.9 - 3.2.	68-0313
		$2k = (1.2 \pm 0.3) \times 10^{10}$ $2k = 5.2 \times 10^9$	1.9 1	0.1 0.2	p.r. f.phot.	D.k.	72-3107
						D.k. in FeCl^{2+} soln. ($10^{-3} M \text{ Fe}^{3+}$, $0.1 M \text{ H}^+$, $0.1 M \text{ Cl}^-$); $\epsilon(366 \text{ nm}) = 1.0 \times 10^4 M^{-1} \text{ cm}^{-1}$.	73-7159
		$2k = (1.8 \pm 0.5) \times 10^{10}$	-	~ 1	p.r.	D.k. in NaCl or HCl soln. assuming $\epsilon = 1.25 \times 10^4 M^{-1} \text{ cm}^{-1}$.	74-1087
		$2k = (4 \text{ to } 14) \times 10^9$	-	< 10	p.r.	D.k. in LiCl soln.; values from graph; k increased as concn. decreased.	74-1149
		$2k = (2.7 \pm 0.5) \times 10^9$	7.0	< 14	p.r.	D.k. in $1.5\text{--}14 M \text{ LiCl}$ soln. at 340 nm ; calcd. from obs. $2k/\epsilon = (5 \text{ to } 8) \times 10^5$.	75-1154
		$2k = (1.3 \pm 0.5) \times 10^{10}$	-	-	p.r.	D.k. in Cl^- soln.; $\epsilon(340 \text{ nm}) = (1.24 \pm 0.08) \times 10^4 M^{-1} \text{ cm}^{-1}$.	76-1048
		$2k = 1.7 \times 10^{10}$	7	0.2	p.r.	D.k. in Cl^- soln.; $\epsilon(340 \text{ nm}) = 1.25 \times 10^4 M^{-1} \text{ cm}^{-1}$; obs. change in $[\text{Cl}_2^-]$ with dose.	77-1097
6.1a	$\text{Cl}_2^- \rightleftharpoons \text{Cl} + \text{Cl}^-$	$k_f = (1.1 \pm 0.4) \times 10^5 \text{ s}^{-1}$	2	0.02	p.r.	Abs. at 340 nm depends on $[\text{Cl}^-]$; $k_f = 2.1 \times 10^{10} M^{-1} \text{ s}^{-1}$.	73-1039
6.2	$\text{Cl}_2^- + \text{O}_2^- \rightarrow 2\text{Cl}^- + \text{O}_2$	$\leq 2 \times 10^9$	-	-	p.r.	D.k. in $\text{O}_2\text{-satd. } 1.3 \text{ and } 5 M \text{ LiCl}$ soln. compared with deaerated soln.	74-1149
6.2a	$\text{Cl}_2^- + \text{HO}_2 \rightarrow 2\text{Cl}^- + \text{H}^+ + \text{O}_2$	$(4.5 \pm 0.5) \times 10^9$	$0.4 M \text{ H}_2\text{SO}_4$	0.5	p.r.	Calcd. from effect of Cl^- on $G(\text{Fe}^{3+})$ in air-satd. Fe^{2+} soln.	77-1170
6.3	$\text{Cl}_2^- + \text{SCN}^- \rightarrow 2\text{Cl}^- + \text{SCN}$	$(2.9 \pm 0.3) \times 10^9$	2.6	-	p.r.	D.k. in $\text{N}_2\text{O-satd. } 6 \times 10^{-3} M \text{ Cl}^-$ soln.; addn. of $0.1 M \text{ NaClO}_4$ gave $k = 3.7 \times 10^9$; product is $(\text{SCN})_2^-$ detd. by absorption spectrum; $K(\text{ClSCN}^- + \text{SCN}^- \rightleftharpoons \text{Cl}^- + (\text{SCN})_2^-) = 3.0 \times 10^4$.	69-0565
6.4	$\text{Cl}_2^- + \text{Ce}^{3+} \rightarrow \text{Ce}^{4+} + 2\text{Cl}^-$	$(0.8 \text{ or } 4.0) \times 10^4$	-	-	$\gamma\text{-r.}$	Calcd. from $G(\text{Ce}^{3+})$ in Ce^{4+} soln. contg. $10 M \text{ LiCl}$ assuming $k(\text{Cl}_2^- + \text{Cl}_2^-) = 2 \times 10^9$.	75-0440
6.5	$\text{Cl}_2^- + \text{OCIO}$	$(1.0 \pm 0.1) \times 10^9$	5	-	f.phot.	D.k. in $\text{ClO}_2\text{-Cl}^-$ soln.	73-7043

TABLE 6. Rates of reaction of Cl_2^- in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
6.6	$\text{Cl}_2^- + \text{Co(II)} \rightarrow \text{CoCl}^{2+} + \text{Cl}^-$	$(1.4 \pm 0.2) \times 10^6$	~ 1	0.3	f.phot.	D.k. in Cl_3^- soln. (0.1 M Cl^- , 0.005 M Cl_2 , 0.1 M H^+) contg. 0.1–0.5 M Co(II) ; $\Delta H^\ddagger = 29 \pm 4 \text{ kJ mol}^{-1}$ and $\Delta S^\ddagger = -31 \pm 10 \text{ J K}^{-1} \text{ mol}^{-1}$ detd. at 13.5 to 41.5°C.; inner sphere substitution reaction.	73-7316
6.7	$\text{Cl}_2^- + \text{CoL}^{2+} \rightarrow \text{Co}^{\text{III}}$ L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene	$(1.0 \pm 0.4) \times 10^9$	1	1.1	f.phot.	D.k.	72-7506
6.8	$\text{Cl}_2^- + \text{Cr}_{\text{aq}}^{2+} \rightarrow [\text{Cr}(\text{OH}_2)_5\text{Cl}]^{2+} + \text{Cl}^-$ $\text{Cl}_2^- + \text{Cr}_{\text{aq}}^{2+} \rightarrow [\text{Cr}(\text{OH}_2)_6]^{3+} + 2\text{Cl}^-$	$(2.4 \pm 0.3) \times 10^9$ $(2.4 \pm 0.3) \times 10^9$	1	0.2	p.r.	D.k. in deaerated Cl^- soln.; $\Delta G^\circ = -263 \text{ kJ mol}^{-1}$; reaction 50% inner sphere and 50% outer sphere.	74-1104
6.9	$\text{Cl}_2^- + \text{Cu}^{2+}$	$< 1.5 \times 10^8$	—	—	p.r.	—	75-1188
6.10	$\text{Cl}_2^- + \text{CuL}^{2+} \rightarrow \text{Cu}^{\text{III}}$ L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradecane	6.5×10^8	2	0.02	p.r.	D.k.	78-1503
6.11	$\text{Cl}_2^- + \text{CuL}^{2+} \rightarrow \text{Cu}^{\text{III}}$ L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene	3.2×10^8	2	0.02	p.r.	D.k.	78-1503
6.12	$\text{Cl}_2^- + \text{Fe}^{2+} \rightarrow \text{FeCl}^{2+} + \text{Cl}^-$ (1) $\text{Cl}_2^- + \text{Fe}_{\text{aq}}^{2+} \rightarrow \text{Fe}_{\text{aq}}^{3+} + 2\text{Cl}^-$ (2) $k_1 = (4.0 \pm 0.6) \times 10^6$ $k_2 = (1.0 \pm 0.2) \times 10^7$	$(3.8 \pm 0.3) \times 10^7$ $(1.4 \pm 0.2) \times 10^7$	2.1 1 1	0.04 0.1 0.2	p.r. p.r. f.phot.	D.k. D.k. in soln. contg. 0.01 M NaCl and 0.1 M perchloric acid. D.k. in FeCl_2^{2+} soln. (10^{-3} M Fe^{3+} , 0.1 M H^+ , 0.1 M Cl^-) in pres. of Fe^{2+} ; $\Delta H^\ddagger$ for the inner sphere reaction (1) = $31.5 \pm 4 \text{ kJ mol}^{-1}$ and $\Delta S^\ddagger = -21 \pm 15 \text{ J K}^{-1} \text{ mol}^{-1}$; $\Delta H^\ddagger$ for the outer sphere reaction (2) = $22.7 \pm 4 \text{ kJ mol}^{-1}$ and $\Delta S^\ddagger = -42 \pm 15 \text{ J K}^{-1} \text{ mol}^{-1}$.	68-0313 73-1039 73-7159
6.13	$\text{Cl}_2^- + \text{Mn}_{\text{aq}}^{2+} \rightarrow \text{Mn}^{\text{III}} + 2\text{Cl}^-$	8×10^6	1	0.25	f.phot.	D.k. in Cl_3^- soln. (0.1 M Cl^- , 0.005 M Cl_2 , 0.1 M H^+) contg. 0.001–0.02 M Mn^{2+} ; $E_a = 34 \pm 4 \text{ kJ mol}^{-1}$ (16 to 40°C); reaction is inner sphere electron transfer controlled; $\Delta H^\ddagger = 32 \pm 4 \text{ kJ mol}^{-1}$, $\Delta S^\ddagger = -4 \pm 6 \text{ J K}^{-1} \text{ mol}^{-1}$.	73-7317
6.14	$\text{Cl}_2^- + \text{N}_2\text{H}_5^+$	8.0×10^6 1.4×10^7	1 6.8	~ 1 ~ 1	p.r. p.r.	D.k. in 1 M Cl^- soln. D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093 78-1093
6.15	$\text{Cl}_2^- + \text{NH}_2\text{OH}$	9.3×10^6	6.7	~ 1	p.r.	D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.16	$\text{Cl}_2^- + \text{NH}_3\text{OH}^+$	$< 10^5$	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
6.17	$\text{Cl}_2^- + \text{N}_3^-$	1.2×10^9	7	~ 1	p.r.	D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093

TABLE 6. Rates of reaction of Cl_2^- in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	l	Method	Comment	Ref.
6.18	$\text{Cl}_2^- + \text{NO}_2^-$	2.5×10^8	7	~ 1	p.r.	D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.19	$\text{Cl}_2^- + \text{NiL}^{2+} \rightarrow \text{Ni}^{\text{III}}$ L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradecane	2.0×10^9	2	~ 0.02	p.r.	D.k. in 0.005 – 0.01 M Cl^- soln.	78-1502
6.20	$\text{Cl}_2^- + \text{NiL}^{2+} \rightarrow \text{Ni}^{\text{III}}$ L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene	9.6×10^9	2	~ 0.02	p.r.	D.k. in 0.005 – 0.01 M Cl^- soln.	78-1502
6.21	$\text{Cl}_2^- + \text{NiL}^{2+} \rightarrow \text{Ni}^{\text{III}}$ L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-1,4,8,11-tetraene	8.7×10^9	2	~ 0.02	p.r.	D.k. in 0.005 – 0.01 M Cl^- soln.	78-1502
6.22	$\text{Cl}_2^- + \text{OH}^-$	7.3×10^6	10–13	–	p.r.	Unpub. data from R.W. Fessenden.	78-1093
6.23	$\text{Cl}_2^- + \text{H}_2\text{O}_2$	1.4×10^5	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
6.24	$\text{Cl}_2^- + \text{PtCl}_4^{2-} \rightarrow \text{Pt(III)}$	$(1.1 \pm 0.2) \times 10^9$	–	–	p.r.	D.k. in 0.5 M HCl; transient with $\epsilon(260 \text{ nm}) \cong 13000 M^{-1} \text{ cm}^{-1}$ was formed.	75-1188
		$(1.60 \pm 0.10) \times 10^9$	~ 1		p.r.	D.k. in 0.1 M HCl soln.; 1.0 M NaCl soln. gave $k = (1.25 \pm 0.3) \times 10^9$.	76-1055
6.25	$\text{Cl}_2^- + \text{PtL}_2^{2+} \rightarrow \text{Pt(III)}$ L = ethylenediamine	$(8.9 \pm 1.9) \times 10^9$	–	–	p.r.	D.k. in 0.5 M HCl; transient with $\epsilon(280 \text{ nm}) = 15900 M^{-1} \text{ cm}^{-1}$ was formed.	75-1188
6.26	$\text{Cl}_2^- + \text{PtLCl}^+ \rightarrow \text{Pt(III)}$ L = diethylenetriamine	$(6.4 \pm 0.4) \times 10^9$	–	–	p.r.	D.k. in 0.5 M HCl; transient with $\epsilon(280 \text{ nm}) = 17000 M^{-1} \text{ cm}^{-1}$ was formed.	75-1188
6.27	$\text{Cl}_2^- + \text{PtLCl}^+ \rightarrow \text{Pt(III)}$ L = tetraethyldiethylenetriamine	$(4.2 \pm 0.4) \times 10^8$	–	–	p.r.	D.k. in 0.5 M HCl; transient with $\epsilon(290 \text{ nm}) = 7240 M^{-1} \text{ cm}^{-1}$ was formed.	75-1188
6.28	$\text{Cl}_2^- + \text{SO}_3^{2-}$	3.3×10^7	7	~ 1	p.r.	D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.28a	$\text{Cl}_2^- + \text{Ti(III)}$	$\sim 4 \times 10^8$	~ 1	–	p.r.	D.k. at 380 nm in 0.02 M HCl contg. formic acid.	73-1057
6.29	$\text{Cl}_2^- + \text{Ti(I)} \rightarrow \text{Ti(II)}$	5×10^9	–	1	p.r.	P.b.k. at 260 nm in 0.99 M HCl soln.	74-1038
6.30	$\text{Cl}_2^- + \text{V}^{2+} \rightarrow 2\text{Cl}^- + \text{V}^{3+}$	$(1.95 \pm 0.2) \times 10^9$	1	0.2	p.r.	D.k. in deaerated Cl^- soln.; $\Delta G^\circ = -241 \text{ kJ mol}^{-1}$; outer sphere diffusion control.	74-1104
6.31	Acetanilide $\text{Cl}_2^- + \text{C}_6\text{H}_4\text{NHCCH}_3$	$\sim 2.0 \times 10^7$	7	~ 1	p.r.	D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.32	Acetic acid	$< 10^4$	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
6.33	Acetone	1.4×10^3	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
6.34	Acriflavin	$\sim 4 \times 10^9$	–	1	p.r.	D.k. at 450 nm (dye) in N_2O -satd. 1 M KCl soln.	70-0241
6.35	Acrylate ion $\text{Cl}_2^- + \text{CH}_2=\text{CHCOO}^-$	1.9×10^7	7	~ 1	p.r.	D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.36	Acrylic acid $\text{Cl}_2^- + \text{CH}_2=\text{CHCOOH}$	5.4×10^6	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
6.37	Acrylonitrile $\text{Cl}_2^- + \text{CH}_2=\text{CHCN}$	2.2×10^6	7	~ 1	p.r.	D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.38	Adenine	$< 5 \times 10^6$	2.7	–	p.r.	D.k. in 0.1 M NaCl soln.	68-0313
6.39	Alanine	1.3×10^5	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093

TABLE 6. Rates of reaction of Cl_2^- in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
6.40	Allyl alcohol $\text{Cl}_2^- + \text{CH}_2=\text{CHCH}_2\text{OH}$	5.9×10^8	1.7	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln.; at pH 7 Cl_2^- was generated by SO_4^{2-} from $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.41	<i>p</i> -Aminobenzoate ion	1.1×10^9	7	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.42	<i>p</i> -Aminobenzoic acid	2.2×10^7	1	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln.	78-1093
6.43	Anilinium ion $\text{Cl}_2^- + \text{C}_6\text{H}_5\text{NH}_3^+$	1.2×10^7	1	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln.	78-1093
6.44	Anthrasemiquinone- 2,6-disulfonate radical ion $\text{Cl}_2^- + \text{A}^- \rightarrow \text{A} + 2\text{Cl}^-$	6.5×10^8	8.0	—	f.phot.	D.k.	73-7569
6.45	Ascorbic acid $\text{Cl}_2^- + \text{AH}_2 \rightarrow$ $\cdot\text{AH} + \text{H}^+ + 2\text{Cl}^-$	6.8×10^8 $(6.0 \pm 1.0) \times 10^8$	2 2	— —	p.r. p.r.	— D.k. in N_2O -satd. 0.5 <i>M</i> Cl^- soln.	72-0266 73-3006
6.46	Benzenesulfonate ion	$< 1 \times 10^5$	7	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.47	Benzoate ion	2×10^6	7	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.48	Benzonitrile	$< 1 \times 10^5$	1.7	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln.; at pH 7 Cl_2^- was generated by SO_4^{2-} from $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.49	<i>p</i> -Bromobenzoate ion	7×10^6	7	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.50	<i>p</i> -Chlorobenzoate ion	3×10^6	7	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.51	5-Chlorouracil	1.0×10^7	2.0	—	p.r.	D.k. in 0.01 <i>M</i> NaCl soln.	72-3107
6.52	<i>p</i> -Cyanobenzoate ion	5×10^6	7	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln.	78-1093
6.53	<i>p</i> -Cyanophenol	4.0×10^7	1	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln.	78-1093
6.54	Cysteine	$(8.5 \pm 0.8) \times 10^8$	1.8	—	p.r.	D.k. in 0.1 <i>M</i> Cl^- soln.	72-0036
6.55	Cytidine	4×10^6	2.0	—	p.r.	D.k. in 0.01 <i>M</i> NaCl soln.	72-3107
6.56	Cytosine	$(9.1 \pm 0.7) \times 10^7$	2.7	—	p.r.	D.k. in 0.1 <i>M</i> NaCl soln.	68-0313
		1.0×10^7	2.0	—	p.r.	D.k. in 0.01 <i>M</i> NaCl soln.	72-3107
6.57	Deoxyadenylic acid	$< 5 \times 10^6$	2.7	—	p.r.	D.k. in 0.1 <i>M</i> NaCl soln.	68-0313
6.58	Deoxycytidylic acid	$< 5 \times 10^6$	2.7	—	p.r.	D.k. in 0.1 <i>M</i> NaCl soln.	68-0313
6.59	Deoxyguanylic acid	$(1.2 \pm 0.1) \times 10^8$	2.7	—	p.r.	D.k. in 0.1 <i>M</i> NaCl soln.	68-0313
6.60	Dithiothreitol	$(3.0 \pm 0.3) \times 10^9$	2	—	p.r.	D.k.	73-1020
6.61	Dodecyl sodium sulfate	3.9×10^6	2.0	—	p.r.	D.k. in 0.5 <i>M</i> NaCl soln.	72-3107
6.62	Ethanol $\text{Cl}_2^- + \text{CH}_3\text{CH}_2\text{OH}$	4.5×10^4	1	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln.	78-1093
6.63	Formate ion $\text{Cl}_2^- + \text{HCO}_2^-$	1.9×10^6	7	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.64	Formic acid $\text{Cl}_2^- + \text{HCOOH}$	1.0×10^6	3.7	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
		3×10^5	3	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln.	78-1093
		6.7×10^3	1	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln.	78-1093
6.65	Fumarate ion $\text{Cl}_2^- + ^-\text{OOCCH}=\text{CHCOO}^- \rightarrow$ $\text{Cl}^- + ^-\text{OOCCHCHClCOO}^-$	4×10^6	7	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$; adduct radical obs. by esr (75-5244).	78-1093
	$\text{Cl}_2^- + ^-\text{OOCCH}=\text{CHCOOH}$	2.4×10^6	3.7	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.66	Fumaric acid $\text{Cl}_2^- + \text{HOOCCH}=\text{CHCOOH}$	$\sim 2 \times 10^5$	1	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln.	78-1093
6.67	Glutamic acid	2.3×10^5	1	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln.	78-1093
6.68	Glycine	$\sim 5 \times 10^6$ $< 10^4$	9.8 1	~ 1	p.r.	D.k. in 1 <i>M</i> Cl^- soln.; at pH 9.8 Cl_2^- generated by SO_4^{2-} from $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.69	Guanine	$(8.1 \pm 0.6) \times 10^7$	2.3	—	p.r.	D.k. in 0.1 <i>M</i> NaCl soln.	68-0313

TABLE 6. Rates of reaction of Cl_2^- in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
6.70	Hexadecyltrimethyl- ammonium chloride	1.2×10^7	2.0	—	p.r.	D.k. in 0.01 M HCl soln.	72-3107
	Hexadienedioate ion <i>See</i> Muconate ion.						
	Hexadienoate ion <i>See</i> Sorbate ion.						
6.71	Hexenedioate ion $\text{Cl}_2^- + ^-\text{OOCCH}_2\text{CH}=\text{CHCH}_2\text{COO}^-$	1.6×10^8	7	~ 1	p.r.	D.k. in 0.1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.72	Histidine	$(1.4 \pm 0.1) \times 10^7$	1.8	—	p.r.	D.k. in 0.1 M Cl^- soln.	72-0036
6.73	Hydroquinone	1.4×10^9	1	~ 1	p.r.	D.k. in 0.1 M Cl^- soln.	78-1093
	$\text{Cl}_2^- + \text{OHC}_6\text{H}_4\text{OH} \rightarrow$ $2\text{Cl}^- + 2\text{H}^+ + ^\cdot\text{OC}_6\text{H}_4\text{O}^-$	1.5×10^9	9.5	~ 1	p.r.	D.k. in 0.1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.74	<i>p</i> -Hydroxybenzoate ion $\text{Cl}_2^- + \text{OHC}_6\text{H}_4\text{COO}^- \rightarrow$ $2\text{Cl}^- + \text{H}^+ + ^\cdot\text{OC}_6\text{H}_4\text{COO}^-$	2.8×10^8	7	~ 1	p.r.	D.k. in 0.1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$; product radical obs. by esr.	78-1093
6.75	<i>p</i> -Hydroxybenzoic acid $\text{Cl}_2^- + \text{OHC}_6\text{H}_4\text{COOH}$	1.3×10^8	3.1	~ 1	p.r.	D.k. in 0.1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
		1.1×10^8	1	~ 1	p.r.	D.k. in 0.1 M Cl^- soln.	78-1093
6.76	Maleate ion	3×10^6	6.5	~ 1	p.r.	D.k. in 0.1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.77	Methanol	3.5×10^3	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
6.78	Methionine	$(7 \pm 0.7) \times 10^6$	1.8	—	p.r.	D.k. in 0.1 M Cl^- soln.	72-0036
6.79	<i>p</i> -Methoxybenzoate ion $\text{Cl}_2^- + \text{CH}_3\text{OC}_6\text{H}_4\text{COO}^- \rightarrow$ $2\text{Cl}^- + \text{CH}_3\text{O}^+\text{C}_6\text{H}_4\text{COO}^-$	2.0×10^8	7	~ 1	p.r.	D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$; product radical identified by optical p.r.	78-1093
6.80	<i>p</i> -Methoxyphenol	1.1×10^9	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
6.81	2-Methyl-2-propanol	$\sim 7 \times 10^2$	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
6.82	Muconate ion	2.1×10^8	7	~ 1	p.r.	D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.83	Phenol	2.5×10^8	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
6.84	<i>p</i> -Phenoxybenzoate ion	1.5×10^8	7	~ 1	p.r.	D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.85	Phenylalanine	$(6 \pm 0.6) \times 10^6$	1.8	—	p.r.	D.k. in 0.1 M Cl^- soln.	72-0036
6.86	Polyoxyethylene(15)nonyl phenol	2.1×10^8	2.0	—	p.r.	D.k. in 0.01 M NaCl soln.	72-3107
6.87	2-Propanol	1.2×10^5	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
		$(1.9 \pm 0.3) \times 10^5$	—	—	p.r.	D.k. in 0.5 M HCl.	75-1188
6.88	Propionic acid	2.2×10^3	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
6.89	Salicylic acid	1.1×10^8	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
6.90	Serine	1.2×10^5	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
6.91	Sorbate ion	6.8×10^8	7	~ 1	p.r.	D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.92	Succinic acid	$\sim 8 \times 10^2$	1	~ 1	p.r.	D.k. in 1 M Cl^- soln.	78-1093
6.93	Terephthalate ion	6×10^6	7	~ 1	p.r.	D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.94	Tetramethylpiperidone <i>N</i> -oxyl (TAN)	1.4×10^9	2	—	p.r.	D.k. at 350 nm.	71-0618
6.95	Thymidylic acid	$(4.4 \pm 0.3) \times 10^7$	2.7	—	p.r.	D.k. in 0.1 M NaCl soln.	68-0313
6.96	Thymine	$(1.2 \pm 0.1) \times 10^8$	2.7	—	p.r.	D.k. in 0.1 M NaCl soln.	68-0313
		7.0×10^7	2.0	—	p.r.	D.k. in 0.01 M NaCl soln.	72-3107
6.97	<i>p</i> -Toluate ion	5×10^6	7	~ 1	p.r.	D.k. in 1 M Cl^- soln. contg. $\text{S}_2\text{O}_8^{2-}$.	78-1093
6.98	Tryptophan	$(2.6 \pm 0.3) \times 10^9$	1.8	—	p.r.	D.k. in 0.1 M Cl^- soln.	72-0036
6.99	Tyrosine	$(2.7 \pm 0.3) \times 10^8$	1.8	—	p.r.	D.k. in 0.1 M Cl^- soln.	72-0036
6.100	Uracil	$(4.1 \pm 0.3) \times 10^7$	2.7	—	p.r.	D.k. in 0.1 M NaCl soln.	68-0313
		3.5×10^7	2.0	—	p.r.	D.k. in 0.01 M NaCl soln.	72-3107
		3.7×10^7	6	—	p.r.	—	75-5244

TABLE 7. Rates of reaction of Br_2^- in aqueous solution

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
7.1	$\text{Br}_2^- + \text{Br}_2^- \rightarrow \text{Br}_3^- + \text{Br}^-$	$k = 1.8 \times 10^9$	2	0.03	p.r.	D.k. in aerated $10^{-4} M \text{ Br}^- - 10^{-3} M \text{ Br}_2$; $k/\epsilon = 2.2 \times 10^5 \text{ cm s}^{-1}$; $\epsilon(360 \text{ nm}) = 8200 M^{-1} \text{ cm}^{-1}$.	65-0382
		$k = (1.8 \pm 0.3) \times 10^9$	7	0.01	p.r.	D.k. in N_2O -satd. $10^{-2} M \text{ Br}^-$; $\epsilon(360 \text{ nm}) = 9600 \pm 800 M^{-1} \text{ cm}^{-1}$.	65-0383
		$2k = (3.3 \pm 1.0) \times 10^9$	7	<0.01	p.r.	D.k. in $10^{-4} - 10^{-2} M \text{ Br}^-$ soln.; $\epsilon(365 \text{ nm}) = 7800 \pm 2000 M^{-1} \text{ cm}^{-1}$.	66-0425
		$2k = (3.7 \pm 0.6) \times 10^9$	6	0.02	f.phot.	D.k. in $0.02 M \text{ Br}^-$ or $\text{HgBr}_2 - \text{HgBr}_4^{2-}$ soln.; $\epsilon(360 \text{ nm}) = (7.8 \pm 0.2) \times 10^4 M^{-1} \text{ cm}^{-1}$.	67-7171
		$k = 2.8 \times 10^9$	12	0.02	p.r.	D.k. in N_2O -satd. $10^{-2} M \text{ Br}^-$ soln.; $\epsilon(360 \text{ nm}) = 8200 M^{-1} \text{ cm}^{-1}$.	68-0153
		$2k = 5.3 \times 10^9$	1	0.2	f.phot.	D.k. in FeBr^{2+} soln. ($10^{-3} M \text{ Fe}^{3+}$, $0.1 M \text{ H}^+$, $0.1 M \text{ Br}^-$); $\epsilon(366 \text{ nm}) = 7.8 \times 10^3 M^{-1} \text{ cm}^{-1}$.	73-7159
		1.6×10^9	-	-	f.phot.	Calcd. from assumed mechanism for d.k. at 350 nm in Br^- soln. 10^{-5} to $10^{-1} M$.	75-7346
		$2k = (3.1 \pm 0.9) \times 10^9$	-	-	p.r.	D.k. in Br^- soln.; $\epsilon(360 \text{ nm}) = (8.56 \pm 0.24) \times 10^3 M^{-1} \text{ cm}^{-1}$.	76-1048
		2.4×10^9	-	0.02	p.r.	D.k. in N_2O -satd. Br^- soln.; $\epsilon(360 \text{ nm}) = 9600 M^{-1} \text{ cm}^{-1}$; rates in micellar systems also detd.	76-1058
7.1a	$\text{Br}_2^- \rightleftharpoons \text{Br} + \text{Br}^-$	$k_i = (2.5 \pm 1) \times 10^4 \text{ s}^{-1}$	2	-	p.r.	Calcd. from dependence of $[\text{Br}_2^-]$ on $[\text{Br}^-]$; $k_i = (5.4 \pm 3) \times 10^9$ and $K^{-1} = (2.2 \pm 1) \times 10^5 M^{-1}$.	65-0383
		$k_i = (7 \pm 2) \times 10^5 \text{ s}^{-1}$	~ 2	0.12	f.phot.	Calcd. from buildup of Br_2^- in $\text{Br}^- - \text{Br}_2$ soln.; $k_i = (1.1 \pm 0.1) \times 10^{10}$.	75-7222
7.2	$\text{Br}_2^- + \text{Br} \rightarrow \text{Br}_3^-$	1.37×10^9	~ 7	-	f.phot.	Calcd. from assumed mechanism for d.k. at 350 nm in Br^- soln. 10^{-5} to $10^{-1} M$.	75-7346
7.3	$\text{Br}_2^- + \text{H} \rightarrow \text{H}^+ + 2\text{Br}^-$	$(7 \pm 1) \times 10^9$	2	-	p.r.	-	70-7726
7.4	$\text{Br}_2^- + \text{HO}_2^- \rightarrow \text{Br}_2 + \text{HO}_2^-$	$(1.6 \pm 0.5) \times 10^9$	2	-	p.r.	D.k. in air-satd. $0.1 M \text{ Br}^-$ soln.	65-0383
7.5	$\text{Br}_2^- + \text{BrO}^- \rightarrow \text{BrO} + 2\text{Br}^-$	$(8.0 \pm 0.7) \times 10^7$	-	0.02	p.r.	D.k. in $\text{Br}^- - \text{BrO}^-$ soln.	68-0153
7.6	$\text{Br}_2^- + \text{BrO}_2^- \rightarrow \text{BrO} + \text{BrO}^- + \text{Br}^-$	$(8.0 \pm 0.8) \times 10^7$	-	1	p.r.	D.k. in $\text{Br}^- - \text{BrO}_2^-$ soln.	68-0153
7.7	$\text{Br}_2^- + \text{BrO}_3^-$	no reaction	-	-	p.r.	D.k. of Br_2^- unaffected by BrO_3^- .	68-0153
7.8 *	$\text{Br}_2^- + \text{ClO}_2$	$(1.2 \pm 0.1) \times 10^9$	-	-	f.phot.	D.k. in $\text{Br}^- - \text{ClO}_2$ soln.	73-7043
7.9	$\text{Br}_2^- + \text{Co}^{2+} \rightarrow \text{Co}^{2+} + 2\text{Br}^-$	$(1.0 \pm 0.3) \times 10^{10}$	-	-	f.phot.	D.k. in Br^- soln. contg. Co^{2+} ; $e_{\text{aq}}^- + \text{Co}^{2+} \rightarrow \text{Co}^+$.	70-7726
7.10	$\text{Br}_2^- + \text{Co}^{2+}$	-	-	-	f.phot.	No reaction.	70-7726 73-7316
7.11	$\text{Br}_2^- + \text{CoL}^{2+} \rightarrow \text{Co}^{\text{III}}$	$(1.4 \pm 0.1) \times 10^9$ 2.0×10^9	1 9.2	0.2 0.1	f.phot. p.r.	D.k. in Br^- soln. D.k. in N_2O -satd. Br^- soln.	72-7506 76-1203

L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene

TABLE 7. Rates of reaction of Br_2^- in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
7.12	$\text{Br}_2^- + \text{Cr}_{\text{aq}}^{2+} \rightarrow [\text{Cr}(\text{OH}_2)_5\text{Br}]^{2+} + \text{Br}^-$	$(1.9 \pm 0.2) \times 10^9$	1	0.2	p.r.	D.k. in deaerated Br^- soln.; $\Delta G^\circ = -213 \text{ kJ mol}^{-1}$; inner sphere diffusion controlled mechanism.	74-1104
7.13	$\text{Br}_2^- + \text{CuL}^{2+} \rightarrow \text{Cu}^{\text{III}}$ L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradecane	2×10^7	6	0.02	p.r.	D.k.	78-1503
7.14	$\text{Br}_2^- + \text{CuL}^{2+} \rightarrow \text{Cu}^{\text{III}}$ L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene	1×10^7	6	0.02	p.r.	D.k.	78-1503
7.15	$\text{Br}_2^- + \text{Fe}_{\text{aq}}^{2+} \rightarrow \text{Br}^- + \text{FeBr}^{2+}$	$(3.6 \pm 0.4) \times 10^6$	1	0.2	f.phot.	D.k. at 366 nm (Br_2^-) or p.b.k. at 405 nm (FeBr^{2+}); $\Delta H^\ddagger = 25.2 \pm 2 \text{ kJ mol}^{-1}$; $\Delta S^\ddagger = -42 \pm 12 \text{ J K}^{-1} \text{ mol}^{-1}$; inner sphere substitution controlled mechanism.	73-7159
7.16	$\text{Br}_2^- + \text{ferricenium ion} \rightarrow \text{ferrocene}$	3×10^8	—	—	p.r.	—	74-1182
7.17	$\text{Br}_2^- + \text{H}_2\text{O}_2 \rightarrow \text{H}^+ + 2\text{Br}^- + \text{HO}_2$	$\leq 10^3$	7	—	p.r.	D.k. in N_2O -satd. 1 M NaBr soln.	78-1093
7.18	$\text{Br}_2^- + \text{Mn}^{2+} \rightarrow 2\text{Br}^- + \text{Mn}^{3+}$	6.3×10^6	1	0.25	f.phot.	D.k. in Br_3^- soln. (0.1 M Br^- , 5×10^{-5} M Br_2 and 0.1 M H^+); $E_a = 36 \pm 4 \text{ kJ mol}^{-1}$; $\Delta H^\ddagger = 33.6 \pm 4 \text{ kJ mol}^{-1}$; $\Delta S^\ddagger = -3 \pm 6 \text{ J K}^{-1} \text{ mol}^{-1}$; inner sphere electron transfer.	73-7317
7.19	$\text{Br}_2^- + \text{NiL}^{2+} \rightarrow \text{Ni}^{\text{III}}$ L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradecane	3.4×10^9	2	—	p.r.	D.k. in 0.005 – 0.01 M Br^- soln.	78-1502
7.20	$\text{Br}_2^- + \text{NiL}^{2+} \rightarrow \text{Ni}^{\text{III}}$ L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene	9.8×10^9	2	—	p.r.	D.k. in 0.005 – 0.01 M Br^- soln.	78-1502
7.21	$\text{Br}_2^- + \text{NiL}^{2+} \rightarrow \text{Ni}^{\text{III}}$ L = 5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraazacyclotetradeca-1,4,8,11-tetraene	9.5×10^9	2	—	p.r.	D.k. in 0.005 – 0.01 M Br^- soln.	78-1502
7.22	$\text{Br}_2^- + \text{PtCl}_4^{2-} \rightarrow \text{Pt(III)}-\text{D}$	$(2.8 \pm 0.2) \times 10^8$	—	$\rightarrow 0$	p.r.	D.k. in N_2O -satd. 0.1 or 0.01 M Br^- soln. contg. PtCl_4^{2-} ; $\text{Pt(III)}-\text{D}$ ($\epsilon_{310 \text{ nm}} = 8.4 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$) may be $\text{PtBr}_n\text{Cl}_{4-n}^-$.	76-1055
7.23	$\text{Br}_2^- + \text{Ru}(\text{bipy})_3^{2+} \rightleftharpoons [\text{Ru}(\text{bipy})_3^{3+}\text{Br}] + \text{Br}^-$	$(3.1 \pm 0.5) \times 10^9$	1 M H_2SO_4	—	f.phot.	D.k. in 10^{-3} M Br^- soln.	73-7066
7.24	$\text{Br}_2^- + \text{SCN}^- \rightarrow 2\text{Br}^- + \text{SCN}$ or $\rightarrow \text{Br}^- + \text{BrSCN}^-$	1.9×10^9	—	—	p.r.	D.k. in N_2O -satd. 5×10^{-3} M Br^- soln. contg. 10^{-5} – 2×10^{-4} M SCN^- as well as p.b.k. at 500 nm ($\text{SCN})_2^-$; $K(\text{BrSCN}^- + \text{SCN}^- \rightleftharpoons \text{Br}^- + (\text{SCN})_2^-) = 1.1 \times 10^2$.	69-0180
7.25	$\text{Br}_2^- + \text{Tl}^+$		1	0.01	f.phot.	no reaction	74-7625
7.26	$\text{Br}_2^- + \text{UO}_2^{2+} \rightarrow 2\text{Br}^- + \text{UO}_2^{2+}$	$0.33 \times 10^3 \text{ s}^{-1}$	1.7	—	f.phot.	D.k. in Br^- – U(VI) soln.; react. probably reoxid. of U(V) to U(VI).	76-7279
7.27	$\text{Br}_2^- + \text{V}^{2+} \rightarrow 2\text{Br}^- + \text{V}^{3+}$	$(1.48 \pm 0.2) \times 10^9$	1	0.2	p.r.	D.k. in Br^- soln.; $\Delta G^\circ = -191 \text{ kJ mol}^{-1}$; outer sphere diffusion control.	74-1104

TABLE 7. Rates of reaction of Br_2^- in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
7.28	Acridflavin	$(3.7 \pm 0.6) \times 10^9$	—	—	p.r.	D.k. of Br_2^- in N_2O -satd. $10^{-3} M \text{Br}^-$ soln. as well as dye bleaching at 450 nm.	70-0241
7.29	Alcohol dehydrogenase (yeast)	2.6×10^9	7	—	p.r.	D.k. in $0.05 M \text{Br}^-$ soln.; $k = 1.0 \times 10^9$ for horse liver enzyme (78-3007).	74-1125
7.30	Ascorbic acid $\text{Br}_2^- + \text{AH}_2 \rightarrow \text{AH} + \text{H}^+ + 2\text{Br}^-$	$(1.1 \pm 0.4) \times 10^8$ $(1.1 \pm 0.4) \times 10^9$ $(8.7 \pm 0.9) \times 10^8$	2 7 7.2	— — 0.5	p.r. p.r.	$pK = 4.17, 11.57$. D.k. in N_2O -satd. Br^- soln.	72-0266 73-3006
7.31	<i>p</i> -Bromophenoxide ion	2.75×10^8	12.5	—	p.r.	—	74-3052
7.32	Carboxypeptidase A	$(1 \text{ to } 2.5) \times 10^9$	8-11	—	p.r.	D.k. in $0.05 M \text{Br}^-$ soln.	73-1060
7.33	<i>p</i> -Chlorophenoxide ion	1.73×10^8	12.5	—	p.r.	—	74-3052
7.34	α -Chymotrypsin	1.6×10^9	6.7	—	p.r.	D.k. in N_2O -satd. $0.04 M \text{Br}^-$; mol. wt. = 20,000.	74-1096
7.34a	Concanavalin A	7×10^9	7.1	—	p.r.	P.b.k. at 530 nm in N_2O -satd. $10^{-2} M \text{KBr}$ soln.	78-3005
7.35	<i>p</i> -Cyanophenoxide ion	1.8×10^8	12.5	—	p.r.	—	74-3052
7.36	Cysteamine	3×10^8	—	—	p.r.	—	74-1168
7.37	Cysteine	$(1.8 \pm 0.2) \times 10^8$	6.6	0.1	p.r.	D.k. in N_2O -satd. Br^- soln.; k increases with pH.	72-0036
7.38	Cytochrome C (ferro) $\text{Br}_2^- + \text{Fe}^{2+} \text{cytC} \rightarrow 2\text{Br}^- + \text{Fe}^{3+} \text{cytC}$	1.5×10^9	7	—	p.r.	—	77-3091
7.39	Diethyl disulfide $\text{Br}_2^- + \text{RSSR} \rightarrow 2\text{Br}^- + \text{RSSR}^{\cdot+}$	1.8×10^9	4-5	0.02	p.r.	D.k. in N_2O -satd. Br^- soln., as well as p.b.k. at 430 nm ($\text{RSSR}^{\cdot+}$).	76-1143
7.39a	Diethyl sulfide $\text{Br}_2^- + (\text{C}_2\text{H}_5)_2\text{S} \rightarrow (\text{C}_2\text{H}_5)_2\text{SBr}^{\cdot+} + \text{Br}^-$	$\sim 2 \times 10^9$	3.9	0.2	p.r.	D.k. at 360 nm in N_2O -satd. KBr soln.	77-1164
7.40	3,5-Diiodotyrosine	$< 1 \times 10^8$ $\sim 1.3 \times 10^9$	6 8	—	p.r.	D.k. in N_2O -satd. $0.04 M \text{Br}^-$ soln.; values from graph.	73-1067
7.41	Dimethyl disulfide	2.2×10^9	4-5	0.02	p.r.	D.k. in N_2O -satd. Br^- soln., as well as p.b.k. at 430 nm ($\text{RSSR}^{\cdot+}$).	76-1143
7.42	<i>m</i> -Fluorotyrosine	$< 1 \times 10^8$ $\sim 1.2 \times 10^9$	6 11	—	p.r.	D.k. in N_2O -satd. $0.04 M \text{Br}^-$ soln.; values from graph.	73-1067
7.43	Formate ion	$< 10^3$	7	—	p.r.	D.k. in N_2O -satd. $1 M \text{Br}^-$ soln.	78-1093
7.44	d-Guanylic acid $\text{Br}_2^- + \text{GMP} \rightarrow 2\text{Br}^- + \text{GMP}^{\cdot+}$	4×10^7 2×10^8	— 12	—	p.r.	—	74-1168, 74-1181
7.45	Histidine	$(1.5 \pm 0.2) \times 10^7$	7.6	0.1	p.r.	D.k. in N_2O -satd. Br^- soln.	72-0036
7.46	Hydroquinone	1×10^8	—	—	p.r.	—	74-1168
7.47	<i>p</i> -Hydroxybenzoate ion	2.28×10^8	12.5	—	p.r.	—	74-3052
7.48	Lactate dehydrogenase	$(5.5 \pm 0.5) \times 10^9$	7.2	0.005	p.r.	D.k. in N_2O -satd. Br^- soln.	77-1132
7.49	Methionine	$(1.1 \pm 0.1) \times 10^7$	7.3	0.1	p.r.	D.k. in N_2O -satd. Br^- soln.; k increases with pH.	72-0036
7.50	<i>p</i> -Methylphenoxide ion	3.74×10^8	12.5	—	p.r.	—	74-3052
7.51	Nicotinamide-adenine dinucleotide, reduced $\text{Br}_2^- + \text{NADH} \rightarrow 2\text{Br}^- + \text{H}^+ + \text{NAD}^{\cdot+}$	9.0×10^8	—	—	p.r.	P.b.k. at ~ 400 nm in N_2O -satd. $0.1 M \text{Br}^-$ soln.	71-0158
7.52	Papain	1.05×10^9 2.30×10^9	7 11.5	—	p.r.	D.k. in $0.05 M \text{Br}^-$ soln.; activated enzyme used.	74-1026

TABLE 7. Rates of reaction of Br_2^- in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
7.53	Phenol	6×10^6	6	—	p.r.	—	74-3003
7.54	Phenoxide ion	5×10^8	10	—	p.r.	—	74-3003
		2.88×10^8	12.5	—	p.r.	—	74-3052
7.55	Phenylalanine	$< 1 \times 10^6$	7.0	0.1	p.r.	D.k. in N_2O -satd. Br^- soln.	72-0036
7.56	Promethazine	4×10^9	—	—	p.r.	—	74-1168
7.57	Pyrene (triplet state) $\text{Br}_2^- + {}^3\text{Py} \rightarrow$ $2\text{Br}^- + \text{Py}^+$	1×10^9	—	—	p.r.	D.k. at 414 nm (${}^3\text{Py}$) or 360 nm (Br_2^-) or p.b.k. at 448 nm (Py^+) in 0.02 M Br^- contg. $10^{-2} M$ hexadecyltrimethyl ammonium bromide; triplet state formed by f.phot.	76-1181
7.58	Ribonuclease	4.6×10^9	4.5	—	p.r.	D.k. in $10^{-2} M \text{Br}^-$ soln.	72-0037
7.59	Subtilisin (serine protease)	1×10^9	7	—	p.r.	D.k. in N_2O -satd. 0.04 $M \text{Br}^-$ soln.; k increases with pH in alk. soln.	73-1147, 74-1119
7.60	Superoxide dismutase	$(0.44 \pm 0.03) \times 10^9$	< 10	—	p.r.	Bovine enzyme; $k = (1.18 \pm 0.04) \times 10^9$ for human enzyme; k decreases at higher pH.	73-1148, 74-3081
7.61	Tetramethylpiperidone <i>N</i> -oxyl (TAN)	1.6×10^9	5-6	—	p.r.	D.k. at 380 nm.	71-0618
7.62	Thymine	$< 1 \times 10^7$ 2×10^8	— 12	—	p.r.	—	74-1168
7.63	Trypsin	$(2.55 \pm 0.3) \times 10^9$	7.0- 8.0 11.5	—	p.r.	D.k. in N_2O -satd. 0.04 $M \text{Br}^-$.	73-1067
7.64	Tryptophan	$(5.30 \pm 0.5) \times 10^9$ $(7.7 \pm 0.8) \times 10^8$	7.0	0.1	p.r.	D.k. in N_2O -satd. Br^- soln.	72-0036
7.65	Tyrosine $\text{Br}_2^- + \text{HOC}_6\text{H}_4\text{CH}_2\text{CHNH}_2\text{COOH} \rightarrow$ $2\text{Br}^- + \text{H}^+ +$ $\cdot\text{OC}_6\text{H}_4\text{CH}_2\text{CHNH}_2\text{COOH}$	$(2.0 \pm 0.2) \times 10^7$ $< 1 \times 10^7$ $\sim 1.4 \times 10^9$	7.5 6 12	0.1 —	p.r. p.r.	D.k. in N_2O -satd. Br^- soln.; k increases with pH. D.k. in N_2O -satd. 0.04 $M \text{Br}^-$ soln.; values from graph.	72-0036 73-1067
		5×10^8	12	—	p.r.	—	74-1168
7.66	Uracil	$< 1 \times 10^7$ 2×10^8	— 12	—	p.r.	—	74-1168

TABLE 8. Rates of reaction of I_2^- in aqueous solution

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
8.1	$I_2^- + I_2^- \rightarrow I_3^- + I^-$	$2k = (7.7 \pm 1.5) \times 10^9$	1.4–6	–	f.phot.	D.k. at 404.7 nm in 3×10^{-5} to $10^{-3} M I^-$ soln.; $\epsilon(404.7 \text{ nm}) = 11,700 M^{-1} \text{ cm}^{-1}$.	57–7007
		$2k = (1.17 \pm 0.2) \times 10^{10}$ $(8.7 \pm 0.5) \times 10^9$ $(6.7 \pm 0.6) \times 10^9$	6.0 1.1 6.5	–	f.phot.	D.k. at 385 nm in N_2O -satd. $10^{-3} M I^-$ soln., or HgI_4^{2-} soln. ($10^{-4} M HgI_2 + 10^{-2} M I^-$); $\epsilon(385 \text{ nm}) = 1.4 \times 10^4 M^{-1} \text{ cm}^{-1}$.	67–7171
		$2k = (9.0 \pm 2.0) \times 10^9$	alk.	–	p.r.	Computer analysis of abs. at 335 and 390 nm in N_2O -satd. I^- soln.; $\epsilon(390 \text{ nm}) = 15,600 M^{-1} \text{ cm}^{-1}$; $\epsilon(335 \text{ nm}) = 8200 M^{-1} \text{ cm}^{-1}$.	76–1105
8.2	$I_2^- \rightleftharpoons I + I^-$	$k_f = 6 \times 10^4 s^{-1}$	–	–	p.r.	$K^{-1} = 1.13 \times 10^5 M^{-1}$; $k_r = 7.6 \times 10^9 M^{-1} s^{-1}$ detd. by effect of $[I^-]$ on $[I_2^-]$.	68–0375
		$k_f = 1.7 \times 10^6 s^{-1}$	–	–	f.phot.	D.k. at 385 nm in HgI_2 - I^- soln.	74–7224
		$k_f = (9 \pm 1) \times 10^5 s^{-1}$	–	–	f.phot.	Detd. in I_3^- soln.	74–7554
8.3	$I_2^- + Co^{2+}$	–	–	–	f.phot.	no reaction	73–7316
8.4	$I_2^- + CoL^{2+} \rightarrow CoLI_2^+$ $L = 5,7,7,12,14,14\text{-hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene}$	$(7.2 \pm 0.8) \times 10^9$	2	0.03	f.phot.	D.k.; I_2^- from ion pair $Co(NH_3)_6^{3+}$, I^- .	72–7506
8.5	$I_2^- + Co(NH_3)_5I^{2+} (+H^+) \rightarrow I_3^- + Co^{2+} + 5NH_4^+$	$\sim 2.5 \times 10^4$	–	–	f.phot.	Estd. from intensity dependence of $\Phi(Co^{II})$.	72–7506
8.6	$I_2^- + Cr^{2+} \rightarrow [Cr(OH_2)_5I]^{2+} + I^-$	$(1.5 \pm 0.2) \times 10^9$	1	0.2	p.r.	D.k.; $\Delta G^\circ = -138 \text{ kJ mol}^{-1}$; inner sphere diffusion control.	74–1104
8.7	$I_2^- + Fe^{2+} \rightarrow FeI^{2+} + I^-$	3.6×10^6	–	–	–	$\Delta G^\circ = -22 \text{ kJ mol}^{-1}$; unpub. work cited; inner sphere substitution.	74–1104
8.8	$I_2^- + IO^- \rightarrow IO^- + 2I^-$	$(5.0 \pm 0.6) \times 10^7$	13.6	–	f.phot.	D.k. in $(1-3) \times 10^{-4} M IO^-$ and I^- soln.	70–0018
8.9	$I_2^- + IOH^- \rightarrow I_3^- + OH^-$	$(1.8 \pm 0.8) \times 10^{10}$	alk	–	p.r.	Computer anal. of abs. at 335 nm and 390 nm in N_2O -satd. I^- soln.	76–1105
8.10	$I_2^- + V^{2+} \rightarrow 2I^- + V^{3+}$	$(1.43 \pm 0.2) \times 10^8$	1	0.2	p.r.	D.k.; $\Delta G^\circ = -116 \text{ kJ mol}^{-1}$; outer sphere.	74–1104
8.11	Acridflavin	$< 2 \times 10^8$	–	–	p.r.	Decay of I_2^- only slightly increased in presence of dye.	70–0241
8.12	Alcohol dehydrogenase (yeast)	$\sim 1.2 \times 10^9$	–	–	p.r.	Est. from d.k. in N_2O -satd. I^- soln.	73–1065

TABLE 8. Rates of reaction of I_2^- in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
8.13	Aldolase (rabbit muscle)	$\sim 3 \times 10^8$	—	—	p.r.	Est. from d.k. in N_2O -satd. I^- soln.	73-1065
8.14	Ascorbate ion	$(1.4 \pm 0.3) \times 10^8$	7	—	p.r.	—	72-0266
8.15	Ascorbic acid	5×10^6	2	—	p.r.	—	72-0266
8.16	<i>p</i> -Bromophenoxide ion	5.0×10^7	12.5	—	p.r.	At pH 7 $k \cong 5 \times 10^6$.	74-3052
8.17	<i>p</i> -Chlorophenoxide ion	5.8×10^7	12.5	—	p.r.	—	74-3052
8.18	Cysteine	$(1.1 \pm 0.1) \times 10^8$	6.8	0.1	p.r.	D.k. in N_2O -satd. I^- soln.; k increases with pH.	72-0036
8.19	Dithiothreitol	$(1.9 \pm 0.2) \times 10^7$	7	—	p.r.	D.k.	73-1020
8.20	Histidine	$< 1 \times 10^6$	7.0	0.1	p.r.	D.k. in N_2O -satd. I^- soln.	72-0036
8.21	Lactate dehydro- genase	$(4.3 \pm 0.5) \times 10^9$	7.2	0.005	p.r.	D.k. at N_2O -satd. I^- soln.	77-1132
8.22	Methionine	$< 1 \times 10^6$	7.0	0.1	p.r.	D.k. at N_2O -satd. I^- soln.	72-0036
8.23	<i>p</i> -Methylphenoxide ion	9.8×10^7	12.5	—	p.r.	—	74-3052
8.24	Nicotinamide-adenine dinucleotide, reduced $I_2^- + NADH \rightarrow$ $2I^- + NAD\cdot + H^+$	$\sim 5 \times 10^7$	—	0.1	p.r.	D.k. at 370 nm in N_2O -satd. I^- soln.	71-0158
8.25	Phenoxide ion	5.7×10^7	12.5	—	p.r.	—	74-3052
8.26	Phenylalanine	$< 1 \times 10^6$	7.0	0.1	p.r.	D.k. in N_2O -satd. I^- soln.	72-0036
8.27	Tetramethylpiperidone <i>N</i> -oxyl (TAN)	1.7×10^9	5-6	—	p.r.	D.k. at 390 nm; final product may be I_3^- .	71-0618
8.28	Trypsin	$(1 \text{ to } 6) \times 10^8$	11-12	—	p.r.	D.k. in N_2O -satd. 0.04 $M I^-$ soln.; values from graph.	73-1067
8.29	Tryptophan	$< 1 \times 10^6$	7.0	0.1	p.r.	D.k. in N_2O -satd. I^- soln.; k increases with pH.	72-0036
8.30	Tyrosine	$< 1 \times 10^6$	7.0	0.1	p.r.	D.k. in N_2O -satd. I^- soln.	72-0036

TABLE 9. Rates of reaction of $(\text{SCN})_2^-$ in aqueous solution

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
9.1	$2(\text{SCN})_2^- \rightarrow (\text{SCN})_2 + 2\text{SCN}^-$	$2k = 2.9 \times 10^9$	7	0.01	p.r.	D.k. in presence or absence of O_2 ; $\epsilon(500 \text{ nm}) = 7100 M^{-1} \text{ cm}^{-1}$.	65-0386
		$2k \approx 3 \times 10^9$	-	$\rightarrow 0$	p.r.	Second order decay in N_2O -satd. CNS^- soln.; value from graph; $\epsilon = 7600 M^{-1} \text{ cm}^{-1}$.	68-0375
		$2k = (2.4 \pm 0.4) \times 10^9$	~ 5.7	0	p.r.	D.k. in O_2 -satd. soln. of $\leq 0.5 M \text{ CNS}^-$; ϵ not given.	72-0475
		$2k = (2.5 \pm 0.2) \times 10^9$	-	0.005	p.r.	Addn. of Cl^- increases k to 3.3×10^9 at $5 M \text{ LiCl}$.	75-1119
9.1a	$(\text{SCN})_2^- \rightleftharpoons \text{SCN} + \text{SCN}^-$	$k_f = 3.4 \times 10^4 s^{-1}$	-	-	p.r.	$K^{-1} = 2 \times 10^5 M$. $k_r = 7.0 \times 10^9 M^{-1} s^{-1}$ detd. by effect of $[\text{SCN}^-]$ on $[(\text{SCN})_2^-]$.	68-0375
		$k_f \approx 3 \times 10^3 s^{-1}$	1,6	-	f.phot.	First-order decay in N_2 or N_2O -satd. 2.5×10^{-5} to $2 \times 10^{-4} M \text{ CNS}^-$ soln.; second order in presence of O_2 with $2k/\epsilon = 2.0 \times 10^6$.	68-7072
9.2	$(\text{SCN})_2^- + \text{I}^- \rightarrow \text{ISCN}^- + \text{SCN}^-$	1.6×10^9	-	0.005	p.r.	D.k. in N_2O -satd. CNS^- soln.	70-0164
9.3	$(\text{SCN})_2^- + \text{NiL}^{2+} \rightarrow \text{Ni}^{\text{III}}$ $\text{L} = 5,7,7,12,14,14\text{-hexamethyl-1,4,8,11-tetraazacyclotetradecane}$	$(1.0 \pm 0.4) \times 10^9$	2	~ 0.02	p.r.	D.k. in $0.005\text{-}0.01 M \text{ SCN}^-$ soln.	78-1502
9.4	$(\text{SCN})_2^- + \text{UO}_2^{2+} \rightarrow 2\text{SCN}^- + \text{UO}_2^{2+}$	$(1.46 \pm 0.34) \times 10^9$	1.0	-	f.phot.	D.k. in $\text{CNS}^- - \text{U(VI)}$ soln.; reaction probably reoxid. of U(V) to U(VI) ; assume $[(\text{SCN})_2^-] = [\text{UO}_2^{2+}]$.	76-7279
9.5	Alcohol dehydrogenase (yeast)	9.6×10^8	7	-	p.r.	D.k.; $k = 5.6 \times 10^8$ in horse liver enzyme (78-3007).	74-1125
9.6	Aldolase	$< 5 \times 10^9$	7	0.1	p.r.	D.k. in SCN^- soln.; value from graph; k increases with pH.	75-3058
9.7	D-Amino acid oxidase	6.6×10^8	-	-	p.r.	D.k. in N_2O -satd. $0.05 M \text{ SCN}^-$ soln.; mol. wt. 50,000.	77-1198
9.8	Ascorbic acid $(\text{SCN})_2^- + \text{AH}_2 \rightarrow \text{H}^+ + 2\text{SCN}^- + \cdot\text{AH}$	$(1.0 \pm 0.5) \times 10^7$	1.8	-	p.r.	P.b.k. at 380 nm in $0.1 M \text{ SCN}^-$ soln.	72-0266
		$< 1.0 \times 10^7$	-	-	p.r.	D.k. in N_2O -satd. $10^{-2} M \text{ SCN}^-$ soln.	73-3006
	$(\text{SCN})_2 + \text{AH}^- \rightarrow \text{H}^+ + 2\text{SCN}^- + \text{A}^-$	$(6.0 \pm 1) \times 10^8$	7	0.1	p.r.	D.k. in SCN^- soln., as well as p.b.k. at 360 nm.	72-0266
		$(4.8 \pm 0.8) \times 10^8$	6.7	0.01	p.r.	D.k. in N_2O -satd. $10^{-2} M \text{ SCN}^-$ soln.	73-3006
9.9	<i>p</i> -Bromophenoxide ion	3.10×10^8	12.5	-	p.r.	-	74-3052
9.10	Carboxypeptidase A	$(4 \text{ to } 12) \times 10^8$	8-11	-	p.r.	D.k.; values from graph.	73-1060
9.11	<i>p</i> -Chlorophenoxide ion	3.40×10^8	12.5	-	p.r.	-	74-3052
9.12	α -Chymotrypsin	9×10^8	6.7	0.04	p.r.	D.k. in N_2O -satd. SCN^- soln.	74-1096
9.13	<i>p</i> -Cyanophenoxide ion	6.3×10^7	12.5	-	p.r.	-	74-3052
9.14	Cysteamine	8×10^7	-	-	p.r.	-	74-1168
9.15	Cysteine	$(5 \pm 0.5) \times 10^7$	6.6	0.1	p.r.	D.k. in N_2O -satd. SCN^- soln.; k increases with pH.	72-0036
9.16	3,4-Dihydroxyphenyl-alanine	3×10^7	-	-	p.r.	-	74-1168
9.17	Dithiothreitol	$(2.1 \pm 0.2) \times 10^7$	7	0.1	p.r.	D.k. in SCN^- soln.	73-1020

TABLE 9. Rates of reaction of $(\text{SCN})_2^-$ in aqueous solution—Continued

No.	Reaction	$k(M^{-1} s^{-1})$	pH	I	Method	Comment	Ref.
9.18	Formate ion	$\leq 2 \times 10^3$	7	~ 1	p.r.	D.k. in N_2O -satd. 1 M SCN^- soln.	78-1093
9.19	Histidine	$< 1 \times 10^6$	7.0	0.1	p.r.	D.k. in N_2O -satd. SCN^- soln.	72-0036
9.20	Hydroquinone	6×10^7	—	—	p.r.	—	74-1168
9.21	<i>p</i> -Hydroxybenzoate ion	2.65×10^8	12.5	—	p.r.	—	74-3052
9.22	Lactate dehydrogenase	1.5×10^9	7.2	0.005	p.r.	D.k. in N_2O -satd. SCN^- soln.	77-1132
9.23	Lysozyme	6.6×10^8	—	0.1	p.r.	D.k. in N_2O -satd. SCN^- soln.	69-3039, 70-0967
9.24	Methionine	$(2 \pm 0.2) \times 10^6$	7.0	0.1	p.r.	D.k. in N_2O -satd. SCN^- soln.; k increases with pH.	72-0036
9.25	<i>p</i> -Methylphenoxide ion	5.52×10^3	12.5	—	p.r.	At pH 7 $k \approx 10^7$.	74-3052
9.26	Nicotinamide-adenine dinucleotide, reduced $(\text{SCN})_2^- + \text{NADH} \rightarrow 2\text{SCN}^- + \text{NADH}^+$	4.7×10^8	—	0.1	p.r.	D.k. in N_2O -satd. SCN^- soln.	71-0158
9.27	<i>p</i> -Nitrophenoxide ion	3.9×10^7	12.5	—	p.r.	—	74-3052
9.28	Papain	9.5×10^9 1.75×10^9	7 11.5	—	p.r.	D.k.; activated enzyme used.	74-1026
9.29	Phenol	$< 1 \times 10^7$	—	—	p.r.	—	74-1168
9.30	Phenoxide ion	3×10^8 3.42×10^8	12 12.5	—	p.r.	—	74-1168 74-3052
9.31	Phenylalanine	$< 1 \times 10^6$	7.0	0.1	p.r.	D.k. in N_2O -satd. SCN^- soln.	72-0036
9.32	Promethazine	2×10^9	—	—	p.r.	—	74-1168
9.33	Ribonuclease	3.4×10^7	7	0.05	p.r.	D.k. in N_2O -satd. SCN^- soln.; k increases at pH > 9.5 .	72-0037
9.34	Serum albumin	1×10^9 (bovine) 5×10^8 (human)	6.0 6.5	0.01	p.r.	D.k. in SCN^- soln.; also detd. reactivity with alkyl sulfate complexes.	76-1185
9.35	Subtilisin (serine protease)	1×10^8	7	—	p.r.	D.k. in N_2O -satd. 0.04 M SCN^- soln.; k increases with pH in alk. soln.	73-1147, 74-1119
9.36	Tetramethyl- piperidone- <i>N</i> -oxyl (TAN)	1.1×10^9 1.0×10^9 1.1×10^9	2 5-6 12	0.1	p.r.	D.k. in 0.1 M CNS^- soln.	71-0618
9.37	Thymine	$< 1 \times 10^6$ 3×10^7	6 12	—	p.r.	—	74-1168
9.38	Trypsin	$(5.1 \pm 0.5) \times 10^8$ $(3.06 \pm 0.3) \times 10^9$	7-8 11.5	0.04	p.r.	D.k. in N_2O -satd. SCN^- soln.	73-1067
9.39	Trypsinogen	$(2.6 \pm 0.3) \times 10^8$ $(3.6 \pm 0.4) \times 10^9$	7-8 12.1	—	p.r.	—	73-1067
9.40	Tryptophan $(\text{SCN})_2^- + \text{TrpH} \rightarrow 2\text{SCN}^- + \text{Trp} \cdot + \text{H}^+$ (pK of $\text{TrpH}^+ = 4.3$, ref. 76-1151)	3×10^8 $(2.7 \pm 0.3) \times 10^8$ 4.6×10^8	7 7.0 11.2	— 0.1 —	p.r. p.r. p.r.	— D.k. in N_2O -satd. SCN^- soln.; k increases with pH. —	69-3039 72-0036 73-1147
9.41	Tyrosine	$(5 \pm 0.5) \times 10^6$ 3.2×10^8	7.0 11.2	0.1 —	p.r. p.r.	D.k. in N_2O -satd. SCN^- soln.; k increases with pH. —	72-0036 73-1147

Formula Index

- AsO_2^- Arsenite ion, 2.2
 Br Bromine atom, 7.2
 Br^- Bromide ion, 1.2a, 2.3, 5.4–5.5
 $\text{BrCoH}_{15}\text{N}_5^{2+}$ Bromopentaamminecobalt(III) ion, 1.7e
 BrO^- Hypobromite ion, 1.3, 7.5
 BrO_2^- Bromite ion, 1.4, 7.6
 BrO_3^- Bromate ion, 7.7
 Br_2 Bromine molecule anion, table 7
 CCl_4 Carbon tetrachloride, 1.36
 $\text{CCoH}_{12}\text{N}_4\text{O}_3^+$ Carbonatopentaamminecobalt(III) ion, 1.6
 CHO_2^- Formate ion, 1.51, 2.67, 5.51, 6.63, 7.43, 9.18
 CHO_3^- Bicarbonate ion, 2.4
 CH_2NO_2^- *aci*-Nitromethane, 1.77
 CH_2O_2 Formic acid, 2.68, 4.12, 6.64
 CH_3NO_2 Nitromethane, 1.76
 $\text{CH}_4\text{N}_2\text{O}$ Urea, 1.100
 CH_4O Methanol, 1.69, 2.78, 4.14, 5.62, 6.77
 CN^- Cyanide ion, 2.5
 CNO^- Cyanate ion, 2.6
 CNS^- Thiocyanate ion, 1.13d, 2.7, 6.3, 7.24
 CO_3^- Carbonate radical ion, table 1
 $\text{C}_2\text{Cl}_3\text{O}_2^-$ Trichloroacetate ion, 1.91
 $\text{C}_2\text{H}_2\text{ClO}_2^-$ Chloroacetate ion, 1.37
 $\text{C}_2\text{H}_3\text{N}$ Acetonitrile, 1.18
 $\text{C}_2\text{H}_3\text{O}_2^-$ Acetate ion, 1.16, 2.32, 5.31
 $\text{C}_2\text{H}_4\text{O}_2$ Acetic acid, 2.33, 4.8, 5.32, 6.32
 $\text{C}_2\text{H}_5\text{NO}_2$ Glycine, 1.55, 2.70, 5.54, 6.68
 $\text{C}_2\text{H}_6\text{O}$ Ethanol, 1.49, 2.65, 3.5, 3.11, 4.10, 5.50, 6.62
 $\text{C}_2\text{H}_6\text{O}_2$ Ethyleneglycol, 4.11
 $\text{C}_2\text{H}_6\text{S}_2$ Dimethyl disulfide, 1.45, 2.64, 7.41
 $\text{C}_2\text{H}_7\text{NS}$ Cysteamine, 7.36, 9.14
 $\text{C}_2\text{H}_{18}\text{CoN}_5\text{O}_2^{2+}$ Acetatopentaamminecobalt(III) ion, 1.7h
 $\text{C}_2\text{N}_2\text{S}_2^-$ Bisthiocyanate radical ion, table 9
 $\text{C}_3\text{H}_2\text{O}_4^{2-}$ Malonate ion, 2.75
 $\text{C}_3\text{H}_3\text{N}$ Acrylonitrile, 2.38, 5.36, 6.37
 $\text{C}_3\text{H}_3\text{O}_2^-$ Acrylate ion, 2.37, 6.35
 $\text{C}_3\text{H}_4\text{N}_2$ Imidazole, 1.64
 $\text{C}_3\text{H}_4\text{O}_2$ Acrylic acid, 5.35, 6.36
 $\text{C}_3\text{H}_4\text{O}_4$ Malonic acid, 5.58
 $\text{C}_3\text{H}_5\text{NO}$ Acrylamide, 2.36, 5.34
 $\text{C}_3\text{H}_5\text{O}_2^-$ Propionate ion, 2.94
 $\text{C}_3\text{H}_5\text{O}_2\text{S}^-$ 3-Mercaptopropionate ion, 1.68
 $\text{C}_3\text{H}_6\text{O}$ Acetone, 1.17, 5.33, 6.33; Allyl alcohol, 2.40, 5.38, 6.40
 $\text{C}_3\text{H}_6\text{O}_2$ Propionic acid, 5.66, 6.88
 $\text{C}_3\text{H}_7\text{NO}_2$ Alanine, 1.24, 2.39, 5.37, 6.39
 $\text{C}_3\text{H}_7\text{NO}_2\text{S}$ Cysteine, 1.40, 6.54, 7.37, 8.18, 9.15
 $\text{C}_3\text{H}_7\text{NO}_3$ Serine, 2.99, 6.90
 $\text{C}_3\text{H}_8\text{O}$ 1-Propanol, 1.85, 2.92; 2-Propanol, 1.86, 2.93, 3.6, 4.15, 5.65, 6.87
 $\text{C}_3\text{H}_8\text{O}_3$ Glycerol, 4.13
 $\text{C}_4\text{H}_2\text{O}_4^{2-}$ Fumarate ion, 2.69, 6.65; Maleate ion, 6.76
 $\text{C}_4\text{H}_3\text{ClN}_2\text{O}_2$ 5-Chlorouracil, 6.51
 $\text{C}_4\text{H}_4\text{N}_2\text{O}_2$ Uracil, 1.99, 5.73, 6.100, 7.66
 $\text{C}_4\text{H}_4\text{O}_4$ Fumaric acid, 5.52, 6.66; Maleic acid, 5.57
 $\text{C}_4\text{H}_4\text{O}_4^{2-}$ Succinate ion, 2.100
 $\text{C}_4\text{H}_5\text{N}$ Allyl cyanide, 2.41, 5.39; Methacrylonitrile, 2.77, 5.61
 $\text{C}_4\text{H}_5\text{N}_3\text{O}$ Cytosine, 6.56
 $\text{C}_4\text{H}_5\text{O}_2^-$ Crotonate ion, 5.45; Methacrylate ion, 5.59
 $\text{C}_4\text{H}_6\text{NO}_4^-$ Aspartate ion, 1.32
 $\text{C}_4\text{H}_6\text{N}_2\text{O}_2$ Dihydrouracil, 5.49
 $\text{C}_4\text{H}_6\text{O}_2$ Crotonic acid, 2.51, 5.46; Methacrylic acid, 2.76, 5.60; Vinyl acetate, 2.116
 $\text{C}_4\text{H}_6\text{O}_4$ Succinic acid, 5.68, 6.92
 $\text{C}_4\text{H}_7\text{NO}_3$ *N*-Acetyl glycine, 1.21
 $\text{C}_4\text{H}_8\text{N}_2\text{O}_3$ Glycylglycine, 1.56
 $\text{C}_4\text{H}_8\text{N}_2\text{O}_4\text{Pt}$ *cis*-Bis(glycinato)platinum(II), 1.11; *trans*-Bis(glycinato)platinum(II), 1.12
 $\text{C}_4\text{H}_8\text{O}$ Tetrahydrofuran, 2.101a
 $\text{C}_4\text{H}_8\text{O}_2$ Dioxane, 2.64a
 $\text{C}_4\text{H}_9\text{NO}_2\text{S}$ Cysteine methyl ester (Methyl 2-amino-3-mercaptopropionate), 1.41; *S*-Methylcysteine, 1.72
 $\text{C}_4\text{H}_{10}\text{O}$ 2-Methyl-2-propanol (*tert*-Butanol), 1.74, 2.85, 5.64, 6.81
 $\text{C}_4\text{H}_{10}\text{O}_2\text{S}_2$ Dithiothreitol (*threo*-2,3-Dihydroxy-1,4-dithiobutane), 1.47, 6.60, 8.19, 9.17
 $\text{C}_4\text{H}_{10}\text{S}_2$ Diethyl disulfide, 1.43, 2.55, 7.39
 $\text{C}_4\text{H}_{12}\text{N}_4\text{Pt}^{2+}$ Bisethylenediamineplatinum(II) ion, 6.25
 $\text{C}_4\text{H}_{13}\text{ClN}_3\text{Pt}^+$ Chlorodiethylenetriamineplatinum(II) ion, 6.26
 $\text{C}_5\text{H}_5\text{N}$ Pyridine, 2.95
 $\text{C}_5\text{H}_5\text{N}_5$ Adenine, 6.38
 $\text{C}_5\text{H}_5\text{N}_5\text{O}$ Guanine, 6.69
 $\text{C}_5\text{H}_6\text{N}^+$ Pyridinium ion, 2.96
 $\text{C}_5\text{H}_6\text{N}_2\text{O}_2$ Thymine, 1.89, 5.71, 6.96, 7.62, 9.37
 $\text{C}_5\text{H}_8\text{O}_2$ Methyl methacrylate, 2.84, 5.63
 $\text{C}_5\text{H}_9\text{NO}_3\text{S}$ *N*-Acetylcysteine, 1.20
 $\text{C}_5\text{H}_9\text{NO}_4$ Glutamic acid, 6.67
 $\text{C}_5\text{H}_9\text{O}_5$ Deoxyribose, 5.48
 $\text{C}_5\text{H}_{10}\text{O}_2$ Ribose, 5.67
 $\text{C}_5\text{H}_{11}\text{NO}_2\text{S}$ Methionine, 1.70, 2.79, 6.78, 7.49, 8.22, 9.24; Penicillamine (2-amino-3-methyl-3-thiobutyric acid), 1.80
 $\text{C}_5\text{H}_{12}\text{O}$ 3-Pentanol, 2.87
 $\text{C}_6\text{FeN}_6^{4+}$ Hexacyanoferrate(II) ion, 1.9, 4.17
 $\text{C}_6\text{H}_4\text{BrO}^-$ *p*-Bromophenoxide ion, 1.35, 7.31, 8.16, 9.9
 $\text{C}_6\text{H}_4\text{ClO}^-$ *p*-Chlorophenoxide ion, 1.38, 7.33, 8.17, 9.11
 $\text{C}_6\text{H}_4\text{NO}_3^-$ *p*-Nitrophenoxide ion, 1.78, 9.27
 $\text{C}_6\text{H}_4\text{O}_4^{2-}$ Muconate ion (2,4-Hexadienedioate ion), 6.82
 $\text{C}_6\text{H}_5\text{O}^-$ Phenoxide ion, 1.82, 5.64a, 7.54, 8.25, 9.30
 $\text{C}_6\text{H}_5\text{O}_3\text{S}^-$ Benzenesulfonate ion, 6.46
 C_6H_6 Benzene, 1.33, 2.44
 $\text{C}_6\text{H}_6\text{O}$ Phenol, 1.81, 6.83, 7.53, 9.29
 $\text{C}_6\text{H}_6\text{O}_2$ Hydroquinone, 6.73, 7.46, 9.20
 $\text{C}_6\text{H}_6\text{O}_4^{2-}$ Hexenedioate ion, 6.71
 $\text{C}_6\text{H}_7\text{N}$ Aniline, 1.25
 $\text{C}_6\text{H}_7\text{O}_2^-$ Sorbate ion (2,4-Hexadienoate ion), 6.91
 $\text{C}_6\text{H}_7\text{O}_6^-$ Ascorbate ion, 1.31, 7.30, 8.14, 9.8
 $\text{C}_6\text{H}_8\text{N}^+$ Anilinium ion, 6.43
 $\text{C}_6\text{H}_8\text{O}_4\text{S}_2^{2-}$ 3,3'-Dithiobis(propionate ion), 1.46
 $\text{C}_6\text{H}_8\text{O}_6$ Ascorbic acid, 6.45, 7.30, 8.15, 9.8
 $\text{C}_6\text{H}_9\text{N}_2\text{O}_4$ *N*-Acetylglycylglycine, 1.22
 $\text{C}_6\text{H}_9\text{N}_3\text{O}_2$ Histidine, 1.62, 2.71, 6.72, 7.45, 8.20, 9.19
 C_6H_{10} Cyclohexene, 2.54
 $\text{C}_6\text{H}_{11}\text{N}_3\text{O}_4$ Glycylglycylglycine, 1.57
 $\text{C}_6\text{H}_{12}\text{O}_6$ Glucose, 1.52, 5.53
 $\text{C}_6\text{H}_{14}\text{N}_4\text{O}_2$ Arginine, 1.30
 $\text{C}_6\text{H}_{24}\text{CoN}_6^{3+}$ Trisethylenediaminecobalt(III) ion, 1.7b
 $\text{C}_7\text{H}_4\text{BrO}_2^-$ *o*-Bromobenzoate ion, 2.48; *p*-Bromobenzoate ion, 2.49, 6.49
 $\text{C}_7\text{H}_4\text{ClO}_2^-$ *p*-Chlorobenzoate ion, 2.50, 5.43, 6.50
 $\text{C}_7\text{H}_4\text{NO}^-$ *p*-Cyanophenoxide ion, 5.47a, 7.35, 9.13
 $\text{C}_7\text{H}_5\text{ClO}_2$ *p*-Chlorobenzoic acid, 5.44

- C_7H_5N Benzonitrile, 2.46, 6.48
 C_7H_5NO *p*-Cyanophenol, 6.53
 $C_7H_5O_2^-$ Benzoate ion, 2.45, 5.41, 6.47
 $C_7H_5O_3^-$ *p*-Hydroxybenzoate ion, 1.63, 2.73, 5.55, 6.74, 7.47, 9.21
 $C_7H_6NO_2^-$ *p*-Aminobenzoate ion, 6.41
 $C_7H_6O_2$ Benzoic acid, 5.42
 $C_7H_6O_3$ *p*-Hydroxybenzoic acid, 6.75; Salicylic acid, 6.89
 C_7H_7NO Benzamide, 2.43
 $C_7H_7NO_2$ *p*-Aminobenzoic acid, 5.40, 6.42
 $C_7H_7O^-$ *p*-Methylphenoxide ion, 1.73, 5.62a, 7.50, 8.23, 9.25
 $C_7H_7O_2^-$ *p*-Methoxyphenoxide ion, 1.71
 C_7H_8 Toluene, 1.90
 C_7H_8O Anisole, 1.25a, 2.42; *p*-Methylphenol, 6.80
 C_7H_9N *N*-Methylaniline, 1.71b
 $C_7H_{13}O$ Cycloheptanol, 2.53
 $C_7H_{20}CoN_5O_2^{2+}$ Benzoatopentaamminecobalt(III) ion, 1.7g
 $C_8H_7NO_2^-$ *p*-Cyanobenzoate ion, 2.52, 5.47, 6.52
 $C_8H_4O_3^{2-}$ Terephthalate ion, 2.101, 5.69, 6.93
 $C_8H_6O_3$ Terephthalic acid, 5.70
 C_8H_7N Indole, 1.65
 $C_8H_7O_2^-$ *o*-Toluate ion, 2.102; *m*-Toluate ion, 2.103; *p*-Toluate ion, 2.104, 6.97
 $C_8H_7O_3^-$ 2-Methoxybenzoate ion, 2.80; 3-Methoxybenzoate ion, 2.81; 4-Methoxybenzoate ion, 2.82, 6.79
 C_8H_8O Acetophenone, 1.19, 2.34
 $C_8H_8O_2$ Toluic acid, 5.72
 C_8H_9NO Acetanilide, 1.15, 2.31, 6.31
 $C_8H_{10}N_2O$ *p*-Nitroso-*N,N*-dimethylaniline, 2.86
 $C_9H_{10}O$ Benzyl methyl ether, 2.47; Ethoxybenzene, 1.50; 1-Phenylethanol, 2.89
 $C_8H_{10}O_2$ 1,2-Dimethoxybenzene, 2.56; 1,3-Dimethoxybenzene, 2.57; 1,4-Dimethoxybenzene, 2.58
 $C_8H_{11}N$ *N,N*-Dimethylaniline, 1.44
 $C_8H_{12}NO_2$ Norpseudopelletierine-*N*-oxyl (9-Azabicyclo[3,3,1]nonan-3-one-9-oxyl), 1.79
 $C_8H_{12}N_4O_3$ Glycylhistidine, 1.59
 $C_9H_4O_4^{2-}$ 2,3-Dimethoxybenzoate ion, 2.59; 2,4-Dimethoxybenzoate ion, 2.60; 3,4-Dimethoxybenzoate ion, 2.61; 2,6-Dimethoxybenzoate ion, 2.62; 3,5-Dimethoxybenzoate ion, 2.63
 $C_9H_3O_6^{3-}$ Trimesate ion, 2.105
 $C_9H_6O_2^{2-}$ Homophthalate ion, 2.72
 $C_9H_7O_3^-$ *p*-Acetylbenzoate ion, 2.35
 $C_9H_9I_2NO_3$ 3,5-Diiodotyrosine, 7.40
 $C_9H_{10}FNO_3$ *m*-Fluorotyrosine, 7.42
 $C_9H_{11}NO_2$ Phenylalanine, 1.83, 6.85, 7.55, 8.26, 9.31
 $C_9H_{11}NO_3$ Tyrosine, 1.98, 2.115, 6.99, 7.65, 8.30, 9.41
 $C_9H_{11}NO_3$ 3,4-Dihydroxyphenylalanine, 9.16
 $C_9H_{12}O$ 1-Phenyl-2-propanol, 2.90; 2-Phenyl-2-propanol, 2.91
 $C_9H_{12}O_3$ 1,2,3-Trimethoxybenzene, 2.106; 1,2,4-Trimethoxybenzene, 2.107; 1,3,5-Trimethoxybenzene, 2.108
 $C_9H_{13}N_3O_5$ Cytidine, 6.55
 $C_9H_{14}N^+$ Trimethylanilinium ion, 2.113
 $C_9H_{14}N_3O_5P$ Deoxycytidylic acid, 6.58
 $C_9H_{16}NO_2$ 2,2,6,6-Tetramethyl-4-oxo-1-piperidinylloxyl (TAN), 1.88, 6.94, 7.61, 8.27, 9.36
 $C_{10}H_2O_8^{4-}$ Pyromellitate ion, 2.97
 $C_{10}H_{10}Fe^+$ Ferricinium ion, 7.16
 $C_{10}H_{11}O_5^-$ 2,3,4-Trimethoxybenzoate ion, 2.109; 3,4,5-Trimethoxybenzoate ion, 2.110; 2,4,5-Trimethoxybenzoate ion, 2.111; 2,4,6-Trimethoxybenzoate ion, 2.112
 $C_{10}H_{12}N_2$ Tryptamine, 1.93
 $C_{10}H_{12}N_2O_8^{4-}$ Ethylenediaminetetraacetate ion, 1.50a
 $C_{10}H_{12}O_2$ Duroquinone, 1.48
 $C_{10}H_{14}N_5O_6P$ Deoxyadenylic acid, 6.57
 $C_{10}H_{14}N_5O_7P$ Deoxyguanylic acid, 6.59
 $C_{10}H_{14}N_5O_8P$ Guanylic acid, 7.44
 $C_{10}H_{14}O$ 1-(*p*-Ethylphenyl)ethanol, 2.66; 1-Phenyl-3-butanol, 2.88
 $C_{10}H_{16}N_2O_9P$ Thymidylic acid, 6.95
 $C_{10}H_{17}N_3O_6S$ Glutathione, 1.53
 $C_{11}H_{10}NO_2^-$ Indole-3-propionate ion, 1.66
 $C_{11}H_{12}N_2O_2$ Tryptophan, 1.94, 2.114, 3.16, 6.98, 7.64, 8.29, 9.40
 $C_{11}H_{13}N_3O$ Tryptophanamide, 1.96
 $C_{11}H_{14}N_2O_3$ Phenylalanylglycine, 1.84
 $C_{11}H_{14}N_2O_4$ Glycyltyrosine, 1.61
 $C_{11}H_{16}O$ 1-Methoxy-2-methyl-1-phenylpropane, 2.83
 $C_{12}H_{14}N_2O_2$ *N*-Methyltryptophan, 1.75; Tryptophan methyl ester (Methyl 2-amino-3-indolylpropionate), 1.95
 $C_{12}H_{25}NaO_4S$ Dodecyl sodium sulfate, 6.61
 $C_{12}H_{29}ClN_3Pt^+$ Chlorotetraethyldiethylenetriamineplatinum(II) ion, 6.27
 $C_{13}H_9O_3^-$ *p*-Phenoxybenzoate ion, 6.84
 $C_{13}H_{10}O$ Benzophenone, 1.34
 $C_{13}H_{14}N_2O_3$ *N*-Acetyltryptophan, 1.23
 $C_{13}H_{15}N_3O_3$ Glycyltryptophan, 1.60; Tryptophanylglycine, 1.97
 $C_{14}H_6O_8S_2^{3-}$ Anthrasemiquinone-2,6-disulfonate radical ion, 1.28, 4.9, 6.44; Anthrasemiquinone-2,7-disulfonate radical ion, 1.29
 $C_{14}H_7O_5S^{2-}$ Anthrasemiquinone-1-sulfonate radical ion, 1.26; Anthrasemiquinone-2-sulfonate radical ion, 1.27
 $C_{14}H_{14}ClN_3$ Acriflavin, 6.34, 7.28, 8.11
 $C_{15}H_{18}N_3O_4$ Glycylglycyltryptophan, 1.58
 $C_{16}H_{10}$ Pyrene (triplet state), 7.57
 $C_{16}H_{28}CoN_4^{2+}$ 5,7,7,12,14,14-Hexamethyl-1,4,8,11-tetraazacyclotetradeca-1,4,8,11-tetraene cobalt(II) ion, 1.5a
 $C_{16}H_{28}Ni^{2+}$ 5,7,7,12,14,14-Hexamethyl-1,4,8,11-tetraazacyclotetradeca-1,4,8,11-tetraenenickel(II) ion, 6.21, 7.21
 $C_{16}H_{32}CoN_4^{2+}$ 5,7,7,12,14,14-Hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-dienecobalt(II) ion, 1.5b, 6.7, 7.11, 8.4
 $C_{16}H_{32}CuN_4^{2+}$ 5,7,7,12,14,14-Hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-dienecopper(II) ion, 6.11
 $C_{16}H_{32}Ni^{2+}$ 5,7,7,12,14,14-Hexamethyl-1,4,8,11-tetraazacyclotetradeca-4,11-dienenicke(II) ion, 6.20, 7.20
 $C_{16}H_{36}CuN_4^{2+}$ 5,7,7,12,14,14-Hexamethyl-1,4,8,11-tetraazacyclotetradecanecopper(II) ion, 6.10
 $C_{16}H_{36}Ni^{2+}$ 5,7,7,12,14,14-Hexamethyl-1,4,8,11-tetraazacyclotetradecanenickel(II) ion, 6.19, 7.19, 9.3
 $C_{17}H_{20}N_2S$ Promethazine, 7.56, 9.32

$C_{18}H_{16}N_2O_4S_2$ Cystine dimethyl ester (Dimethyl 3,3'-dithiobis(2-aminopropionate)), 1.42
 $C_{19}H_{42}ClN$ Hexadecyltrimethylammonium chloride, 6.70
 $C_{20}H_{32}N_6O_{12}S_2$ Glutathione disulfide, 1.54
 $C_{21}H_{29}N_7O_{10}P_2$ Nicotinamide-adenine dinucleotide, reduced, 7.51, 8.24, 9.26
 $C_{30}H_{24}N_6Ru^{2+}$ Tris(2,2'-bipyridine)ruthenium(II) ion, 1.13, 7.23
 Ce^{3+} Cerium(III) ion, 2.8, 4.3, 6.4
 Cl^- Chloride ion, 2.9, 4.4, 5.6–5.7
 $ClCoH_{15}N_5^{2+}$ Chloropentaamminecobalt(III) ion, 1.7d
 $ClCrH_{15}N_5^{2+}$ Chloropentaamminechromium(III) ion, 1.8a
 $ClH_{15}IrN_5^{2+}$ Chloropentaammineiridium(III) ion, 1.10a
 $ClH_{15}N_5Rh^{2+}$ Chloropentaamminerhodium(III) ion, 1.12a
 $ClH_{15}N_5Ru^{2+}$ Chloropentaammineruthenium(III) ion, 1.13b
 ClO_2 Chlorine dioxide, 6.5, 7.8
 Cl_2^- Chlorine molecule anion, table 6
 Cl_4Pt^{2-} Tetrachloroplatinate(II) ion, 6.24, 7.22
 Co^+ Cobalt(I) ion, 7.9
 Co^{2+} Cobalt(II) ion, 1.5, 6.6, 7.10, 8.3
 $CoH_{15}IN_5^{2+}$ Iodopentaamminecobalt(III) ion, 8.5
 $CoH_{15}N_5O_4S^+$ Sulfatopentaamminecobalt(III) ion, 1.7j
 $CoH_{15}N_5O_4S^+$ Sulfatopentaamminecobalt(III) ion, 1.7f
 $CoH_{15}N_6O_2^{2+}$ Nitritopentaamminecobalt(III) ion, 1.7i
 $CoH_{16}N_4O_2^{3+}$ Diaquotetraamminecobalt(III) ion, 1.7
 $CoH_{16}N_5O_4P^+$ Phosphatopentaamminecobalt(III) ion, 1.7c
 $CoH_{16}N_6^{3+}$ Hexaamminecobalt(III) ion, 1.7a
 Cr^{2+} Chromium(II) ion, 2.10, 6.8, 7.12, 8.6
 $CrH_{17}N_5O_3^{3+}$ Aquopentaamminechromium(III) ion, 1.8b
 Cu^{2+} 1.8c, 6.9
 Fe^{2+} Iron(II) ion, 2.11, 4.5, 6.12, 7.15, 8.7
 H Hydrogen atom, 7.3
 HIO^- 8.9
 HNO_3^- 4.21
 HO^- Hydroxide ion, 2.19, 5.16, 6.22
 HO_2 Hydroperoxy radical, 6.2, 7.4
 HO_3P^{2-} 2.23, 5.19–5.20
 HO_3S^- Hydrogen sulfite ion, 5.28
 HO_3Se 3.14
 HO_4P^- Phosphate radical ion, table 5
 HO_4P^{2-} 2.26
 HO_5P^{2-} 5.25
 HO_5S^- Hydrogen peroxysulfate ion, 2.28
 H_2O Water, 2.20, 4.19, 4.20, 4.24, 4.25
 H_2O_2 Hydrogen peroxide, 1.9a, 2.21, 5.17–5.18, 6.23, 7.17
 $H_2O_2P^-$ Hypophosphite ion, 2.22, 5.22–5.24
 $H_2O_3P^-$ Phosphite ion, 2.24, 5.21
 H_2O_3Se 3.15
 $H_2O_4P^-$ Phosphate radical, table 5
 $H_2O_4P^-$ Phosphate ion, 2.25, 4.23
 H_3NO Hydroxylamine, 2.12, 5.13, 6.15
 H_4NO^+ Hydroxylammonium ion, 2.13, 5.14, 6.16
 H_4N_2 Hydrazine, 2.14, 5.09
 $H_5N_2^+$ Hydrazinium ion, 2.15, 5.10–5.11, 6.14
 $H_{17}N_5ORh^{3+}$ Aquopentaamminerhodium(III) ion, 1.12b
 $H_{17}N_5ORu^{3+}$ Aquopentaammineruthenium(III) ion, 1.13c
 $H_{18}N_6Ru^{3+}$ Hexaammineruthenium(III) ion, 1.13a
 I^- Iodide ion, 1.10, 5.8, 9.2
 IO^- Hypiodite ion, 8.8
 I_2^- Iodine molecule anion, table 8
 Mn^{2+} 1.10b, 6.13, 7.18
 NO Nitric oxide, 4.18
 NO_2 Nitrogen dioxide, 1.10d; table 4
 NO_2^- Nitrite ion, 1.10c, 2.17, 4.6, 5.15, 6.18
 NO_2^{2-} Nitrite radical anion, 4.25
 NO_3 Nitrate radical, table 4
 NO_3^- Nitrate ion, 2.18
 NO_3^{2-} Nitrate radical anion, table 4
 N_2O Nitrous oxide, 3.3
 N_2O_3 Dinitrogen trioxide, 4.24
 N_2O_4 Dinitrogen tetroxide, table 4
 N_3^- Azide ion, 2.16, 5.12, 6.17
 Ni^{2+} Nickel(II) ion, 1.10c
 O_2 Oxygen, 3.4, 4.22
 O_2^- Superoxide ion, 1.2, 6.2
 O_2S^- Hyposulfite radical ion, table 3
 O_2Se^- 3.12
 $O_2S_2^-$ Thiosulfite radical ion, table 3
 O_2U^+ Uranyl(IV) ion, 7.26, 9.4
 O_2U^{2+} Uranyl(VI) ion, 1.14
 O_3S^- Sulfite radical ion, 1.13f; table 3
 O_3S^{2-} Sulfite ion, 1.13e, 2.27, 5.26–5.27, 6.28
 $O_3S_2^-$ Thiosulfate radical ion, table 3
 $O_3S_2^{2-}$ Thiosulfate ion, 3.8, 5.29
 O_3Se^- 3.13
 O_3Si^{2-} Silicate ion, 2.29
 O_4P^{2-} Phosphate radical ion, table 5
 O_4S^- Sulfate radical ion, table 2
 O_5S^- Peroxysulfate radical ion, table 3
 $O_5S_2^{2-}$ Peroxythiosulfate ion, 5.30
 $Ti(III)$ Titanium(III) ion, 6.28a
 Tl^+ Thallium(I) ion, 2.30, 4.7, 6.29, 7.25
 V^{2+} Vanadium(II) ion, 6.30, 7.27, 8.10
 Zn^{2+} Zinc(II) ion, 1.14a

References

- 57-7007 Grossweiner, L.I.; Matheson, M.S., The kinetics of the dihalide ions from the flash photolysis of aqueous alkali halide solutions, *J. Phys. Chem.* 61: 1089-95 (1957).
- 64-7009 Martin, T.W.; Rummel, R.E.; Gross, R.C., Electron exchange kinetics of the NO_3 free radical in solution, *J. Am. Chem. Soc.* 86: 2595-600 (1964).
- 65-0382 Sutton, H.C.; Adams, G.E.; Boag, J.W.; Michael, B.D., Radical yields and kinetics in the pulse radiolysis of potassium bromide solutions, *Pulse Radiolysis*, M. Ebert, J.P. Keene, A.J. Swallow, J.H. Baxendale (eds.), Academic Press, New York, 1965, p.61-81.
- 65-0383 Cercek, B.; Ebert, M.; Gilbert, C.W.; Swallow, A.J., Pulse radiolysis of aerated aqueous potassium bromide solutions, *Ibid.*, p.83-98.
- 65-0386 Adams, G.E.; Boag, J.W.; Currant, J.; Michael, B.D., The pulse radiolysis of aqueous solutions of thiocyanate ion, *Ibid.*, p.117-29.
- 66-0001 Adams, G.E.; Boag, J.W.; Michael, B.D., Transient species produced in irradiated water and aqueous solutions containing oxygen, *Proc. Roy. Soc. (London) Ser. A* 289: 321-41 (1966).
- 66-0019 Heckel, E.; Henglein, A.; Beck, G., Pulsradiolytische Untersuchung des Radikalanions SO_4^- , *Ber. Bunsenges. Physik. Chem.* 70(2): 149-54 (1966).
- 66-0139 Weeks, J.L.; Rabani, J., The pulse radiolysis of deaerated aqueous carbonate solutions. I. Transient optical spectrum and mechanism. II. pK for OH radicals, *J. Phys. Chem.* 70(7): 2100-6 (1966).
- 66-0425 Matheson, M.S.; Mulac, W.A.; Weeks, J.L.; Rabani, J., The pulse radiolysis of deaerated aqueous bromide solutions, *J. Phys. Chem.* 70: 2092-9 (1966).
- 67-0002 Broszkiewicz, R.K., The radiation-induced formation of NO_3 in aqueous solutions, *Intern. J. Appl. Radiat. Isot.* 18: 25-32 (1967).
- 67-0032 Daniels, M.; Wigg, E.E., Radiation chemistry of the aqueous nitrate system. I. γ radiolysis of dilute solutions, *J. Phys. Chem.* 71(4): 1024-33 (1967).
- 67-7012 Hayon, E.; McGarvey, J.J., Flash photolysis in the vacuum ultraviolet region of SO_4^{2-} , CO_3^{2-} , and OH^- ions in aqueous solutions, *J. Phys. Chem.* 71(5): 1472-7 (1967).
- 67-7058 Dogliotti, L.; Hayon, E., Flash photolysis of persulfate ions in aqueous solutions. Study of the sulfate and ozonide radical anions, *J. Phys. Chem.* 71(8): 2511-6 (1967).
- 67-7171 Langmuir, M.E.; Hayon, E., Flash photolysis study of mercury(II) halide complexes in aqueous solution. Rates of reaction of X_2^- radical anions, *J. Phys. Chem.* 71(12): 3808-14 (1967).
- 67-7274 Dogliotti, L.; Hayon, E., Transient species produced in the photochemical decomposition of ceric salts in aqueous solution. Reactivity of NO_3 and HSO_4 free radicals, *J. Phys. Chem.* 71: 3802-8 (1967).
- 68-0153 Buxton, G.V.; Dainton, F.S., The radiolysis of aqueous solutions of oxybromine compounds. The spectra and reactions of BrO and BrO_2 , *Proc. Roy. Soc. (London) Pt. A* 304: 427-39 (1968).
- 68-0313 Ward, J.F.; Kuo, I., Steady state and pulse radiolysis of aqueous chloride solutions of nucleic acid components, *Advan. Chem. Ser.* 81: 368-73 (1968).
- 68-0375 Baxendale, J.H.; Bevan, P.L.T.; Stott, D.A., Pulse radiolysis of aqueous thiocyanate and iodide solutions, *Trans. Faraday Soc.* 64(9): 2389-97 (1968).
- 68-7024 Ottolenghi, M.; Rabani, J., Photochemical generation of nitrogen dioxide in aqueous solutions, *J. Phys. Chem.* 72(2): 593-8 (1968).
- 68-7072 Dogliotti, L.; Hayon, E., Flash photolysis study of sulfite, thiocyanate, and thiosulfate ions in solution, *J. Phys. Chem.* 72(5): 1800-7 (1968).
- 68-7281 Dogliotti, L.; Hayon, E., Optical spectrum of SO_3^- radicals produced from the photolysis of dithionate ions in solution, *Nature (London)* 218: 949-50 (1968).
- 68-9084 Pennington, D.E.; Haim, A., Stoichiometry and mechanism of the Cr(II) -peroxydisulfate reaction, *J. Am. Chem. Soc.* 90(14): 3700-4 (1968).
- 69-0158 Roebke, W.; Renz, M.; Henglein, A., Pulsradiolyse der Anionen $\text{S}_2\text{O}_8^{2-}$ und HSO_5^- in waessriger Loesung, *Int. J. Radiat. Phys. Chem.* 1(1): 39-44 (1969).
- 69-0180 Schoeneshoefer, M.; Henglein, A., Pulsradiolytische Untersuchung der Dissoziationsgleichgewichte des gemischten Radikal-Anion-Komplexes BrSCN^- , *Ber. Bunsenges. Physik. Chem.* 73(3): 289-93 (1969).
- 69-0417 Daniels, M., Radiation chemistry of the aqueous nitrate system. III. Pulse electron radiolysis of concentrated sodium nitrate solutions, *J. Phys. Chem.* 73(11): 3710-7 (1969).
- 69-0439 Graetzel, M.; Henglein, A.; Lilie, J.; Beck, G., Pulsradiolytische Untersuchung einiger Elementarprozesse der Oxydation und Reduktion des Nitrits, *Ber. Bunsenges. Physik. Chem.* 73(7): 646-53 (1969).
- 69-0565 Schoeneshoefer, M., Pulsradiolytische Untersuchung des gemischten Radikal-Anion-Komplexes ClSCN^- , *Int. J. Radiat. Phys. Chem.* 1: 505-13 (1969).
- 69-3039 Adams, G.E.; Willson, R.L.; Aldrich, J.E.; Cundall, R.B., On the mechanism of the radiation-induced inactivation of lysozyme in dilute aqueous solution, *Int. J. Radiat. Biol.* 16(4): 333-42 (1969).
- 69-7297 Blyumenfel'd, L.A.; Karyakin, A.V.; Kuz'min, V.A.; Fomin, G.V.; Chibisov, A.K.; Sholina, S.I., Photochemical reduction of organic electron acceptors in aqueous solutions, *Bull. Acad. Sci. USSR Div. Chem. Sci. (English Transl.)* (11): 2463-4 (1969), translated from *Izv. Akad. Nauk, Ser. Khim.* (11): 2615-7 (1969).
- 70-0018 Amichai, O.; Treinin, A., On the oxyiodine radicals in aqueous solution, *J. Phys. Chem.* 74(4): 830-5 (1970).
- 70-0151 Graetzel, M.; Henglein, A.; Taniguchi, S., Pulsradiolytische Beobachtungen ueber die Reduktion des NO_3^- -Ions und ueber Bildung und Zerfall der persalpetrigen Saeure in waessriger Loesung, *Ber. Bunsenges. Physik. Chem.* 74(3): 292-8 (1970).
- 70-0164 Schoeneshoefer, M.; Henglein, A., Pulsradiolytische Untersuchung der Komplexe ISCN^- und I_2SCN^- , *Ber. Bunsenges. Physik. Chem.* 74(4): 393-8 (1970).
- 70-0228 Graetzel, M.; Taniguchi, S.; Henglein, A., Pulsradiolytische Untersuchung der NO-Oxydation und des Gleichgewichts $\text{N}_2\text{O}_3 \rightleftharpoons \text{NO} + \text{NO}_2$ in waessriger Loesung, *Ber. Bunsenges. Physik. Chem.* 74(5): 488-92 (1970).
- 70-0241 Pruetz, W.A.; Land, E.J., Chemiluminescent reactions after pulse radiolysis of aqueous solutions of acriflavin. Effects of halides and pseudo halides, *J. Phys. Chem.* 74(10): 2107-14 (1970).
- 70-0247 Behar, D.; Czapski, G.; Duchovny, I., Carbonate radical in flash photolysis and pulse radiolysis of aqueous carbonate solutions, *J. Phys. Chem.* 74(10): 2206-10 (1970).

- 70-0254 Barker, G.C.; Fowles, P.; Stringer, B., Pulse radiolytic induced transient electrical conductance in liquid solutions. Part 2. Radiolysis of aqueous solutions of NO_3^- , NO_2^- and $\text{Fe}(\text{CN})_6^{3-}$, *Trans. Faraday Soc.* 66(6): 1509-19 (1970).
- 70-0302 Black, E.D.; Hayon, E., Pulse radiolysis of phosphate anions H_2PO_4^- , HPO_4^{2-} , PO_4^{3-} , and $\text{P}_2\text{O}_7^{4-}$ in aqueous solutions, *J. Phys. Chem.* 74(17): 3199-203 (1970).
- 70-0326 Nakashima, M.; Hayon, E., Rates of reaction of inorganic phosphate radicals in solution, *J. Phys. Chem.* 74(17): 3290-1 (1970).
- 70-0967 Adams, G.E.; Cundall, R.B.; Willson, R.L., Pulse radiolysis: an oxidative-reductive probe in general biochemical and enzyme studies, *Chemical Reactivity and Biological Role of Functional Groups in Enzymes*, R.M.S. Smellie (ed.), Academic Press, London, 1970, No. 31, p.171-2.
- 70-7234 Kraljic, I., Photolytic determination of SO_4^- rate constants, *Int. J. Radiat. Phys. Chem.* 2(2): 59-68 (1970).
- 70-7262 Kuz'min, V.A.; Chibisov, A.K., The OH radical in condensed systems, studied by using pulsed illumination, *High Energy Chem. (Engl. Transl.)* 4(2): 146-7 (1970), translated from *Khim. Vys. Energ.* 4(2): 171-2 (1970).
- 70-7264 Treinin, A.; Hayon, E., Absorption spectra and reaction kinetics of NO_2 , N_2O_3 , and N_2O_4 in aqueous solution, *J. Am. Chem. Soc.* 92(20): 5821-8 (1970).
- 70-7726 Thornton, A.T.; Laurence, G.S., The rates of oxidation of Fe^{2+} , Mn^{2+} , and Co^{+} by $\cdot\text{Br}_2^-$ radical ions, *Chem. Commun.* 443-4 (1970).
- 70-7727 Martin, T.W.; Glass, R.W., Competitive electron transfer. Activity-defined formation constants of cerium(III) nitrate complexes based on reaction with the nitrate free radical, *J. Am. Chem. Soc.* 92(17): 5075-83 (1970).
- 70-7728 Glass, R.W.; Martin, T.W., Flash generation and decay kinetics of the nitrate radical in aqueous nitric acid solutions, *J. Am. Chem. Soc.* 92(17): 5084-93 (1970).
- 71-0061 Fielden, E.M.; Roberts, P.B., Pulse radiolysis studies of the radiosensitizer norpseudopelletierine-N-oxyl (NPPN). I. Radiation chemistry, *Int. J. Radiat. Biol.* 20(4): 355-62 (1971).
- 71-0158 Land, E.J.; Swallow, A.J., One-electron reactions in biochemical systems as studied by pulse radiolysis. IV. Oxidation of dihydronicotinamide-adenine dinucleotide, *Biochim. Biophys. Acta* 234: 34-42 (1971).
- 71-0461 Zagorski, Z.P.; Sehested, K.; Nielsen, S.O., Pulse radiolysis of aqueous alkaline sulfite solutions, *J. Phys. Chem.* 75(23): 3510-7 (1971).
- 71-0618 Willson, R.L., Pulse radiolysis studies on reaction of triacetoneamine-N-oxyl with radiation-induced free radicals, *Trans. Faraday Soc.* 67(10): 3008-19 (1971).
- 71-7574 Kuz'min, V.A.; Chibisov, A.K., Photochemical oxidation of water by aromatic ketones, benzo-, and naphthoquinones, *Theor. Exper. Chem. (Engl. Transl.)* (10): 337-9 (1971), translated from *Teor. Eksp. Khim.* 7(3): 403-7 (1971).
- 72-0036 Adams, G.E.; Aldrich, J.E.; Bisby, R.H.; Cundall, R.B.; Redpath, J.L.; Willson, R.L., Selective free radical reactions with proteins and enzymes: Reactions of inorganic radical anions with amino acids, *Radiat. Res.* 49(2): 278-89 (1972).
- 72-0037 Adams, G.E.; Bisby, R.H.; Cundall, R.B.; Redpath, J.L.; Willson, R.L., Selective free radical reactions with proteins and enzymes: The inactivation of ribonuclease, *Radiat. Res.* 49(2): 290-9 (1972).
- 72-0094 Matthews, R.W.; Mahlman, H.A.; Sworski, T.J., Elementary processes in the radiolysis of aqueous sulfuric acid solutions. Determinations of both G_{OH} and $\text{G}_{\text{SO}_4^-}$, *J. Phys. Chem.* 76(9): 1265-72 (1972).
- 72-0266 Schoeneshoefer, M., Pulsradiolytische Untersuchung zur Oxidation der Ascorbinsäure durch OH-Radikale und Halogen-Radikal-Komplexe in wässriger Lösung, *Z. Naturforsch. Teil B* 27(6): 649-59 (1972).
- 72-0475 Zhestkova, T.P.; Pikaev, A.K.; Spitsyn, V.N., Pulse radiolysis of concentrated aqueous solutions of potassium thiocyanate, *Dokl. Phys. Chem. (Engl. Transl.)* 204(4-6): 482-5 (1972), transl. from *Dokl. Akad. Nauk SSSR* 204(5): 1147-50 (1972).
- 72-3107 Patterson, L.K.; Bansal, K.M.; Bogan, G.; Infante, G.A.; Fendler, E.J.; Fendler, J.H., Micellar effects on Cl_2^- reactivity. Reactions with surfactants and pyrimidines, *J. Am. Chem. Soc.* 94(26): 9028-32 (1972).
- 72-5049 Behar, D.; Fessenden, R.W., Electron spin resonance studies of inorganic radicals in irradiated aqueous solutions. I. Direct observation, *J. Phys. Chem.* 76(12): 1706-10 (1972).
- 72-5050 Behar, D.; Fessenden, R.W., Electron spin resonance studies of inorganic radicals in irradiated aqueous solutions. II. Radical trapping with nitromethane, *J. Phys. Chem.* 76(12): 1710-21 (1972).
- 72-7008 Hayon, E.; Treinin, A.; Wilf, J., Electronic spectra, photochemistry, and autoxidation mechanism of the sulfite-bisulfite-pyrosulfite systems. The SO_2^- , SO_3^- , SO_4^- , and SO_5^- radicals, *J. Am. Chem. Soc.* 94(1): 47-57 (1972).
- 72-7335 Kuzmin, V.A.; Chibisov, A.K.; Karyakin, A.V., Decay kinetics of the triplet and anthrasemiquinone radicals in aqueous organic media, *Int. J. Chem. Kinet.* 4(6): 639-44 (1972).
- 72-7383 Kuz'min, V.A., Reactions of the CO_3^- and SiO_3^- radical anions, *High Energy Chem. (Engl. Transl.)* 6(4): 338-9 (1972), transl. from *Khim. Vys. Energ.* 6(4): 381-2 (1972).
- 72-7464 Kuz'min, V.A.; Karyakin, A.B.; Chibisov, A.K., Direct and sensitized photoreduction of anthraquinone derivatives in aqueous and aqueous alcoholic solutions, *High Energy Chem. (Engl. Transl.)* 6(6): 442-5 (1972), transl. from *Khim. Vys. Energ.* 6(6): 502-6 (1972).
- 72-7506 Malone, S.D.; Endicott, J.F., The photochemical behavior of cobalt complexes containing macrocyclic (N_4) ligands. Oxidation-reduction chemistry of dihalogen radical anions, *J. Phys. Chem.* 76(16): 2223-9 (1972).
- 73-0030 Anbar, M.; Bamnolker, M.; Ross, A.B., Selected specific rates of reactions of transients from water. I. Hydrated electron, *NSRDS-NBS* 43, 1973, 67p.
- 73-1020 Redpath, J.L., Pulse radiolysis of dithiothreitol, *Radiat. Res.* 54(3): 364-74 (1973).
- 73-1027 Schoeneshoefer, M., Pulsradiolytische Untersuchung ueber das $\text{S}_4\text{O}_6^{3-}$ -Radikal als Zwischenprodukt der Thiosulfatoxidation und Tetrathionatreduktion, *Int. J. Radiat. Phys. Chem.* 5(4): 375-86 (1973).
- 73-1030 Lesigne, B.; Ferradini, C.; Pucheault, J., Pulse radiolysis study of the direct effect on sulfuric acid, *J. Phys. Chem.* 77(17): 2156-8 (1973).
- 73-1039 Jayson, G.G.; Parsons, B.J.; Swallow, A.J., Some simple, highly reactive, inorganic chlorine derivatives in aqueous solution. Their formation using pulses of radiation and their role in the mechanism of the Fricke dosimeter, *J. Chem. Soc. Faraday Trans. I* 69: 1597-607 (1973).
- 73-1050 Grabner, G.; Getoff, N.; Schwoerer, F., Pulsradiolyse von H_3PO_4 , H_2PO_4^- , HPO_4^{2-} und $\text{P}_2\text{O}_7^{4-}$ in wässriger Lösung. II. Spektren und Kinetik der Zwischenprodukte, *Int. J. Radiat. Phys. Chem.* 5(5): 405-17 (1973).
- 73-1057 Ellis, J.D.; Green, M.; Sykes, A.G.; Buxton, G.V.; Sellers, R.M., Pulse radiolysis of titanium(III) and other metal(III) ions in the presence of formic acid, *J. Chem. Soc. Dalton Trans.* (16): 1724-8 (1973).

- 73-1060 Roberts, P.B., A radiation chemical study of the inactivation and active site composition of carboxypeptidase A, *Int. J. Radiat. Biol.* 24(2): 143-52 (1973).
- 73-1065 Quintiliani, M.; Shejbal, J.; Davies, J.V.; Ebert, M.; Gilbert, C.W., Reactions of radiation-induced I^- , I_2^- and I_3^- with alcohol dehydrogenase and aldolase, *Int. J. Radiat. Biol.* 24(3): 243-55 (1973).
- 73-1067 Adams, G.E.; Redpath, J.L.; Bisby, R.H.; Cundall, R.B., Selective free radical reactions with proteins and enzymes. Reactions of inorganic radical anions with trypsin, *J. Chem. Soc. Faraday Trans. I* 69(9): 1608-17 (1973).
- 73-1147 Bisby, R.H.; Cundall, R.B.; Adams, G.E.; Redpath, J.L., Free radical reactions with subtilisin Carlsberg, *Fast Processes in Radiation Chemistry and Biology*, G.E. Adams, E.M. Fielden and B.D. Michael (eds.), Wiley, 1975, p.309-18 (Proc. of the Fifth L.H. Gray Conference, University of Sussex, Sept. 10-14, 1973).
- 73-1148 Roberts, P.B.; Fielden, E.M., Superoxide dismutase inactivation by Br_2^- radicals, *Ibid.*, p.319-26.
- 73-3006 Redpath, J.L.; Willson, R.L., Reducing compounds in radioprotection and radio-sensitization: Model experiments using ascorbic acid, *Int. J. Radiat. Biol.* 23(1): 51-65 (1973).
- 73-5022 Chawla, O.P.; Arthur, N.L.; Fessenden, R.W., An electron spin resonance study of the photolysis of aqueous sulfite solutions, *J. Phys. Chem.* 77(6): 772-6 (1973).
- 73-5403 Chawla, O.P., An electron spin resonance study of the mechanism and kinetics of photochemical reactions in aqueous solutions, Ph.D., Thesis, Carnegie-Mellon University, Pittsburgh, PA, 1973, 145p.
- 73-7043 Mialocq, J.C.; Barat, F.; Gilles, L.; Hickel, B.; Lesigne, B., Flash photolysis of chlorine dioxide in aqueous solution, *J. Phys. Chem.* 77(6): 742-9 (1973).
- 73-7066 Natarajan, P.; Endicott, J.F., Direct observation of the dibromide radical anion oxidation of tris(bipyridyl)ruthenium(II). Evidence for a triplet-to-triplet energy transfer mechanism in the photosensitized redox decomposition of cobalt(III) substrates, *J. Phys. Chem.* 77(7): 971-2 (1973).
- 73-7109 Chen, S.; Cope, V.W.; Hoffman, M.Z., Behavior of CO_3^- radicals generated in the flash photolysis of carbonatoamine complexes of cobalt(III) in aqueous solution, *J. Phys. Chem.* 77(9): 1111-6 (1973).
- 73-7159 Thornton, A.T.; Laurence, G.S., Kinetics of oxidation of transition-metal ions by halogen radical anions. I. The oxidation of iron(II) by dibromide and dichloride ions generated by flash photolysis, *J. Chem. Soc. Dalton Trans.* (8): 804-13 (1973).
- 73-7316 Thornton, A.T.; Laurence, G.S., Kinetics of oxidation of transition-metal ions by halogen radical anions. Part II. The oxidation of cobalt(II) by dichloride ions generated by flash photolysis, *J. Chem. Soc. Dalton Trans.* (16): 1632-6 (1973).
- 73-7317 Laurence, G.S.; Thornton, A.T., Kinetics of oxidation of transition-metal ions by halogen radical anions. Part III. The oxidation of manganese(II) by dibromide and dichloride ions generated by flash photolysis, *J. Chem. Soc. Dalton Trans.* (16): 1637-44 (1973).
- 73-7352 Chen, S.; Hoffman, M.Z., Rate constants for the reaction of the carbonate radical with compounds of biochemical interest in neutral aqueous solution, *Radiat. Res.* 56(1): 40-7 (1973).
- 73-7569 Kuz'min, V.A.; Chibisov, A.K., One-electron oxidation of inorganic anions by 2,6-disulfo-9,10-anthraquinone in the triplet state, *Dokl. Phys. Chem. (Engl. Transl.)* 212(4-6): 853-5 (1973), transl. from *Dokl. Akad. Nauk SSSR* 212(5): 1146-8 (1973).
- 74-1026 Adams, G.E.; Redpath, J.L., Selective free-radical reactions with proteins and enzymes: pulse radiolysis of inactivation studies on papain, *Int. J. Radiat. Biol.* 25(2): 129-38 (1974).
- 74-1033 Eriksen, T.E., pH Effects on the pulse radiolysis of deoxygenated aqueous solutions of sulfur dioxide, *J. Chem. Soc. Faraday Trans. I* 70(2): 208-15 (1974).
- 74-1038 Dodson, R.W.; Schwarz, H.A., Pulse radiolysis studies of chloride complexes of thallium(II). Absorption spectra and stability constants of $TlCl^+$, $TlCl_2^-$, and $TlCl_3^-$, *J. Phys. Chem.* 78(9): 892-9 (1974).
- 74-1087 Broszkiewicz, R.K., Pulse radiolysis studies on chloro-complexes of palladium, *Int. J. Radiat. Phys. Chem.* 6(4): 249-58 (1974).
- 74-1096 Baverstock, K.; Cundall, R.B.; Adams, G.E.; Redpath, J.L., Selective free radical reactions with proteins and enzymes: the inactivation of α -chymotrypsin, *Int. J. Radiat. Biol.* 26(1): 39-46 (1974).
- 74-1104 Laurence, G.S.; Thornton, A.T., Kinetics of oxidation of transition-metal ions by halogen radical anions, Part IV. The oxidation of vanadium(II) and chromium(II) by di-iodide, dibromide, and dichloride ions generated by pulse radiolysis, *J. Chem. Soc. Dalton Trans.* (11): 1142-8 (1974).
- 74-1119 Bisby, R.H.; Cundall, R.B.; Adams, G.E.; Redpath, J.L., Selective free radical reactions with proteins and enzymes. The inactivation of subtilisin Carlsberg and subtilisin Novo, *J. Chem. Soc., Faraday Trans. I* 70(12): 2210-8 (1974).
- 74-1125 Badiello, R.; Tamba, M.; Quintiliani, M., Primary and secondary radicals in the radiation-induced inactivation of yeast alcoholdehydrogenase, *Int. J. Radiat. Biol.* 26(4): 311-9 (1974).
- 74-1140 Pikaev, A.K.; Sibirskaya, G.K.; Shirshov, E.M.; Glazunov, P.Ya.; Spitsyn, V.I., Pulsed radiolysis of concentrated aqueous solutions of nitric acid, *Dokl. Phys. Chem. (Engl. Transl.)* 215(1-3): 328-31 (1974), transl. from *Dokl. Akad. Nauk SSSR* 215(3): 645-8 (1974).
- 74-1149 Zhestkova, T.P.; Pikaev, A.K., Destruction rate of Cl_2^- anion-radicals during pulse radiolysis of concentrated aqueous lithium chloride solutions, *Bull. Acad. Sci. USSR, Div. Chem. Sci. (Engl. Transl.)* 23(4): 877-8 (1974), transl. from *Izv. Akad. Nauk SSSR, Ser. Khim.* (4): 913-4 (1974).
- 74-1168 Willson, R.L., Electron-transfer reactions studied by pulse radiolysis, *Biochem. Soc. Trans.* 2(5): 1082-4 (1974).
- 74-1181 Willson, R.L.; Wardman, P.; Asmus, K.D., Interaction of dGMP radical with cysteamine and promethazine as possible model of DNA repair, *Nature (London)* 252(5481): 323-4 (1974).
- 74-1182 Teitelbaum, Z.; Meyerstein, D., Kinetics of reaction of ferricenium ion with some free radicals in aqueous solution. A pulse radiolytic study, *Proc. Int. Conf. Coord. Chem.*, 16th, Dublin, Ireland, Aug. 19-24, 1974, R56, 2p. (Chem. Abs. 85: 191822g).
- 74-3003 Simic, M.; Powers, E.L.; Centilli, M.; Cross, M., Radiation sensitization of bacterial spores and phage T7 by NaBr, *Int. J. Radiat. Biol.* 25(1): 43-50 (1974).
- 74-3052 Kemsley, K.; Moore, J.S.; Phillips, G.O.; Sosnowski, A., Reaction of radical probes with substituted phenols as models for the investigation of tyrosine in aldolase and chemically modified aldolase, *Acta Vitaminol. Enzymol.* 28: 263-7 (1974).
- 74-3081 Roberts, P.B.; Fielden, E.M.; Rotilio, G.; Calabrese, L.; Bannister, J.V.; Bannister, W.H., Superoxide dismutase inactivation by radiation-induced radicals: Evidence for histidine residues in the active site, *Radiat. Res.* 60(3): 441-52 (1974).

- 74-7224 Fournier de Violet, P.; Bonneau, R.; Logan, S.R., Laser flash photolytic study of mercury(II) iodide in aqueous solution, *J. Phys. Chem.* 78(17): 1698-701 (1974).
- 74-7296 Chen, S.; Hoffman, M.Z., Reactivity of the carbonate radical in aqueous solution. Tryptophan and its derivatives, *J. Phys. Chem.* 78(21): 2099-102 (1974).
- 74-7554 Barkatt, A.; Ottolenghi, M., Laser flash photolysis of aqueous triiodide solutions, *Mol. Photochem.* 6(3): 253-61 (1974).
- 74-7625 Falcinella, B.; Felgate, P.D.; Laurence, G.S., Aqueous chemistry of thallium(II). Part I. Kinetics of reaction of thallium(II) with cobalt(II) and iron(III) ions and oxidation reduction potentials of thallium(II), *J. Chem. Soc. Dalton Trans.* 1367-73 (1974).
- 74-9006 Snook, M.E.; Hamilton, G.A., Oxidation and fragmentation of some phenyl-substituted alcohols and ethers by peroxydisulfate and Fenton's reagent, *J. Am. Chem. Soc.* 96(3): 860-9 (1974).
- 75-0001 Anbar, M.; Farhataziz; Ross, A.B., Selected specific rates of reactions of transients from water in aqueous solution. II. Hydrogen atom, NSRDS-NBS 51, 1975, 56p.
- 75-0002 Ross, A.B., Selected specific rates of reactions of transients from water in aqueous solution. I. Hydrated electron, Supplemental data, NSRDS-NBS 43-Supp., 1975, 43p.
- 75-0440 Vladimirova, M.V.; Milovanova, A.S., Radiation-chemical oxidation of Ce^{3+} and reduction of Ce^{4+} in the α and γ radiolysis of LiCl solutions, *High Energy Chem. (Engl. Transl.)* 9(4): 309-11 (1975), transl. from *Khim. Vys. Energ.* 9(4): 357-60 (1975).
- 75-1069 Redpath, J.L.; Willson, R.L., Chain reactions and radiosensitization: Model enzyme studies, *Int. J. Radiat. Biol.* 27(4): 389-98 (1975).
- 75-1118 Eriksen, T.E., Pulse radiolytic investigation of the $SO_2^{\cdot-}$ radical ion, *Radiochem.-Radioanal. Lett.* 22(1): 33-40 (1975).
- 75-1119 Zhestkova, T.P.; Pikaev, A.K., Pulse radiolysis of concentrated aqueous solutions of lithium chloride, *High Energy Chem. (Engl. Transl.)* 9(2): 138-42 (1975), transl. from *Khim. Vys. Energ.* 9(2): 165-70 (1975).
- 75-1154 Woods, R.J.; Lesigne, B.; Gilles, L.; Ferradini, C.; Pucheault, J., Pulse radiolysis of aqueous lithium chloride solutions, *J. Phys. Chem.* 79(24): 2700-4 (1975).
- 75-1171 O'Neill, P.; Steenken, S.; Schulte-Frohlinde, D., Formation of radical cations of methoxylated benzenes by reaction with OH radicals, Ti^{2+} , Ag^{2+} , and $SO_4^{\cdot-}$ in aqueous solution. An optical and conductometric pulse radiolysis and in situ radiolysis electron spin resonance study, *J. Phys. Chem.* 79(25): 2773-9 (1975).
- 75-1188 Storer, D.K.; Waltz, W.L.; Brodovitch, J.C.; Eager, R.L., A pulse radiolysis study of some platinum(II) and platinum(IV) complex ions in aqueous solutions. The formation and characterization of platinum(I) and platinum(III) transients, *Int. J. Radiat. Phys. Chem.* 7(6): 693-704 (1975).
- 75-3058 Felicioli, R.; Montagnoli, G.; Monti, S.; Moore, J.S.; Phillips, G.O.; Sosnowski, A., Radiation inactivation of rabbit muscle aldolase, *Int. J. Radiat. Biol.* 27(6): 525-32 (1975).
- 75-5244 Chawla, O.P.; Fessenden, R.W., Electron spin resonance and pulse radiolysis studies of some reactions of $SO_4^{\cdot-}$, *J. Phys. Chem.* 79(24): 2693-700 (1975).
- 75-7110 Chen, S.; Hoffman, M.Z., Effect of pH on the reactivity of the carbonate radical in aqueous solution, *Radiat. Res.* 62(1): 18-27 (1975).
- 75-7222 Treinin, A.; Hayon, E., Charge transfer spectra of halogen atoms in water. Correlation of the electronic transition energies of iodine, bromine, chlorine, hydroxyl, and hydrogen radicals with their electron affinities, *J. Am. Chem. Soc.* 97(7): 1716-21 (1975).
- 75-7313 Chen, S.; Hoffman, M.Z.; Parsons, G.H., Jr., Reactivity of the carbonate radical toward aromatic compounds in aqueous solution, *J. Phys. Chem.* 79(18): 1911-2 (1975).
- 75-7346 Wong, D.; Di Bartolo, B., Evolution of the dihalide ion $Br_2^{\cdot-}$ in aqueous solutions, *J. Photochem.* 4(4): 249-68 (1975).
- 76-1048 Broszkiewicz, R.K., On the effect of Br^- on formation of $Cl_2^{\cdot-}$ in the pulse radiolysis of the aqueous chloride solutions, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 24(2): 123-31 (1976).
- 76-1055 Broszkiewicz, R.K.; Grodkowski, J., Pulse radiolysis studies on aqueous systems $PtCl_4^{2-}-Cl^-$ and $PtCl_4^{2-}-Br^-$, *Int. J. Radiat. Phys. Chem.* 8(3): 359-65 (1976).
- 76-1058 Frank, A.J.; Graetzel, M.; Kozak, J.J., On the reduction of dimensionality in radical decay kinetics induced by micellar systems, *J. Am. Chem. Soc.* 98(11): 3317-21 (1976).
- 76-1105 Buechler, Hch.; Buehler, R.E., The radical ion complex $IOH^{\cdot-}$: spectrum and reactions studied by pulse radiolysis of aqueous iodide solutions, *Chem. Phys.* 16(1): 9-18 (1976).
- 76-1112 Bansal, K.M.; Fessenden, R.W., Pulse radiolysis studies of the oxidation of phenols by $\cdot SO_4^{\cdot-}$ and $Br_2^{\cdot-}$ in aqueous solutions, *Radiat. Res.* 67(1): 1-8 (1976).
- 76-1141 Kim, K.-J.; Hamill, W.H., Direct and indirect effects in pulse irradiated concentrated aqueous solutions of chloride and sulfate ions, *J. Phys. Chem.* 80(21): 2320-5 (1976).
- 76-1143 Bonifacic, M.; Asmus, K.-D., Free radical oxidation of organic disulfides, *J. Phys. Chem.* 80(21): 2426-30 (1976).
- 76-1151 Posener, M.L.; Adams, G.E.; Wardman, P.; Cundall, R.B., Mechanism of tryptophan oxidation by some inorganic radical-anions: A pulse radiolysis study, *J. Chem. Soc., Faraday Trans. 1* 72(10): 2231-9 (1976).
- 76-1171 Broszkiewicz, R.K., The pulse radiolysis study of $NaNO_2$ and $NaNO_3$ solutions, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.* 24(3): 221-9 (1976).
- 76-1181 Frank, A.J.; Graetzel, M.; Henglein, A.; Janata, E., Kinetics of the heterogeneous electron transfer reaction of triplet pyrene in micelles to $Br_2^{\cdot-}$ radicals in aqueous solution, *Int. J. Chem. Kinet.* 8(6): 817-24 (1976).
- 76-1185 Eadsforth, C.V.; Power, D.M.; Thomas, E.W.; Davies, J.V., Investigation of the interaction of alkyl sulphates with serum albumin using the thiocyanate radical ion $\cdot(SCN)_2^-$, *Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med.* 30(5): 449-57 (1976).
- 76-1187 Zemel, H., Electron spin resonance and pulse radiolysis studies of the mechanism and kinetics of $SO_4^{\cdot-}$ reactions with aromatic compounds in aqueous solution, Ph.D., Thesis, Carnegie-Mellon Univ., Pittsburgh, PA, Aug. 5, 1976, 158p.
- 76-1203 Tait, A.M.; Hoffman, M.Z.; Hayon, E., The reaction of radiation-generated free radicals with cobalt(II) and cobalt(III) complexes containing macrocyclic ligands, *Int. J. Radiat. Phys. Chem.* 8(6): 691-6 (1976).
- 76-7279 Burrows, H.D.; Pedrosa de Jesus, J.D., A flash photolytic study of the photo-oxidation of some inorganic anions by the uranyl ion, *J. Photochem.* 5(3/4): 265-75 (1976).
- 76-7587 Scheerer, R.; Graetzel, M., Photo-induced oxydation of carbonate ions by duroquinone, a pathway of oxygen evolution from water by visible light, *Ber. Bunsenges. Phys. Chem.* 80(10): 979-82 (1976).

- 77-0011 Farhataziz; Ross, A.B., Selected specific rates of reactions of transients from water in aqueous solution. III. Hydroxyl radical and perhydroxyl radical and their radical ions, NSRDS-NBS 59, 1977, 113p.
- 77-1001 Neta, P.; Madhavan, V.; Zemel, H.; Fessenden, R.W., Rate constants and mechanism of reaction of $\text{SO}_4^{\cdot-}$ with aromatic compounds, J. Am. Chem. Soc. 99(1): 163-4 (1977).
- 77-1006 Steenken, S.; O'Neill, P.; Schulte-Frohlinde, D., Formation of radical zwitterions from methoxylated benzoic acids. 1. One electron oxidation by Ti^{2+} , Ag^{2+} , and $\text{SO}_4^{\cdot-}$, J. Phys. Chem. 81(1): 26-30 (1977).
- 77-1047 Maruthamuthu, P.; Neta, P., Radiolytic chain decomposition of peroxomonophosphoric and peroxomonosulfuric acids, J. Phys. Chem. 81(10): 937-40 (1977).
- 77-1053 Waltz, W.L.; Brodovitch, J.C.; Kundu, K.P., A pulse radiolysis study of the reactions of platinum(II) glycine complexes in aqueous media, Radiat. Phys. Chem. 10(2): 77-87 (1977).
- 77-1093 Meisel, D.; Matheson, M.S.; Mulac, W.A.; Rabani, J., Transients in the flash photolysis of aqueous solutions of tris(2,2-bipyridine) ruthenium(II) ion, J. Phys. Chem. 81(15): 1449-55 (1977).
- 77-1097 Zansokhova, A.A.; Kabakchi, S.A.; Pikaev, A.K., Mechanism of $\text{Cl}_2^{\cdot-}$ formation in the pulse radiolysis of neutral aqueous solutions of alkali metal chlorides, High Energy Chem. (Engl. Transl.) 11(1): 50-4 (1977), transl. from Khim. Vys. Energ. 11(1): 67-71 (1977).
- 77-1098 Moore, J.S.; Phillips, G.O.; Sosnowski, A., Reaction of the carbonate radical anion with substituted phenols, Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med. 31(6): 603-5 (1977).
- 77-1106 Maruthamuthu, P.; Neta, P., Reactions of phosphate radicals with organic compounds, J. Phys. Chem. 81(17): 1622-5 (1977).
- 77-1132 Buchanan, J.D.; Armstrong, D.A.; Greenstock, C.L.; Ruddock, G.W., Pulse radiolysis of lactate dehydrogenase, Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med. 32(3): 247-57 (1977).
- 77-1164 Asmus, K.-D.; Bahnmann, D.; Bonifacic, M.; Gillis, H.A., Free radical oxidation of organic sulphur compounds in aqueous solution, Faraday Discuss. Chem. Soc. 63: 213-25 (1977).
- 77-1170 Gilbert, C.W.; Ingalls, R.B.; Swallow, A.J., Pulse irradiation of aqueous solutions containing ferrous and chloride ions: reaction between $\text{Cl}_2^{\cdot-}$ and $\text{HO}_2^{\cdot}$, Radiat. Phys. Chem. 10(4): 221-5 (1977).
- 77-1173 Tamba, M.; Badiello, R., Selenium dioxide free radicals: a study by pulse radiolysis, Radiat. Phys. Chem. 10(4): 283-8 (1977).
- 77-1198 Anderson, R.F.; Patel, K.B.; Adams, G.E., Critical residues in D-amino acid oxidase. A pulse-radiolysis and inactivation study, Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med. 32(6): 523-31 (1977).
- 77-1503 Maruthamuthu, P.; Neta, P., unpubl. data.
- 77-3091 Seki, H.; Imamura, M.; Illian, X.A.; Illan, Y.; Stein, G., Oxidation and reduction of cytochrome C with $\text{Br}_2^{\cdot-}$, $\text{O}_2^{\cdot-}$, and $\text{CO}_2^{\cdot-}$ studied by pulse radiolysis, J. Radiat. Res. 18(1): 6 (1977).
- 77-5209 Maruthamuthu, P.; Taniguchi, H., An in situ photolysis-electron spin resonance study of some reactions of phosphate radicals, J. Phys. Chem. 81(20): 1944-8 (1977).
- 78-1075 Maruthamuthu, P.; Neta, P., Phosphate radicals. Spectra, acid-base equilibria, and reactions with inorganic compounds, J. Phys. Chem. 82(6): 710-3 (1978).
- 78-1076 Eibenberger, H.; Steenken, S.; O'Neill, P.; Schulte-Frohlinde, D., Pulse radiolysis and electron spin resonance studies concerning the reaction of $\text{SO}_4^{\cdot-}$ with alcohols and ethers in aqueous solution, J. Phys. Chem. 82(6): 749-50 (1978).
- 78-1093 Hasegawa, K.; Neta, P., Rate constants and mechanisms of reaction of $\text{Cl}_2^{\cdot-}$ radicals, J. Phys. Chem. 82(8): 854-7 (1978).
- 78-3005 Davies, J.V.; Moore, J.S.; Mudher, S., Implication of protein aromatic amino acids in concanavalin A-carbohydrate interactions, Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med. 33(1): 11-9 (1978).
- 78-3007 Tamba, M.; Pini, L.; Badiello, R., Radiation chemical evidence for differences in the amino-acid composition of yeast and horse-liver alcohol dehydrogenases, Int. J. Radiat. Biol. Relat. Stud. Phys. Chem. Med. 33(1): 95-101 (1978).
- 78-1501 Cope, V.W.; Hoffman, M.Z.; Chen, S., Reactivity of the carbonate radical toward metal complexes in aqueous solution, J. Phys. Chem., 82(25): 2665-9 (1978).
- 78-1502 Maruthamuthu, P.; Patterson, L.K.; Ferraudi, G., Redox reactions of free radicals with Ni(II) complexes. A pulse radiolytic study, Inorg. Chem., 11(5): 225-7 (1978).
- 78-1503 Patterson, L.K., to be publ.
- 78-1504 Zemel, H.; Fessenden, R.W., The mechanism of reaction of $\text{SO}_4^{\cdot-}$ with some derivatives of benzoic acid, J. Phys. Chem. 82(25): 2670-6 (1978).
- 78-1505 Madhavan, V.; Levanon, H.; Neta, P., Decarboxylation by $\text{SO}_4^{\cdot-}$ radicals, Radiat. Res. 76(1): 15-22 (1978).
- 78-1506 Lilie, J.; Henglein, A.; Hanrahan, R.J., Reactions of the carbonate radical anion with organic and inorganic solutes in aqueous solution, Abstracts of the American Chemical Soc. Meeting, Miami Beach FL, 11-15 Sep 1978, and Radiat. Phys. Chem., 17(11): 3157-63 (1978).

U.S. DEPT. OF COMM. BIBLIOGRAPHIC DATA SHEET	1. PUBLICATION OR REPORT NO. NSRDS-NBS 65	2. Gov't. Accession No.	3. Recipient's Accession No.
4. TITLE AND SUBTITLE Rate Constants for Reactions of Inorganic Radicals in Aqueous Solution		5. Publication Date June 1979	
		6. Performing Organization Code	
7. AUTHOR(S) Alberta B. Ross and Pedatsur Neta		8. Performing Organ. Report No.	
9. PERFORMING ORGANIZATION NAME AND ADDRESS NATIONAL BUREAU OF STANDARDS DEPARTMENT OF COMMERCE WASHINGTON, DC 20234		10. Project/Task/Work Unit No.	
		11. Contract/Grant No.	
12. SPONSORING ORGANIZATION NAME AND COMPLETE ADDRESS (Street, City, State, ZIP) Same as Item 9.		13. Type of Report & Period Covered N/A	
		14. Sponsoring Agency Code	
15. SUPPLEMENTARY NOTES Library of Congress Catalog Card Number: 79-607030 ☐ Document describes a computer program; SF-185, FIPS Software Summary, is attached.			
16. ABSTRACT (A 200-word or less factual summary of most significant information. If document includes a significant bibliography or literature survey, mention it here.) Rate constants have been compiled for reactions of various transient inorganic radicals produced by radiolysis or photolysis in aqueous solution. Data are included for the carbonate radical, sulfate radical, phosphate radical, nitrate radical and other nitrogen-, sulfur- and selenium-containing radicals, and the halide and pseudohalide radicals Cl_2^- , Br_2^- , I_2^- , and $(\text{SCN})_2^-$. The radicals react with other inorganic ions, as well as aliphatic, aromatic and heterocyclic compounds.			
17. KEY WORDS (six to twelve entries; alphabetical order; capitalize only the first letter of the first key word unless a proper name; separated by semicolons) Aqueous solution; carbonate radical; chemical kinetics; halogen radical anions; nitrate radical; oxyanion radicals; phosphate radical; photolysis; radiolysis; rates; selenium radicals; sulfate radical; thiocyanate radical anion.			
18. AVAILABILITY ☒ Unlimited ☐ For Official Distribution. Do Not Release to NTIS ☒ Order From Sup. of Doc., U.S. Government Printing Office, Washington, DC 20402, SD Stock No. 003-003-02072-9 ☐ Order From National Technical Information Service (NTIS), Springfield, VA. 22161		19. SECURITY CLASS (THIS REPORT) UNCLASSIFIED	21. NO. OF PRINTED PAGES 62
		20. SECURITY CLASS (THIS PAGE) UNCLASSIFIED	22. Price \$3.50

**Where can you find
all the reference data
you need?**

**Right in the
Journal of Physical
and Chemical
Reference Data!**


Now in its sixth year, this valuable publication has proved that it fills the important gaps for you in the literature of the physical sciences.

Published by the American Institute of Physics and the American Chemical Society for the National Bureau of Standards, this quarterly gives you quantitative numerical data, with recommended values and uncertainty limits chosen by experts in the field.

Critical commentary on methods of measurement and sources of error, as well as full references to the original literature, is an integral part of each of your four issues a year.

Can you afford to be without this prime source of reliable data on physical and chemical properties? To start receiving your copies, just fill in the order form and drop into the mail. If you do use a purchase order, please attach the printed form as this will help us to expedite your order.

Send for complete list of reprints!



Journal of Physical and Chemical Reference Data
American Chemical Society
1155 Sixteenth Street, N.W., Washington, D.C. 20036

Yes, I would like to receive the JOURNAL OF PHYSICAL AND CHEMICAL REFERENCE DATA at the one-year rate checked below:

Name

☐ Home
☐ Business

Street

City State Zip

Bill me ☐

Bill company or school ☐

Payment enclosed ☐

Members Nonmembers

*U.S., Canada,
Mexico*

☐ \$25.00 ☐ \$100.00

Other Countries

☐ \$29.00 ☐ \$104.00

Please Attach This Order Form To Purchase Order.

Important Additions

TO THE JOURNAL OF PHYSICAL AND CHEMICAL REFERENCE DATA

Three comprehensive reference volumes, each, as the Journal itself, published by the American Institute of Physics and the American Chemical Society for the National Bureau of Standards . . . your triple assurance of their accuracy, immediacy, and usefulness.

Supplement No. 1 to Vol. 2

"PHYSICAL AND THERMODYNAMIC PROPERTIES OF ALIPHATIC ALCOHOLS"

by R. C. Wilhoit and B. J. Zwolinski, *Thermodynamics Research Center, Department of Chemistry, Texas A & M University*

Represents the most exhaustive review and critical analysis of selected physical and thermodynamic properties of aliphatic alcohols that has been published in the world literature of chemistry.

Supplement No. 1 to Vol. 3

"THERMAL CONDUCTIVITY OF THE ELEMENTS: A COMPREHENSIVE REVIEW"

by C. Y. Ho, R. W. Powell, and P. E. Liley, *Thermophysical Properties Research Center, Purdue University, West Lafayette, Indiana*

This comprehensive review of the world's thermal conductivity data presents recommended or estimated values for all 105 elements.

Supplement No. 1 to Vol. 6

ENERGETICS OF GASEOUS IONS

by H. M. Rosenstock, K. Draxl, B. Steiner, and J. T. Herron, *National Bureau of Standards*

Provides a comprehensive body of critically evaluated information on ionization potentials, appearance potentials, electron affinities and heats of formation of gaseous positive and negative ions. It is a complete revision and extension of the earlier reference work, "Ionization Potentials, Appearance Potentials and Heats for Formation of Gaseous Positive Ions," NSRDS-NBS 26.

Business Operations—Books and Journals Dept.
American Chemical Society
1155 16th Street, N.W.
Washington, D.C. 20036

Please send _____ copies of _____ at \$_____.

A. "PHYSICAL AND THERMODYNAMIC PROPERTIES OF ALIPHATIC ALCOHOLS." (First supplement to Vol. 2 of the *Journal of Physical and Chemical Reference Data*.) Hard Cover: \$33.00. Soft Cover: \$30.00.

B. "THERMAL CONDUCTIVITY OF THE ELEMENTS. A COMPREHENSIVE REVIEW." (First supplement to Vol. 3 of the *Journal of Physical and Chemical Reference Data*.) Hard Cover: \$60.00. Soft Cover: \$55.00.

C. "ENERGETICS OF GASEOUS IONS." (First supplement to Vol. 6 of the *Journal of Physical and Chemical Reference Data*.) Hard Cover: \$70.00. Soft Cover: \$65.00.

☐ I am enclosing a check ☐ I am enclosing a money order

Name _____

Address _____

City _____ State _____ Zip Code _____

Title _____ Employer _____

Please add \$1.50 extra for foreign postage and handling.

**Announcement of New Publications in
National Standard Reference Data Series**

Superintendent of Documents,
Government Printing Office,
Washington, D.C. 20402

Dear Sir:

Please add my name to the announcement list of new publications to be issued in the series: National Standard Reference Data Series—National Bureau of Standards.

Name_____

Company_____

Address_____

City_____State_____Zip Code_____

(Notification Key N-519)

NBS TECHNICAL PUBLICATIONS

PERIODICALS

JOURNAL OF RESEARCH—The Journal of Research of the National Bureau of Standards reports NBS research and development in those disciplines of the physical and engineering sciences in which the Bureau is active. These include physics, chemistry, engineering, mathematics, and computer sciences. Papers cover a broad range of subjects, with major emphasis on measurement methodology, and the basic technology underlying standardization. Also included from time to time are survey articles on topics closely related to the Bureau's technical and scientific programs. As a special service to subscribers each issue contains complete citations to all recent NBS publications in NBS and non-NBS media. Issued six times a year. Annual subscription: domestic \$17.00; foreign \$21.25. Single copy, \$3.00 domestic; \$3.75 foreign.

Note: The Journal was formerly published in two sections: Section A "Physics and Chemistry" and Section B "Mathematical Sciences."

DIMENSIONS/NBS

This monthly magazine is published to inform scientists, engineers, businessmen, industry, teachers, students, and consumers of the latest advances in science and technology, with primary emphasis on the work at NBS. The magazine highlights and reviews such issues as energy research, fire protection, building technology, metric conversion, pollution abatement, health and safety, and consumer product performance. In addition, it reports the results of Bureau programs in measurement standards and techniques, properties of matter and materials, engineering standards and services, instrumentation, and automatic data processing.

Annual subscription: Domestic, \$11.00; Foreign \$13.75

NONPERIODICALS

Monographs—Major contributions to the technical literature on various subjects related to the Bureau's scientific and technical activities.

Handbooks—Recommended codes of engineering and industrial practice (including safety codes) developed in cooperation with interested industries, professional organizations, and regulatory bodies.

Special Publications—Include proceedings of conferences sponsored by NBS, NBS annual reports, and other special publications appropriate to this grouping such as wall charts, pocket cards, and bibliographies.

Applied Mathematics Series—Mathematical tables, manuals, and studies of special interest to physicists, engineers, chemists, biologists, mathematicians, computer programmers, and others engaged in scientific and technical work.

National Standard Reference Data Series—Provides quantitative data on the physical and chemical properties of materials, compiled from the world's literature and critically evaluated. Developed under a world-wide program coordinated by NBS. Program under authority of National Standard Data Act (Public Law 90-396).

NOTE: At present the principal publication outlet for these data is the Journal of Physical and Chemical Reference Data (JPCRD) published quarterly for NBS by the American Chemical Society (ACS) and the American Institute of Physics (AIP). Subscriptions, reprints, and supplements available from ACS, 1155 Sixteenth St. N.W., Wash., D.C. 20056.

Building Science Series—Disseminates technical information developed at the Bureau on building materials, components, systems, and whole structures. The series presents research results, test methods, and performance criteria related to the structural and environmental functions and the durability and safety characteristics of building elements and systems.

Technical Notes—Studies or reports which are complete in themselves but restrictive in their treatment of a subject. Analogous to monographs but not so comprehensive in scope or definitive in treatment of the subject area. Often serve as a vehicle for final reports of work performed at NBS under the sponsorship of other government agencies.

Voluntary Product Standards—Developed under procedures published by the Department of Commerce in Part 10, Title 15, of the Code of Federal Regulations. The purpose of the standards is to establish nationally recognized requirements for products, and to provide all concerned interests with a basis for common understanding of the characteristics of the products. NBS administers this program as a supplement to the activities of the private sector standardizing organizations.

Consumer Information Series—Practical information, based on NBS research and experience, covering areas of interest to the consumer. Easily understandable language and illustrations provide useful background knowledge for shopping in today's technological marketplace.

Order above NBS publications from: Superintendent of Documents, Government Printing Office, Washington, D.C. 20402.

Order following NBS publications—NBSIR's and FIPS from the National Technical Information Services, Springfield, Va. 22161.

Federal Information Processing Standards Publications (FIPS PUB)—Publications in this series collectively constitute the Federal Information Processing Standards Register. Register serves as the official source of information in the Federal Government regarding standards issued by NBS pursuant to the Federal Property and Administrative Services Act of 1949 as amended, Public Law 89-306 (79 Stat. 1127), and as implemented by Executive Order 11717 (38 FR 12315, dated May 11, 1973) and Part 6 of Title 15 CFR (Code of Federal Regulations).

NBS Interagency Reports (NBSIR)—A special series of interim or final reports on work performed by NBS for outside sponsors (both government and non-government). In general, initial distribution is handled by the sponsor; public distribution is by the National Technical Information Services (Springfield, Va. 22161) in paper copy or microfiche form.

BIBLIOGRAPHIC SUBSCRIPTION SERVICES

The following current-awareness and literature-survey bibliographies are issued periodically by the Bureau:

Cryogenic Data Center Current Awareness Service. A literature survey issued biweekly. Annual subscription: Domestic, \$25.00; Foreign, \$30.00.

Liquidified Natural Gas. A literature survey issued quarterly. Annual subscription: \$20.00.

Superconducting Devices and Materials. A literature survey issued quarterly. Annual subscription: \$30.00. Send subscription orders and remittances for the preceding bibliographic services to National Bureau of Standards, Cryogenic Data Center (275.02) Boulder, Colorado 80302.

U.S. DEPARTMENT OF COMMERCE
National Bureau of Standards
Washington, D.C. 20234

OFFICIAL BUSINESS

Penalty for Private Use, \$300

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF COMMERCE
COM-215



SPECIAL FOURTH-CLASS RATE
BOOK
