


Library
National Wetlands Research Center
U. S. Fish and Wildlife Service
700 Cajundome Boulevard
Lafayette, La. 70506

Biological Services Program

FWS/OBS-82/10.1
FEBRUARY 1982

HABITAT SUITABILITY INDEX MODELS: COASTAL STOCKS OF STRIPED BASS



U. S. Fish and Wildlife Service

S. Department of the Interior

SK
361
.U54
no.82-
10.1

The Biological Services Program was established within the U.S. Fish and Wildlife Service to supply scientific information and methodologies on key environmental issues that impact fish and wildlife resources and their supporting ecosystems. The mission of the program is as follows:

- To strengthen the Fish and Wildlife Service in its role as a primary source of information on national fish and wildlife resources, particularly in respect to environmental impact assessment.
- To gather, analyze, and present information that will aid decisionmakers in the identification and resolution of problems associated with major changes in land and water use.
- To provide better ecological information and evaluation for Department of the Interior development programs, such as those relating to energy development.

Information developed by the Biological Services Program is intended for use in the planning and decisionmaking process to prevent or minimize the impact of development on fish and wildlife. Research activities and technical assistance services are based on an analysis of the issues, a determination of the decisionmakers involved and their information needs, and an evaluation of the state of the art to identify information gaps and to determine priorities. This is a strategy that will ensure that the products produced and disseminated are timely and useful.

Projects have been initiated in the following areas: coal extraction and conversion; power plants; geothermal, mineral and oil shale development; water resource analysis, including stream alterations and western water allocation; coastal ecosystems and Outer Continental Shelf development; and systems inventory, including National Wetland Inventory, habitat classification and analysis, and information transfer.

The Biological Services Program consists of the Office of Biological Services in Washington, D.C., which is responsible for overall planning and management; National Teams, which provide the Program's central scientific and technical expertise and arrange for contracting biological services studies with states, universities, consulting firms, and others; Regional Staffs, who provide a link to problems at the operating level; and staffs at certain Fish and Wildlife Service research facilities, who conduct in-house research studies.

FWS/OBS-82/10.1
February 1982

**HABITAT SUITABILITY INDEX MODELS:
COASTAL STOCKS OF STRIPED BASS**

by

**Kark B. Bain
and**

Jane L. Bain

**Massachusetts Cooperative Fishery Research Unit
Department of Forestry and Wildlife Management
University of Massachusetts
Amherst, Massachusetts 01003**

Project Officer

**Carroll L. Cordes
National Coastal Ecosystems Team
U.S. Fish and Wildlife Service
1010 Gause Boulevard
Slidell, Louisiana 70458**

Performed for

**National Coastal Ecosystems Team
Office of Biological Services
Fish and Wildlife Service
U.S. Department of the Interior
Washington, D.C. 20240**

This report should be cited:

Bain, K.B., and J.L. Bain. 1982. Habitat suitability index models: Coastal stocks of striped bass. U.S. Fish and Wildlife Service, Office of Biological Services, Washington, D.C. FWS/OBS-82/10.1. 29 pp.

PREFACE

The habitat use information and habitat suitability index (HSI) models presented in this report on coastal stocks of striped bass are intended for use in impact assessment and habitat management activities. The models were developed from a review and synthesis of existing information and are scaled to produce an index of habitat suitability between 0 (unsuitable habitat) and 1 (optimally suitable habitat). Assumptions used to transform habitat use information into the HSI models, and guidelines for model applications, including techniques for measuring model variables, are described.

The HSI models presented herein are hypotheses of species-habitat relationships, not statements of proven cause and effect relationships. The models have not been field-tested, but they have been applied to four hypothetical data sets which are presented and discussed. For this reason, the U.S. Fish and Wildlife Service encourages model users to convey comments and suggestions that may help increase the utility and effectiveness of this habitat-based approach to fish and wildlife planning. Please send any comments or suggestions you may have on the striped bass HSI models to the National Coastal Ecosystem Team, U.S. Fish and Wildlife Service, 1010 Gause boulevard, Slidell, Louisiana 70458.



CONTENTS

	<u>Page</u>
PREFACE	iii
ACKNOWLEDGMENTS	vi
HABITAT USE INFORMATION	1
Age, Growth, and Food	1
Reproduction	2
Specific Habitat Requirements	3
HABITAT SUITABILITY INDEX MODEL	8
Model Applicability	8
Model Description	9
Suitability Index Graphs for Model Variables	13
Component Index Equations	17
HSI Getermination	17
Field Use of the Model.....	18
Interpreting Model Outputs	18
ADCITIONAL HABITAT SUITABILITY INDEX MODELS	22
LITERATURE CITEG	23

ACKNOWLEDGMENTS

Development of the habitat suitability index models for the striped bass was monitored and evaluated by Dr. John Eoreman, National Marine Fisheries Service, and Dr. John J. Ney, Virginia Polytechnic Institute and State University. Messrs. James Terrell, James M. Barkuloo, and John Cooper, Fish and Wildlife Service, respectively provided technical assistance in model construction, peer review, and preparation of the manuscript for publication. Useful comments and suggestions were also received from field biologists. Funding for model construction and publication was provided by the Fish and Wildlife Service.

STRIPEC BASS (Morone saxatilis)

HABITAT USE INFORMATION

The striped bass is a major sport and commercial fish. The species has a large natural range that includes freshwater, estuarine, and marine habitats. Although this fish is tolerant of a variety of environmental conditions, several physicochemical and biological factors limit striped bass survival and abundance.

Striped bass populations occur along the Atlantic coast from the St. Lawrence River, Canada, to the St. Johns River, Florida. Along the Gulf of Mexico, striped bass were once common in riverine and estuarine habitats from the Suwannee River in north Florida to Lake Pontchartrain, Louisiana (Hardy 1978; Setzler et al. 1980; Crateau et al. 1981). Following introduction into California's Sacramento and San Joaquin River estuary in 1879, the striped bass population expanded tremendously and now dominates the sport fishery in the estuary (Chadwick et al. 1977). Since 1879, striped bass have been found from the Columbia River, Washington, to Ensenada, Mexico. Striped bass populations have also been established in many freshwater reservoirs across the United States.

One of the most important spawning and nursery grounds for striped bass is Chesapeake Bay and its tributaries (Kernehan et al. 1981). Chesapeake Bay may account for as much as 90% of the recruitment in the Atlantic coastal fishery and is also a major contributor to the fishery in the lower Hudson River and New York Bight (Berggren and Lieberman 1978). Albemarle Sound and the Hudson River are also important sources of striped bass on the Atlantic coast (Raney et al. 1954; Berggren and Lieberman 1978).

Age, Growth, and Food

For purposes of this habitat account, three major developmental stages are recognized for striped bass: adult, larval, and juvenile. Female striped bass generally reach maturity at 3 to 4 years of age and at total lengths (TL) of 432 to 457 mm, while males mature at 2 years and 174 to 254 mm TL (Merriman 1941; Raney 1952; Lewis 1962). Age and size at maturity, however, may vary significantly between stocks. Adult striped bass feed primarily on fish. Manooch (1973) suggested that soft-rayed fish species are preferred prey, but other workers consider striped bass to be generalists in prey choice (Merriman 1941; deSilva 1973).

The larval stage of development begins with the hatching of the fertilized egg and ends after 35 to 50 days. At approximately 8 days and 6 to 7 mm TL, striped bass larvae begin active feeding (Doroshev 1970). Larvae less than 10 mm feed primarily on crustacean nauplii. At sizes greater than 10 mm, they feed primarily on larger zooplankton and macroinvertebrates (Humphries

and Cumming 1973). The availability of sufficient concentrations of suitable prey during the first several days of feeding is a critical factor influencing larval survival (Setzler et al. 1980; Cooper and Polgar 1981; Eldridge et al. 1981). Miller (1977) estimated that a minimum concentration of 1,864 nauplii/l is required for successful initial feeding.

The juvenile stage is relatively long compared to other life stages and encompasses the period from larval transformation at 35 to 50 days until maturity. Young striped bass feed on zooplankton, macroinvertebrates, and fish. Stevens (1966) reported that juveniles in the Sacramento and San Joaquin River estuary feed mainly on invertebrates in the winter and spring. When small fish prey become available in the summer, juvenile striped bass change their diet and feed upon them. Juvenile striped bass begin feeding on larger fish in their second summer (Stevens 1966). In general, juveniles are not highly selective and consume food items based on availability (Heubach et al. 1963; Stevens 1966; Hester and Stevens 1970; Kanooch 1973; Goynton et al. 1981).

Reproduction

Spawning may begin in mid-February in the southern portion of the striped bass range, whereas, in the extreme northern portions of the range, spawning does not begin until June or July (Hardy 1978). A reservoir of small eggs is maintained in the ovaries of mature females. Egg maturation begins slowly in summer and proceeds rapidly as the spawning season is approached (Jackson and Tiller 1952; Lewis 1962). Female striped bass produce approximately 14,000 to 65,000 eggs annually in initial spawning years, while very old and large individuals may produce over 5,000,000 eggs (Jackson and Tiller 1952; Raney 1952).

Adult striped bass migrate to fresh or nearly freshwater to spawn. Typically, spawning takes place in riverine habitat closely associated with the upper estuary, approximately within the first 45 km (28 mi) of freshwater (Talbot 1966). In some stocks, particularly those in estuarine systems lacking pronounced tidal cycles spawning may occur at greater distances upstream of the estuary.

Males reach the spawning grounds before females. As spawning occurs, several males will surround a single female and create commotion by splashing on the surface of the water. Females broadcast their semibuoyant eggs near the water surface when fertilization occurs (Merriman 1941; Raney 1952). Striped bass refrain from feeding only immediately before and during spawning (Morgan and Gerlach 1950; Hollis 1952). Depending on the estuary, there may be one to three peaks in spawning each season which are thought to result from major increases in water temperature (Hardy 1978). Reports documenting spawning time for striped bass vary widely. Spawning activity has been noted in late afternoon and early evening (Morgan and Gerlach 1950), as well as late evening and early morning (Hardy 1978).

Environmental factors play a major role in determining the spawning success of striped bass stocks (Talbot 1966; Ulanowicz and Polgar 1980). Water volume in the spawning river is a significant factor determining spawning habitat suitability (Fish and McCoy 1959; Turner and Chadwick 1972). Suitability of a spawning area appears to increase with greater river discharge (expressed as the percent of natural flow). Fish and McCoy (1959)

found that a sustained minimum flow was necessary for suitable spawning conditions in the Roanoke River, North Carolina, and rapid fluctuations in streamflow were detrimental to spawning.

Shannon and Smith (1967) reported that optimal spawning temperatures are approximately 17° to 19°C (63° to 66°F) with no spawning observed below 13°C (55°F) or above 22°C (72°F). With minor variations, these temperature ranges correspond with observations by Calhoun et al. (1950), Raney (1952), Dickson (1957), Farley (1966), and Talbot (1966). Calhoun et al. (1950) and Farley (1966) reported that a sudden drop in water temperature can terminate spawning.

Salinity and total dissolved solids (TDS) concentrations are important factors during the spawning period and may be responsible for deterring spawning or reducing spawning success. Farley (1966) found that the number of eggs deposited reached a maximum when TDS were less than 0.18 parts per thousand (ppt). Spawning migrations into the San Joaquin River, California, did not occur at a critical salinity concentration of 0.35 ppt (TDS) (Radtke and Turner 1967). However, highly successful spawning was observed in the Chesapeake and Delaware Canal at salinities of 0.70 to 1.5 ppt (TDS) by Johnson and Koo (1975). Stevens (1979) reported that striped bass may not spawn where salinities exceed 5 ppt.

Specific Habitat Requirement²

Striped bass occupy various habitats depending on life stage. Variations in habitat utilization also occur between individual stocks. Known striped bass habitat requirements are presented for each life stage in the following sections.

Adult. In some striped bass stocks from Cape Hatteras, North Carolina, north to New England (predominantly Chesapeake Bay), substantial numbers of striped bass leave the estuaries and migrate along the coast. The migrating fish move north in the summer and south in the winter (Merriman 1941; Clark 1968). Large adults, predominantly females, move farther distances on their migratory sojourns. Adults in the southern portion of the range and on the Pacific coast tend to remain in the estuary or in nearby offshore waters. Sandy beaches, rocky shores, and shallow bays are inhabited in both marine and estuarine environments (Bigelow and Schroeder 1953). Striped bass are generally not found more than 6 to 8 km (4 to 5 mi) offshore at any time.

Limited information is available concerning the physicochemical requirements and tolerances of adult striped bass. The adults appear to have similar water temperature and dissolved oxygen requirements as juveniles. Preferred temperatures vary depending on ambient acclimation temperatures (Meldrim and Gift 1971). The maximum upper avoidance temperature is 34°C (93°F) for striped bass acclimated to 27°C (81°F) in late August, while striped bass acclimated to 5°C (41°F) in December avoid 13°C (55°F) water. A maximum preferred temperature of 25° to 27°C (77° to 81°F) during the growing season was reported by Merriman (1941). Adult striped bass are known to avoid areas of 44% (of saturation) or less in dissolved oxygen concentration (Meldrim et al. 1974).

Information on specific causes of mortality in adults is lacking, although sport and commercial fishing are suspected of having a significant effect on some populations.

Egg. Striped bass eggs are spherical, semibuoyant, nonadhesive, and approximately 1.3 mm in diameter at fertilization (Pearson 1938). Within 12 hours of release, eggs will be water hardened and approximately 3.4 to 3.8 mm in diameter (Pearson 1938; Johnson and Koo 1975). Survival of striped bass eggs to hatching is primarily associated with narrow tolerances to certain physicochemical factors including temperature, dissolved oxygen, and current velocity (Cooper and Polgar 1981).

A temperature range of 17° to 20°C (63° to 68°F) is optimal for egg survival (Barkuloo 1970; Goroshev 1970; Morgan et al. 1981). Morgan and Rasin (1973) and Rogers et al. (1977) indicated that egg survival rapidly declines as water temperature approaches 23°C (73°F) and gradually declines as water temperature drops below 17°C (63°F), with few eggs surviving at 12°C (54°F). These temperature ranges are similar to those required for spawning.

Low dissolved oxygen concentrations (2.0 to 3.5 mg/l) were determined to be responsible for the absence of eggs and larvae in the Celaware River (Murawski 1969; Chittenden 1971). Turner and Farley (1971) reported that even moderate reductions in dissolved oxygen concentrations (from 5 to 4 mg/l) decreased the survival of eggs.

A critical factor for egg survival is sufficient current velocity to maintain egg suspension (Talbot 1966). Either tidal turbulence or river discharge can provide sufficient water movement to suspend the eggs. A minimum average velocity of 30 cm/s (1 ft/s) is necessary to prevent the concentration of eggs on the bottom (Albrecht 1964). Egg distribution in the water column appears to be random at velocities in excess of 30 cm/s. The density gradient of the water column may also influence egg suspension since striped bass eggs are only slightly heavier than pure water (specific gravity = 1.0005, Hardy 1978).

Certain factors that are usually regarded as important determinants of egg survival in other fish species do not appear to be important to striped bass. Salinities typically encountered by eggs are not detrimental to survival, although low salinity is considered optimal for water hardening (Albrecht 1964; Morgan et al. 1981). While high turbidity and heavy suspended sediment loads have been found to be detrimental to egg survival for many fish species, striped bass eggs appear to be adapted to these conditions. Neither turbidity nor suspended sediments have been observed to significantly decrease hatching success (Talbot 1966; Schubel and Auld 1974). Although predation may have a significant effect on egg survival, quantitative estimates of the magnitude of predation are lacking for larval fish in general (Dahlberg 1979).

The egg stage is brief in comparison to other striped bass life stages. Several authors documented hatching at approximately 48 hours after fertilization at a temperature of 18°C (64°F). However, hatching time varies from 29 hours at 22°C (72°F) to 80 hours at 11°C (52°F) (Pearson 1938; Raney 1952; Mansueti 1958; Hardy 1978).

Larval. The larval stage is the period from hatching to the time when the appearance of the young corresponds with that of an adult. The duration of this stage and the size ranges of striped bass larvae have been summarized from a detailed review by Setzler et al. (1980). The larval stage is commonly divided into three distinct phases: yolk sac, finfold, and post-finfold. At hatching, striped bass range from 2.0 to 3.7 mm long and possess an easily identified yolk sac. The yolk sac phase lasts 3 to 9 days and is terminated when the yolk sac is absorbed at about 5 to 8 mm TL (Figure 1a). The finfold (metamorphosing) phase lasts approximately 11 days during which larvae attain a maximum length of 12 mm (Figure 1b). The final metamorphosing phase, post-finfold, lasts 20 to 30 days with a maximum larval length of 20 mm.

Larval survival is a crucial factor in the determination of adult population size (Eldridge et al. 1981). As in the egg stage, larval survival is associated with a narrow tolerance range for certain physicochemical factors. In natural estuarine systems, both food availability and environmental conditions influenced by climatic events are particularly important for early striped bass survival (Cooper and Polgar 1981).


Dissolved oxygen, temperature, and salinity appear to play important roles in determining larval survival. Striped bass larvae need a minimum of 3 mg/l dissolved oxygen to survive (Chittenden 1971). Even moderate reductions in dissolved oxygen (from 5 to 4 mg/l) reduced the survival of larvae (Turner and Farley 1971). Dissolved oxygen requirements for larvae are essentially identical to those reported for striped bass eggs. A temperature range of 18° to 21°C (64° to 70°F) is considered optimal for larvae (Rogers et al. 1977), but temperatures from 12° to 23°C (54° to 73°F) are tolerated (Doroshev 1970). A salinity range of 3 to 7 ppt is optimal for growth and survival (Lal et al. 1977). However, striped bass larvae can tolerate salinities from 0 to 15 ppt (Albrecht 1964).

Other factors which may influence striped bass larval survival are not well understood or do not appear to be significant. While they may initially require turbulence to maintain position in the water column, striped bass larvae rapidly become very motile, positively phototactic, and continually self suspended (Doroshev 1970). The larvae characteristically remain in the open surface waters of the estuary (Raney 1952).

A potentially significant cause of larval mortality is predation. Data on losses of larvae to predators, however, are lacking for most fish species (Flay 1974; Dahlberg 1979) and striped bass in particular (Setzler et al. 1980).

Juvenile. Juvenile striped bass (Figure 1c) are able to tolerate a considerably broader range of environmental conditions than egg and larval life stages. Some habitat-requirements for juveniles are similar to those of adult striped bass.


As young juvenile striped bass grow, they move into nearshore areas and gradually migrate to higher salinity portions of the lower estuary (Raney 1952; Calhoun 1953). Bason (1971) reported that the largest concentrations of juveniles were in protected areas of moderate salinity. Shoals have also been reported as nursery areas for striped bass by Sasaki (1966) and Carlson and McCann (1968). Rathjen and Miller (1957) noted that the largest catches of



a.



b.



c.

Figure 1. Larval and juvenile development stages of the striped bass. Figure 1a depicts the yolk sac larval phase of a 5.3-mm TL specimen (from Mansueti 1958). Figure 1b shows a 12-mm TL striped bass larva beginning the post-finfold phase, prior to full development of the second dorsal fin (from Kansueti 1958). Figure 1c depicts the juvenile development stage of a 130-mm TL specimen (from Pearson 1940). Development stages are not drawn to scale.

juvenile striped bass were near clean sandy bottoms, although differential sampling efficiency may have been responsible for high catches in these areas. Apparently juvenile striped bass will use various nearshore microhabitats and do not appear to require specific microhabitat conditions.

Temperature and dissolved oxygen are important factors affecting growth and survival of juvenile striped bass. A temperature range of 14° to 21°C (57° to 70°F) during the growing season is optimal (Krouse 1968; Doroshev 1970). However, a temperature range of 10° to 27°C (50° to 81°F) can be tolerated by juveniles (Davies 1970; Hester and Stevens 1970). For dissolved oxygen levels, Krouse (1968) found no survival of juvenile striped bass at 1 mg/l, intermediate survival at 3 mg/l, and high survival at 5 mg/l.

Although large piscivores, such as bluefish and weakfish, probably prey on juvenile striped bass, definitive information on, or confirmation of, the extent of predation is not currently available (Setzler et al. 1980).

Special considerations. Striped bass populations are characterized by dramatic fluctuations in year-class size. Traditionally, survival during the larval stage has been considered to determine year-class size in species exhibiting large fluctuations in abundance (May 1974; Dahlberg 1979). Recent research indicates that this concept is particularly applicable to striped bass (Polgar et al. 1975; Cooper and Polgar 1981; Eldridge et al. 1981). Merriman (1941) first noted the occasional occurrence of unusually large year classes of striped bass and suggested a relationship between certain climatic conditions and strong year classes.

While some conclusions are tentative, current thinking links strong striped bass year classes to high food availability (abundant zooplankton) during the larval stage (Heinle et al. 1975; Eldridge et al. 1981). Although zooplankton abundance fluctuates widely over time in any estuary, the potential for an abundance of zooplankton appears to be related to estuarine productivity. The level of productivity in an estuary is a function of both freshwater nutrient (detritus) input to the estuary (Biggs and Flemer 1972; Hobbie et al. 1973; Saila 1973; Day et al. 1975; Polgar et al. 1975) and detritus production in the salt marsh (Teal 1962; Odum and Heald 1973; Reimold et al. 1973; Stevenson et al. 1975). Detrital input to the estuary from freshwater inflow is typically greatest during the late winter and early spring. Heinle et al. (1975) and others have proposed that certain climatic events, which affect nutrient release from the salt marsh and nutrient contribution by freshwater input, can influence plankton abundance during the critical larval stage and consequently affect year-class size in striped bass.

The complex relationship of estuarine productivity, zooplankton abundance, and larval striped bass survival appears to have important ramifications for the success of striped bass in estuarine systems and on the evaluation of habitat suitability. Apparently, even optimal habitat will only occasionally produce ideal conditions for striped bass survival. However, the loss of habitat suitability, which would diminish the potential to produce strong year classes, may ultimately have serious consequences for maintenance of a viable striped bass population.

HABITAT SUITABILITY INDEX (HSI) MODEL

Model Applicability

Striped bass are widely distributed and thrive in many habitats. Few generalized statements concerning habitat requirements will be precisely applicable to all populations. Consequently, it is desirable that information pertaining to a particular striped bass stock under consideration be evaluated with regard to model criteria.

The use of this model is not always appropriate, such as where partial or extensive reduction in habitat availability has occurred due to toxic wastes. The unavailability of potential habitat resulting from the presence of environmental contaminants will interfere with proper interpretation of model applications.

Several model variables utilize average values for certain habitat parameters. While no clear guidance is available in striped bass literature regarding tolerances to rapidly fluctuating conditions, sudden changes in water temperature and river discharge during spawning and early life are detrimental. Caution should be used in calculating average values in situations marked by extreme variability.

Geographic area. This model is suitable for use on estuarine-associated striped bass stocks located on the Atlantic, Pacific, or Gulf of Mexico coasts of the United States. The model and included variables are generalized to reflect the life cycle and requirements of striped bass throughout its coastal range. Therefore, minor deviations in habitat requirements may occur for any specific stock.

Season. The habitat suitability index model is designed for use in year-round applications. In applications involving only portions of the life cycle, the HSI values obtained will apply to seasons in which the life stage(s) of interest occur. Some of the variables included in the model pertain to environmental conditions that would only occur during a particular season. However, it is assumed that as long as those life requisites are available, habitat suitability in other seasons is not limiting.

Cover types. Striped bass typically use at least two habitats during their life cycle: riverine and estuarine. In many stocks, spawning occurs near the transition between riverine and estuarine habitats, whereas, in other stocks, spawning occurs well within riverine habitat. Similarly, egg development varies in its proximity to either type of habitat.

This model is not intended for the evaluation of marine habitat, although in some stocks the adult stage uses this habitat type. When particular life stages are not associated with the estuarine and near-riverine habitats, the model can be manipulated to account for the absence of these life stages.

Minimum habitat area. The minimum habitat area is that area of contiguous suitable habitat that is required for striped bass to live and reproduce. No minimum habitat size requirements for this species or the distinct life stages have been established.

Verification level. The acceptable output of this HSI model is an index between 0 and 1 which is believed to have a positive relationship to carrying capacity. Hypothetical data sets were used to verify that HSI's determined with the striped bass models were reasonable and acceptable. These data sets and their relationship to model verification are discussed later.

Two biological experts outside of the U.S. Fish and Wildlife Service were identified to review and evaluate the striped bass HSI models throughout their development. Ideas and suggestions from these experts were incorporated into the model-building effort. These experts were: Dr. John Boreman, National Marine Fisheries Service, Woods Hole, Massachusetts; and Dr. John Ney, Virginia Polytechnic Institute and State University, Blacksburg, Virginia.

Model Description

The striped bass HSI model is composed of individual components corresponding to the spawning, egg, larval, juvenile, and adult life stages. When all components are used, the model will produce one habitat suitability index value for the entire life cycle. Individual life stage HSI values can be calculated by using individual components. Partial life cycle HSI values can also be calculated.

The model considers the quality of habitat requirements (variables) for each life stage of the species. Habitat requirements are combined for life requisite categories. It is assumed that habitat suitability is primarily associated with water quality (physicochemical conditions) during most life stages. Food availability and water quantity are considered particularly important life requisites in some life stages. The spawning and egg stages are generally associated with riverine habitat, while the later life stages predominantly occur in estuarine habitats. Figure 2 illustrates how the HSI is related to variables, life requisites, and life stages for striped bass.

Following sections document the assumptions and justification for the variables and equations included in this model. Specifically, these sections cover the identification of variables used in each model component and a description of the assumed relationship between variables within each model component.

Adult component. Limited information is available concerning specific habitat requirements of adult striped bass in estuaries, but they are assumed to be adaptable with respect to food and aquatic cover. Many stocks of adult striped bass remain in the estuaries, while others undertake coastal migrations during the growing season. This model is not intended for evaluation of marine habitat and, in the case of some migratory stocks, the component index for adults (CIa) should be deleted.

Habitat suitability for adult striped bass is limited primarily by water quality. Adult requirements appear to be similar to those for juveniles, and the same variables and calculations used to compute the juvenile component index are used for the adult component index.

Spawning component. Striped bass have limited tolerances to water temperature and total dissolved solids during spawning. Spawning is reduced at temperatures above and below the optimal range. A total dissolved solids




Figure 2. Tree diagram illustrating the relationship of habitat variables, life requisites, life stages, and habitat types to the Habitat Suitability Index (HSI) for coastal stocks of striped bass.

(TDS) concentration over 5 ppt may block spawning. Maximum egg deposition has been observed at TDS concentrations less than 0.18 ppt, but discrepancies have been noted in the literature. Therefore, specific data for available stocks should be used to determine the applicability of this variable.

River discharge (m^3/s) influences spawning habitat suitability. Although critical values have not been established, it is assumed that 100% of the natural river discharge (average during spawning) will provide optimal spawning conditions. River discharge above the average (100% of natural) is not assumed to reduce habitat suitability. Natural river discharge may be considered to be mean discharge prior to human modifications of the system or the discharge that would occur if diversions and withdrawals ceased. Reduced river discharge is assumed to result in a proportional reduction in habitat suitability. In situations involving limitations to upstream passage due to low flows, a minimum discharge should be determined that would still allow access to spawning habitat.

It is assumed that a riverine habitat will not provide suitable conditions for spawning if the suitability index (SI) for any variable is equal to zero. Therefore, the geometric mean of the suitability indexes for these variables is used to compute the component index (CI) for spawning habitat (CIs). Using this approach to combine variables, the resulting CI will be zero if any variable SI is zero.

Egg component. Egg survival to hatching primarily depends on defined tolerances to certain water quality parameters. The temperature optima and tolerances reported for striped bass eggs closely correspond with those reported for spawning. Therefore, a single suitability index graph for water temperature is used in calculating both component indexes. Dissolved oxygen levels below 5 mg/l result in reduced egg survival. There is little chance of egg survival at dissolved oxygen levels below 3.5 mg/l. Striped bass eggs require suspension in the water column for survival. Lack of suspension results in limited hatching success, presumably due to siltation and insufficient dissolved oxygen. It is assumed that egg suspension is related to maintenance of proper water quality around the eggs. A critical value for current velocity of 30 cm/s (1 ft/s) has been documented for adequate egg suspension. However, the suitability index graph for water velocity allows a limited range for intermediate index values because the necessary suspension velocity will vary somewhat due to the precise density of the river water.

It is assumed in the calculation of the component index for eggs (CIE) that low SI values for water temperature are not as crucial as low SI values for dissolved oxygen and current velocity. When temperature has the lowest SI, the component index is calculated by the geometric mean. However, if dissolved oxygen or current velocity are limiting, it is assumed to be unlikely that even optimal temperatures will mitigate the undesirable conditions. As a result, in situations in which dissolved oxygen or current velocity variables have the lowest SI, the lowest value is considered limiting and becomes the component index.

Larval component. Larval survival appears to be primarily associated with narrow tolerances for certain physicochemical factors. Striped bass larvae can withstand a considerable range of water temperatures from 12° to

23°C (54" to 73°F), although the highest survival occurs within a narrow range from 18" to 21°C (64" to 70"). As in the egg stage, dissolved oxygen levels below 5 mg/l result in reduced survival. A critical and optimal range for salinity has also been identified. These factors are averaged geometrically to yield a subcomponent index for water quality (WQ) during larval development.

The importance of sufficient food (zooplankton) during larval development emphasizes the need to evaluate this requirement in determining habitat suitability. While it is currently impossible to easily predict zooplankton abundance at a given point in time, indirect measures can be used to reflect the potential of an estuarine habitat to provide sufficient zooplankton prey. It is assumed that potential zooplankton abundance is associated with estuarine productivity, which is believed to be a function of freshwater input during late winter and early spring and the extent of salt marsh. Optimal habitat suitability is assumed to occur when freshwater input and extent of salt marsh are in their original condition.

Reductions in both the freshwater input to the estuary during early year high flows and in the original areas of salt marsh are associated with reduced habitat suitability. Original conditions are those known to have occurred prior to human modification or as the conditions that would occur if current or planned alterations were eliminated. The late winter and spring high flow period is defined as the period in which the river hydrograph near the estuary exceeds the average annual discharge.

Although the true relationships between habitat suitability and the indirect variables of freshwater input and extent of salt marsh are not known, a sigmoid curve is assumed to be an adequate approximation. This form of the SI relationship was chosen to allow for the resilience known to exist in many ecological systems. A sigmoid curve reflects a healthy system's ability to withstand minor perturbations. More importantly, this relationship reflects a system's declining ability to absorb incremental deterioration. Both variables are averaged together to form a subcomponent index for food (F).

The component index for the larval stage of striped bass (CI1) is the geometric mean of the subcomponent indexes for water quality and food. The food subcomponent is weighted due to the emphasis placed on the importance of food availability in the recent literature on striped bass biology.

In calculating the overall habitat suitability index (HSI) for striped bass, the larval stage is given critical importance because the literature indicates that success of this stage principally determines population size. It is assumed that limitations in larval striped bass habitat suitability will not be compensated for by better conditions in other life stages.

Juvenile component. Limited information is available for specific habitat requirements of juvenile striped bass. Juvenile survival does not appear to be limited by food and aquatic cover. Therefore, these habitat variables are not included in model. However, juvenile striped bass do appear to have a limited tolerance to certain physicochemical factors associated with water quality. The optimal temperature range for growth and survival during the growing season is approximately 14° to 21°C (57" to 70°F), although water temperatures from 10" to 27°C (50" to 81°F) are tolerated. Juvenile striped


bass cannot survive dissolved oxygen levels of 1 mg/l. Intermediate survival occurs at 3 mg/l, with optimal survival at dissolved oxygen levels of 5 mg/l or more. The component index for the juvenile stage (CI_j) is the geometric mean of the dissolved oxygen and temperature suitability indexes.

Suitability Index (SI) Graphs for Model Variables


This section provides graphic representation of the relationships between various measurements of habitat variables and habitat suitability for striped bass. The variables are identified with the habitat category (R = riverine, E = estuarine) indicated. The suitability index (SI) values are read directly off the graph (1.0 = optimal habitat, 0.0 = no habitat) for any variable value.

Habitat Variable

R V₁ Percent natural river discharge during the spawning season.




R V₂ Maximum total dissolved solids (TDS) concentration during the spawning season.




Habitat Variable


R V_3 Average water temperature during the spawning season and period of egg development



R,E V_4 Minimum dissolved oxygen level during egg and larval development.




V_5 Average current velocity in water column during period of egg development.




Habitat Variable


V₆ **Percent original salt marsh in estuary.**



V₇ **Percent of original freshwater input (average volume) to estuary during the late winter and spring high flow period.**



V₈ **Average water temperature during period of larval development.**




Habitat Variable

E


V_9

Average salinity during period of larval development.



V_{10}


Average dissolved oxygen during the growing season.



E

V_{11}

Average water temperature during growing season.



Component Index Equations

To obtain component index scores for each life stage of striped bass, the SI values for appropriate variables must be combined by the use of the equations presented below.

<u>LIFE STAGE</u>	<u>EQUATION</u>
Adult	$CIa = V_{10}$ or VII, whichever is lowest
Spawning	$CI_s = (VI \times V_2 \times V_3)^{1/3}$
Egg	if $V_3 < V_4$ and V_5 , then $CI_e = (V_3 \times V_4 \times V_5)^{1/3}$ if $V_3 \geq V_4$ or V_5 , then $CI_e = V_4$ or V_5 , whichever is lowest
Larval	$CI_l = (F^2 \times WQ)^{1/3}$ where $F = (V_6 + V_7)/2$ and $WQ = (V_4 \times V_8 \times V_9)^{1/3}$
Juvenile	$CI_j = V_{10}$ or V_{11} , whichever is lowest

HSI Determination

The following steps must be taken to determine an HSI for any application.

1. Following a review of habitat use information, determine which life stages are found in the habitats (riverine, estuarine, or both) to be evaluated. Also, identify the time periods in which each life stage may be present. This step identifies the appropriate components for the application.
2. Review the section on model applicability for validity of the model for the intended application.
3. Obtain necessary data for each model variable used in the model components to be included in the application. Using the SI graphs and equations, calculate the component indexes.
4. Calculate the HSI:
 - a. If the component index for the larval stage (CI_l) is the lowest component index, then

$$HSI = CI_l;$$

- b. If the component index for the larval stage is not the lowest component index, then

$$HSI = (CI_s \times CI_e \times CI_l \times CI_j \times CI_a)^{1/5}$$

or

$$HSI = (CI_1 \times \dots \times CI_n)^{1/n}$$

if less than 5 components are used.

Four sample data sets from which suitability indices, component indices, and habitat suitability index values have been generated using the model equations are presented in Table 1. Data set 2 represents conditions where adult striped bass are absent from the estuary during the growing season. Data set 4 represents a situation where only the estuarine habitat is being evaluated.

The data sets are not actual field measurements, but represent the kinds of values that one could expect to obtain in riverine and estuarine habitats used by striped bass. The HSI's calculated from these hypothetical data sets reflect carrying capacity trends which the authors believe are appropriate for water bodies with the characteristics listed in Table 1. Thus, the models meet the acceptance goal of producing an index between 0 and 1 that is believed to have a positive relationship to a habitat's carrying capacity for striped bass.

Field Use of the Model

The level of detail needed for a particular application of this model will vary depending on time, money, and accuracy constraints. Detailed evaluation of all variables will produce the most reliable and replicable HSI values. Use of previously collected data for any or all variables may produce a satisfactory application of the model with minimal expense. Data required by this model are frequently available from published or resource agency sources. Table 2 presents suggested techniques for measuring model variables and notes references to consult for more detailed guidance.

Any or all variables can be estimated for preliminary applications of the model. Subjective estimates will decrease model reliability and replicability. When subjective estimates are used, they should be made by experienced professionals, if possible, and accompanied by full documentation of the basis on which estimates were made.

Interpreting Model Outputs

A striped bass HSI determined by field application of these models may not reflect the population density of the species in the study area, since factors other than habitat related ones may be significant in determining population size. However, in coastal areas where striped bass populations are primarily regulated by habitat-based factors, the models presented here should yield HSI's that are positively correlated with long-term average population levels. This correlation, however, has not been tested.

Table 1. Calculations of the suitability indices (SI), component indices (CI) and the habitat suitability index (HSI) for four sample data sets, using the striped bass habitat variables (V) and model equations.

Model component	Data set 1		Data set 2^a		Data set 3		Data set 4^b	
	Data	SI	Data	SI	Data	SI	Data	SI
V_1	90	0.90	75	0.75	75	0.75	-	-
V_2	1.3	1.00	2.9	0.60	1.1	1.00	-	-
V_3	16	0.75	22	0.48	18	1.00	-	-
V_4	4.5	0.70	4.5	0.70	4.2	0.50	4.6	0.75
V_5	32	1.00	30	0.50	30	0.50	-	-
V_6	85	0.95	55	0.65	50	0.50	95	0.99
V_7	85	0.95	75	0.90	50	0.50	90	0.97
V_8	19	1.00	22	0.46	22	0.46	15.5	0.60
V_9	3	1.00	12	0.37	12	0.37	4	1.00
V_{10}	4.5	0.88	3	0.50	4.2	0.80	5	1.00
V_{11}	20	1.00	16	1.00	23	0.67	23	0.67
CI _s		0.88		0.60		0.91		
CI _e		0.70		0.55		0.50		-
CI _l		0.93		0.67		0.48		0.90
CI _j		0.88		0.50		0.67		0.67
CI _a		0.88				0.67		0.67
HSI		0.85		0.58		0.48		0.74

^a Adults absent from estuary during the growing season.

^b Only estuarine habitat is evaluated.

Table 2. Description of striped bass HSI model variables and suggested techniques for measuring variables for riverine and estuarine habitats.

Habitat	Variable	Variable description	Suggested technique	Reference source
Riverine	V ₁	% of natural river discharge in the spawning area during the spawning season. ^a	Historical U.S. Geological Survey gage data or river information to estimate altered and unaltered conditions.	Stalnaker and Arnette 1976 Buchanan and Somers 1976
Riverine	V ₂	Maximum total dissolved solids (TDS) during the spawning season.	Existing data, literature, or field sampling in spawning area and laboratory analysis.	American Public Health Association 1976 U.S. Environmental Protection Agency 1979
20 Riverine	V ₃	Average water temperature during spawning and egg development.	Existing data, literature, or field sampling in areas of spawning and egg development using thermometers.	American Public Health Association 1976 Lind 1974
Riverine Estuarine	V ₄	Minimum dissolved oxygen during egg and larval development.	Existing data, literature, or field sampling in areas which contain eggs and larvae by using dissolved oxygen meters or laboratory analysis	American Public Health Association 1976 U.S. Environmental Protection Agency 1979 Lind 1974
Riverine	V ₅	Average current velocity in water column during egg development.	Existing data, literature, or field measurements in areas with eggs.	Stalnaker and Arnette 1976 Buchanan and Somers 1976
Estuarine	V ₆	% of original salt marsh in estuary. ^b	Historical maps and information or current topographic maps and data to estimate prior conditions.	Lind 1974

Table 2. (Concluded).

Habitat	Variable	Variable description	Suggested technique	Reference source
Estuarine	V ₇	% of original freshwater input to estuary during early-year high flow period. ^c	Historical U.S. Geological Survey gage data or river basin information to estimate altered and unaltered conditions.	Stalnaker and Arnette 1976
Estuarine	V ₈	Average surface water temperature during larval development.	Existing data, literature, or field measurements in areas with larvae by using thermometers.	American Public Health Association 1976 Lind 1974
Estuarine	V ₉	Average surface water salinity during larval development.	Existing data, literature or field sampling in areas with larvae by using salinometers or laboratory analysis.	American Public Health Association 1976 U.S. Environmental Protection Agency 1979
Estuarine	V ₁₀	Average dissolved oxygen during the growing season.	Existing data, literature, or field sampling of the estuary by using dissolved oxygen meters or laboratory analysis.	American Public Health Association 1976 U.S. Environmental Protection Agency 1979 Lind 1974
Estuarine	V ₁₁	Average water temperature during the growing season.	Existing data, literature or field sampling of the estuary by using thermometers.	American Public Health Association 1976 Lind 1974

^aDefined as percent of original or unaltered average discharge (m^3/s) during spawning period.

^bDefined as estimated hectares of salt marsh prior to human alteration.

^cDefined as average water volume entering the estuary during time period when river hydrograph exceeds average annual flow.

The proper interpretation of the HSI is one of comparison. If two different areas, or the same area at different points in time, have different HSI's, then the area with the higher HSI should have the potential to support more striped bass than that with the lower HSI.

ADDITIONAL HABITAT SUITABILITY INDEX MODELS

The Western Energy and Land Use Team (WELUT) of the U.S. Fish and Wildlife Service at Fort Collins, Colorado, has developed an HSI model for freshwater striped bass habitats. Striped bass stocks in certain estuarine systems undergo spawning migrations that typically extend inland, beyond the first 45 km (28 mi) of the freshwater riverine habitat. The riverine HSI model developed by WELUT may be substituted for the spawning and egg components of this model. While both HSI models are similar with respect to the riverine habitat, the inland freshwater model does include some additional variables. The estuarine striped bass HSI model is primarily intended for use on stocks which complete the spawning and egg stages close to the estuary.

LITERATURE CITED

- Albrecht, A.B. 1964. Some observations on factors associated with the survival of striped bass eggs and larvae. Calif. Fish Game 50:101-113.
- American Public Health Association. 1976. Standard Methods for the examination of water, sewage, and wastewater. American Public Health Association, Washington, D.C.
- Barkuloo, J.M. 1970. Taxonomic status and reproduction of striped bass (Morone saxatilis) in Florida. U.S. Bur. Sport Fish. Wildl. Tech. Pap. 44. 16 pp.
- Bason, W.H. 1971. Ecology and early life history of striped bass, Morone saxatilis, in the Delaware estuary. Ichthyol. Assoc. Bull. 4. 122 pp.
- Berggren, T.J., and J.T. Lieberman. 1978. Relative contribution of Hudson, Chesapeake, and Roanoke striped bass, Korone saxatilis, stocks to the Atlantic coast fishery. U.S. Natl. Mar. Fish. Serv. Bull. 76:335-345.
- Bigelow, H.B., and W.C. Schroeder. 1953. Fishes of the Gulf of Maine. U.S. Fish Wildl. Serv. Fish. Bull. 53.
- Biggs, R.B., and D.A. Flemer. 1972. The flux of particulate carbon in an estuary. Mar. Biol. (Berl.) 12:11-17.
- Boynton, W.R., T.T. Polgar, and H.H. Zion. 1981. Importance of juvenile striped bass food habits in the Potomac estuary. Trans. Am. Fish. Soc. 111:56-63.
- Buchanan, J.J., and W.P. Somers. 1976. Discharge measurements at gaging stations. U.S. Geological Survey techniques of water-resources investigations, Book 3, Chapter A8.
- Calhoun, A.J. 1953. Distribution of striped bass fry in relation to major water diversions. Calif. Fish Game 39:279-299.
- Calhoun, A.J., C.A. Woodhull, and W.C. Johnson. 1950. Striped bass reproduction in the Sacramento River system in 1948. Calif. Fish Game 36: 135-145.
- Carlson, F.T., and J.A. McCann. 1968. Hudson River fisheries investigations 1965-1968. Evaluations of a proposed pump storage project at Cornwall. New York, in relation to fish in the Hudson River. Pages 320-372 in G.P. tiowells and G.J. Lauer, eds. Proceedings of the second symposium on Hudson River Ecology. Consolidated Edison Co., New York.

- Chadwick, H.K., D.E. Stevens, and L.W. Miller. 1977. Some factors regulating the striped bass population in the Sacramento-San Joaquin estuary, California. Pages 18-35 in W. Van Winkle, ed. Proceedings of the conference on assessing the effects of power-plant-induced mortality on fish populations. Pergammon Press, New York.
- Chittenden, M.E., Jr. 1971. Status of the striped bass, Morone saxatilis, in the Delaware River. Chesapeake Sci. 12(3):131-136.
- Clark, J.R. 1968. Seasonal movements of striped bass contingents of Long Island Sound and the New York Bight. Trans. Am. Fish. Soc. 97:320-343.
- Cooper, J.C., and T.T. Polgar. 1981. Recognition of year-class dominance in striped bass management. Trans. Am. Fish. Soc. 110:180-187.
- Crateau, E.J., P.A. Moon, and C.M. Wooley. 1981. Biology, population dynamics and management of Morone sp., with emphasis on native Gulf of Mexico race and introduced Atlantic race striped bass, Apalachicola River, Florida. Apalachicola River Striped Bass Project, Annual Progress Report. U.S. Fish and Wildlife Service, Office of Fisheries Assistance, Panama City, Fla. 105 pp.
- Dahlberg, M.D. 1979. A review of survival rates of fish eggs and larvae in relation to impact assessment. Mar. Fish. Rev. 41(3):1-12.
- Davies, W.D. 1970. The effects of temperature, pH, and total dissolved solids on survival of immature striped bass, Morone saxatilis (Walbaum). Ph.D. Thesis. North Carolina State University, Raleigh, N.C. 100 pp.
- Day, J.W., Jr., T.J. Butler, and W.H. Connor. 1975. Productivity and nutrient export studies in a cypress swamp and lake system in Louisiana. Pages 255-269 in M. Wiley, ed. Estuarine processes. Volume II. Circulation, sediments, and transfer of material in the estuary. Academic Press, New York.
- deSilva, D.P. 1973. Nektonic food webs in estuaries. Pages 420-447 in L.E. Cronin, ed. Estuarine research. Volume I. Chemistry, biology, and the estuarine system. Academic Press, New York.
- Dickson, A.W. 1957. The status of striped bass (Roccus saxatilis Walbaum) in North Carolina waters. Proc. 11th Annu. Conf. Southeast. Assoc. Game Fish Comm.:264-268.
- Doroshev, S.I. 1970. Biological features of the eggs, larvae and young of the striped bass (Roccus saxatilis Walbaum) in connection with the problem of its acclimatization in the USSR. J. Ichthyol. 10(2):235-248.
- Eldridge, M.B., J.A. Whipple, D. Eng, M.J. Bowers, and B.M. Jarvis. 1981. Effects of food and feeding factors on laboratory-reared striped bass larvae. Trans. Am. Fish. Soc. 110:111-120.

- Farley, T.C. 1966. Striped bass, Roccus saxatilis, spawning in the Sacramento-San Joaquin River System, during 1963-1964. Pages 28-43 in J.L. Turner and D.W. Kelly, compilers. Ecological studies of the Sacramento-San Joaquin estuary. Part 2. Fishes of the delta. Calif. Dep. Fish Game Fish. Bull. 136.
- Fish, F.F., and E.G. McCoy. 1959. The river discharges required for an effective spawning of striped bass in the rapids of the Roanoke River of North Carolina. N.C. Wildl. Res. Comm., Raleigh. 38 pp.
- Goode, G.B., and associates. 1884. Part III. Fishes. Pages 163-683 in G.B. Goode, The fisheries and fishing industry of the United States, Section 1. National history of useful aquatic animals. U.S. Conim. Fish., Washington, D.C.
- Hardy, J.D., Jr. 1978. Development of fishes of the mid-Atlantic Bight: An atlas of the egg, larval, and juvenile stages. Vol. 3. Aphredoderidae through Rachycentridae. U.S. Fish Wildl. Serv., Biol. Serv. Program FWS/OBS-78/12. 394 pp.
- Heinle, D.R., D.A. Flemer, and J.P. Ustach. 1975. Contribution of tidal marshlands to mid-Atlantic food chains. Pages 309-320 in M. Wiley, ed. Estuarine processes. Volume II. Circulation, sediments, and transfer of material in the estuary. Academic Press, New York.
- Hester, F.E., and R.E. Stevens. 1970. Factors affecting survival of immature striped bass. Natl. Mar. Fish. Serv., NOAA-72053113. Arlington, Va. 139 pp.
- Heubach, W., R.J. Toth, and A.M. McCready. 1963. Food of young-of-the-year striped bass (Roccus saxatilis) in the Sacramento-San Joaquin River System. Calif. Fish Game 49:224-259.
- Hobbie, J.E., B.J. Copeland, and W.G. Harrison. 1973. Sources and fates of nutrients of the Pamlico River estuary, North Carolina. Pages 287-302 in L.E. Cronin, ed. Estuarine research. Volume I. Chemistry, biology, and the estuarine system. Academic Press, New York.
- Hollis, E.H. 1952. Variations in the feeding habits of the striped bass, Roccus saxatilis (Walbaum), in Chesapeake Eay. Bull. Bingham Oceanogr. Collect. Yale Univ. 14(1):111-131.
- Humphries, E.T., and K.B. Cumming. 1973. An evaluation of striped bass fingerling culture. Trans. Am. Fish. Soc. 102(1):13-20.
- Jackson, H.W., and R.E. Tiller. 1952. Preliminary observations on spawning potential in the striped bass, Roccus saxatilis (Walbaum). Md. Dep. Res. Educ. Publ. 93. 169 pp.
- Johnson, R.K., and T.S.Y. Koo. 1975. Production and distribution of striped bass (Morone saxatilis) eggs in the Chesapeake and Delaware Canal. Chesapeake Sci. 16:39-55.

- Kernehan, R.J., M.R. Headrick, and R.E. Smith. 1981. Early life-history of striped bass in the Chesapeake and Delaware Canal and vicinity. Trans. Am. Fish. Soc. 110:137-150.
- Krouse, J.S. 1968. Effects of dissolved oxygen, temperature, and salinity on survival of young striped bass, Roccus saxatilis (Walbaum). M.S. Thesis. North Carolina State University, Raleigh. 100 pp.
- Lal, K., R. Lasker, and A. Kuljis. 1977. Acclimation and rearing of striped bass larvae in sea water. Calif. Fish Game 63:210-218.
- Lewis, R.M. 1962. Sexual maturity as determined from ovum diameters in striped bass from North Carolina. Trans. Am. Fish. Soc. 91:279-282.
- Lind, G.T. 1974. Handbook of common methods in limnology. C.V. Mosby Company, St. Louis, Mo.
- Manooch, C.S., III. 1973. Food habits of yearling and adult striped bass, Morone saxatilis (Walbaum), from Albemarle Sound, North Carolina. Chesapeake Sci. 14:73-86.
- Mansueti, R.J. 1958. Eggs, larvae and young of the striped bass, Roccus saxatilis. Md. Dep. Res. Educ., Chesapeake Biol. Lab., Contrib. 112. 35 pp.
- May, R.C. 1974. Larval mortality in marine fishes and the critical period concept. Pages 3-19 in J.H.S. Blaxter, ed. The early life history of fish. Springer-Verlag, New York.
- Meldrim, J.W., and J.J. Gift. 1971. Temperature preference, avoidance and shock experiments with estuarine fishes. Ichthyol. Assoc. Bull. 7. 75 pp.
- Meldrim, J.W., J.J. Gift, and B.R. Petrosky. 1974. The effect of temperature and chemical pollutants on the behavior of several estuarine organisms. Ichthyol. Assoc. Bull. 11. 129 pp.
- Merriman, D. 1941. Studies on the striped bass (Roccus saxatilis) of the Atlantic coast. U.S. Fish Wildl. Serv. Fish. Bull. 50:1-77.
- Miller, P.E. 1977. Experimental study and modeling of striped bass egg and larval mortality. Ph.D. Thesis. The John Hopkins University, Baltimore, Md. 99 pp.
- Morgan, A.R., and A.R. Gerlach. 1950. Striped bass studies on Coos Bay in 1949 and 1950. @reg. Fish. Comm. Contrib. 14. 31 pp.
- Morgan, R.P., II, and V.J. Rasin, Jr. 1973. Effects of salinity and temperature on the development of eggs and larvae of striped bass and white perch. Appendix X in Hydrographic and ecological effects of enlargement of the Chesapeake and Delaware Canal. Final Rep. DACW-61-71-C-0062, Philadelphia Dist., U.S. Army Corps of Engineers.

- Morgan, R.P., II, V.J. Rasin, Jr., and R.L. Copp. 1981. Temperature and salinity effects on development of striped bass eggs and larvae. Trans. Am. Fish. Soc. 110:95-99.
- Murawski, W.S. 1969. The distribution of striped bass, Roccus saxatilis, eggs and larvae in the lower Delaware River. N.J. Dep. Conserv. Econ. Dev., Div. Fish and Game, Nacote Creek Res. Stn., Misc. Rep. 1M, 39 pp.
- Odum, W.E., and E.J. Heald. 1973. The detritus-based food web of an estuarine mangrove community. Pages 265-286 in L.E. Cronin, ed. Estuarine research. Volume I. Chemistry, biology, and the estuarine system. Academic Press, New York.
- Pearson, J.C. 1938. The life history of the striped bass, or rockfish, Roccus saxatilis (Walbaum). U.S. Bur. Fish. Bull. 49:825-860.
- Pearson, J.C. 1940. The young of some marine fishes taken in lower Chesapeake Bay, Virginia, with special reference to gray trout, Cynoscion regalis (Block). U.S. Fish Wildl. Serv. Fish. Bull. 50:79-102.
- Polgar, T.T., J.A. Mihursky, R.E. Ulanowicz, R.P. Morgan II, and J.S. Wilson. 1975. An analysis of 1974 striped bass spawning success in the Potomac estuary. Pages 151-165 in M. Wiley, ed. Estuarine processes. Volume I. Uses, stresses, and adaptation to the estuary. Academic Press, New York.
- Radtke, L.C., and J.L. Turner. 1967. High concentrations of total dissolved solids block spawning migration of striped bass, Roccus saxatilis, in the San Joaquin River, California. Trans. Am. Fish. Soc. 96:405-407.
- Raney, E.C. 1952. The life history of the striped bass, Roccus saxatilis (Walbaum). Bull. Bingham Oceanogr. Collect. Yale Univ. 14:5-177.
- Raney, E.C., W.S. Woolcott, and A.G. Mehring. 1954. Migratory pattern and racial structure of Atlantic coast striped bass. Trans. 19th. N. Am. Wildl. Conf. 376-396.
- Rathjen, W.F., and L.C. Miller. 1957. Aspects of the early life history of the striped bass (Roccus saxatilis) in the Hudson River. N.Y. Fish Game J. 4:43-60.
- Reimold, R.J., J.L. Gallagher, R.A. Linthurst, and W.J. Pfeiffer. 1973. Detritus production in coastal Georgia salt marshes. Pages 217-228 in L.E. Cronin, ed. Estuarine research. Volume I. Chemistry, biology, and the estuarine system. Academic Press, New York.
- Rogers, B.A., D.T. Westin, and S.B. Saila. 1977. Life stage duration studies on Hudson River striped bass. Univ. Rhode Island, Applied Mar. Res. Group, NOAA Sea Grant Mar. Tech. Rep. 31. 111 pp.
- Saila, S.B. 1973. Some aspects of fish production and cropping in estuarine systems. Pages 473-493 in L.E. Cronin, ed. Estuarine research. Volume I. Chemistry, biology, and the estuarine system. Academic Press, New York.

- Sasaki, S. 1966. Distribution of juvenile striped bass, Roccus saxatilis, in the Sacramento-San Joaquin Delta. Pages 59-67 in J.L. Turner and D.W. Kelly, compilers. Ecological studies of the Sacramento-San Joaquin Delta, Part II. Fishes of the delta. Calif. Dep. Fish Game Fish Bull. 136.
- Schubel, J.R., and A.H. Auld. 1974. Hatching success of blueback herring and striped bass eggs with various time versus temperature histories. Pages 164-190 in J.W. Gibbons and R.R. Sharitz, eds. Thermal ecology. Tech. Inf. Cent., Off. Int. Serv., U.S. AEC, Oak-Ridge, Tenn.
- Setzler, E.M., W.R. Boynton, K.V. Wood, H.H. Zion, L. Lubbers, N.K. Mountford, P. Frere, L. Tucker, and J.A. Mihursky. 1980. Synopsis of biological data on striped bass, Morone saxatilis (Walbaum). NOAA Tech. Rep., NMFS Circ. 433. 69 pp.
- Shannon, E.H. and W.B. Smith. 1967. Preliminary observations on the effect of temperature on striped bass eggs and sac fry. Proc. 21st. Annu. Conf. Southeast. Assoc. Game Fish Comm.:257-260.
- Stalnaker, C.B., and J.L. Arnette. 1976. Basic streamflow measurements and relationships. Pages 16-33 in C.B. Stalnaker and J.L. Arnette, eds. Methodologies for the determination of stream resource flow requirements: An assessment. U.S. Fish and Wildlife Service, Office of Biological Services. FWS/OBS-76/03.
- Stevens, D.E. 1966. Food habits of striped bass, Roccus saxatilis, in the Sacramento-San Joaquin Delta. Pages 68-96 in J.L. Turner and D.W. Kelly, compilers. Ecological studies of the Sacramento-San Joaquin Delta. Part II. Fishes of the delta. Calif. Dep. Fish Game Fish. Bull. 136.
- Stevens, D.E. 1979. Environmental factors affecting striped bass (Morone saxatilis) in the Sacramento-San Joaquin estuary. Pages 469-478 in T.J. Conomos, ed. San Francisco Bay: The urbanized estuary. Am. Assoc. Ad. Sci., Pacific Division, San Francisco, Calif.
- Stevenson, J.C., D.R. Heinle, D.A. Flemer, R.J. Small, R.A. Rowland, and J.F. Ustach. 1975. Nutrient exchanges between brackish water marshes and the estuary. Pages 219-240 in M. Wiley, ed. Estuarine processes. Volume II. Circulation, sediments, and transfer of material in the estuary. Academic Press, New York.
- Talbot, G.B. 1966. Estuarine environmental requirements and limiting factors for striped bass, Roccus saxatilis. Pages 37-49 in R.F. Smith, ed. A symposium on estuarine fisheries. Am. Fish. Soc. Spec. Publ. 3. 154 pp.
- Teal, J.M. 1962. Energy flow in the salt marsh ecosystem of Georgia. Ecology 43(4):614-624.
- Turner, J.L., and H.K. Chadwick. 1972. Distribution and abundance of young-of-the-year striped bass, Morone saxatilis, in relation to river flow in the Sacramento-San Joaquin estuary. Trans. Am. Fish. Soc. 101(3):442-452.

- Turner, J.L., and T.C. Farley. 1971. Effects of temperature, salinity, and dissolved oxygen on the survival of striped bass eggs and larvae. Calif. Fish Game 57:268-273.
- Ulanowicz, R.E., and T.T. Folgar. 1980. Influences of anadromous spawning behavior and optimal environmental conditions upon striped bass (Morone saxatilis) year-class success. Can. J. Fish. Aquat. Sci. 37:143-154.
- U.S. Environmental Protection Agency. 1979. Methods for chemical analysis of water and wastes. U.S. Environmental Protection Agency. Office of Research and Development. EPA-600/4-79-020.

REPORT DOCUMENTATION 1. REPORT N ^o . PAGE FWS/OBS-82/10.1		3. Recipient's Accession No.
4. Title and Subtitle Habitat Suitability Index Models: Coastal Stocks of Striped Bass		5. Report Date February 1982 6.
7. Author(s) Mark B. Bain and Jane L. Bain		8. Performing Organization Rept. No.
9. Performing Organization Name and Address U.S. Fish and Wildlife Service Massachusetts Cooperative Fishery Research Unit Department of Forestry and Wildlife Management University of Massachusetts Amherst, MA 01003		10. Project/Task/Work Unit No. 11. Contract(C) or Grant(G) No. (C) (G)
12. Sponsoring Organization Name and Address U.S. Fish and Wildlife Service Office of Biological Services National Coastal Ecosystems Team 1010 Gause Boulevard, Slidell, Louisiana 70458		13. Type of Report & Period Covered 14.
15. Supplementary Notes		
16. Abstract (Limit: 200 words) A review and synthesis of existing information were used to develop estuarine habitat models for coastal stocks of striped bass (<i>Morone saxatilis</i>). Models for five life stages are scaled to produce an index of habitat suitability between 0 (unsuitable habitat) and 1 (optimally suitable habitat) for estuarine areas of the continental United States. Habitat suitability indexes are designed for use with Habitat Evaluation Procedures previously developed by the U.S. Fish and Wildlife Service. Guidelines for coastal striped bass model applications and techniques for estimating model variables are described.		
17. Document Analysis a. Descriptors Habitability Fishes Estuaries b. Identifiers/Open-Ended Terms Habitat Habitat Suitability Index Striped bass Morone saxatilis c. COSATI Field/Group		
18. Availability Statement Unlimited		19. Security Class (This Report) Unclassified 20. Security Class (This Page) Unclassified
		21. No. of Pages vi + 29 22. Price



DEPARTMENT OF THE INTERIOR

U.S. FISH AND WILDLIFE SERVICE



As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of **our nationally** owned public lands and natural resources. This includes fostering the wisest use of our land and water resources, protecting our fish and wildlife, preserving the environmental and cultural values of our national parks and historical places, and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to assure that their development is in the best interests of all our people. The Department also has a major responsibility for American Indian reservation communities and for people who live in island territories under U.S. administration.