

(3) The heavy metals (as Pb) content cannot be more than 10 parts per million, as determined by the "Heavy Metals Test," of the "Food Chemicals Codex," 4th ed. (1996), pp. 760-761, Method II (with a 2-gram sample and 20 microgram of lead ion in the control), which is incorporated by reference in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. Copies are available from the National Academy Press, Box 285, 2101 Constitution Ave. NW., Washington, DC 20055 (Internet address <http://www.nap.edu>), or may be examined at the Dockets Management Staff (HFA-305), Food and Drug Administration, 5630 Fishers Lane, Rm. 1061, Rockville, MD 20852, 240-402-7500, between 9 a.m. and 4 p.m., Monday through Friday, or at the National Archives and Records Administration (NARA). For information on the availability of this material at NARA, call 202-741-6030, or go to: http://www.archives.gov/federal-register/code-of-federal-regulations/ibr_locations.html.

(c) The additive is used as a halogen stabilizer in brominated soybean oil at a level not to exceed 1 percent.

[60 FR 32903, June 26, 1995, as amended at 64 FR 1759, Jan. 12, 1999; 78 FR 14665, Mar. 7, 2013; 81 FR 5591, Feb. 3, 2016; 88 FR 17719, Mar. 24, 2023]

§ 172.725 Gibberellic acid and its potassium salt.

The food additives gibberellic acid and its potassium salt may be used in the malting of barley in accordance with the following prescribed conditions:

(a) The additives meet the following specifications:

(1) The gibberellic acid is produced by deep-culture fermentation of a suitable nutrient medium by a strain of *Fusarium moniliforme* or a selection of this culture.

(2) The gibberellic acid produced is of 80 percent purity or better.

(3) The empirical formula of gibberellic acid is represented by $C_{19}H_{22}O_6$.

(4) Potassium gibberellate is the potassium salt of the specified gibberellic acid.

(5) The potassium gibberellate is of 80 percent purity or better.

(6) The gibberellic acid or potassium gibberellate may be diluted with substances generally recognized as safe in foods or with salts of fatty acids conforming to § 172.863.

(b) They are used or intended for use in the malting of barley under conditions whereby the amount of either or both additives present in the malt is not in excess of 2 parts per million expressed as gibberellic acid, and the treated malt is to be used in the production of fermented malt beverages or distilled spirits only, whereby the finished distilled spirits contain none and the finished malt beverage contains not more than 0.5 part per million of gibberellic acid.

(c) To insure the safe use of the food additives the label of the package shall bear, in addition to the other information required by the Act:

(1) The name of the additive, "gibberellic acid" or "potassium gibberellate", whichever is appropriate.

(2) An accurate statement of the concentration of the additive contained in the package.

(3) Adequate use directions to provide not more than 2 parts per million of gibberellic acid in the finished malt.

(4) Adequate labeling directions to provide that the final malt is properly labeled as described in paragraph (d) of this section.

(d) To insure the safe use of the additive the label of the treated malt shall bear, in addition to the other information required by the Act, the statements:

(1) "Contains not more than 2 parts per million _____", the blank being filled in with the words "gibberellic acid" or "potassium gibberellate", whichever is appropriate; and

(2) "Brewer's malt—To be used in the production of fermented malt beverages only" or "Distiller's malt—To be used in the production of distilled spirits only", whichever is appropriate.

§ 172.730 Potassium bromate.

The food additive potassium bromate may be safely used in the malting of barley under the following prescribed conditions:

(a)(1) It is used or intended for use in the malting of barley under conditions

§ 172.735

whereby the amount of the additive present in the malt from the treatment does not exceed 75 parts per million of bromate (calculated as Br), and the treated malt is used only in the production of fermented malt beverages or distilled spirits.

(2) The total residue of inorganic bromides in fermented malt beverages, resulting from the use of the treated malt plus additional residues of inorganic bromides that may be present from uses in accordance with other regulations in this chapter promulgated under sections 408 and/or 409 of the act, does not exceed 25 parts per million of bromide (calculated as Br). No tolerance is established for bromide in distilled spirits because there is evidence that inorganic bromides do not pass over in the distillation process.

(b) To assure safe use of the additive, the label or labeling of the food additive shall bear, in addition to the other information required by the Act, the following:

(1) The name of the additive.

(2) Adequate directions for use.

(c) To assure safe use of the additive, the label or labeling of the treated malt shall bear, in addition to other information required by the Act, the statement, "Brewer's Malt—To be used in the production of fermented malt beverages only", or "Distiller's Malt—To be used in the production of distilled spirits only", whichever is the case.

§ 172.735 Glycerol ester of rosin.

Glycerol ester of wood rosin, gum rosin, or tall oil rosin may be safely used in food in accordance with the following prescribed conditions:

(a) It has an acid number of 3 to 9, a drop-softening point of 88 to 96 °C; and a color of N or paler as determined in accordance with Official Naval Stores Standards of the United States. It is purified by countercurrent steam distillation or steam stripping.

(b) It is used to adjust the density of citrus oils used in the preparation of beverages whereby the amount of the additive does not exceed 100 parts per million of the finished beverage.

[42 FR 14491, Mar. 15, 1977, as amended at 70 FR 15758, Mar. 29, 2005; 72 FR 46896, Aug. 22, 2007]

21 CFR Ch. I (4–1–23 Edition)

§ 172.736 Glycerides and polyglycides of hydrogenated vegetable oils.

The food additive glycerides and polyglycides of hydrogenated vegetable oils may be safely used in food in accordance with the following prescribed conditions:

(a) The additive is manufactured by heating a mixture of hydrogenated oils of vegetable origin and polyethylene glycol in the presence of an alkaline catalyst followed by neutralization with any acid that is approved or is generally recognized as safe for this use to yield the finished product.

(b) The additive consists of a mixture of mono-, di- and tri-glycerides and polyethylene glycol mono- and di-esters of fatty acids (polyglycides) of hydrogenated vegetable oils and meets the following specifications:

(1) Total ester content, greater than 90 percent as determined by a method entitled "Determination of Esterified Glycerides and Polyoxyethylene Glycols," approved November 16, 2001, printed by Gattefosse S.A.S., and incorporated by reference. The Director of the Office of the Federal Register approves this incorporation by reference in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. You may obtain a copy from the Office of Food Additive Safety (HFS-200), Center for Food Safety and Applied Nutrition, Food and Drug Administration, 5001 Campus Dr., College Park, MD 20740, 240-402-1200 or you may examine a copy at the Dockets Management Staff (HFA-305), Food and Drug Administration, 5630 Fishers Lane, Rm. 1061, Rockville, MD 20852, 240-402-7500, between 9 a.m. and 4 p.m., Monday through Friday, or at the National Archives and Records Administration (NARA). For information on the availability of this material at NARA, call 202-741-6030, or go to http://www.archives.gov/federal-register/code_of_federal_regulations/ibr_locations.html.

(2) Acid value, not greater than 2, and hydroxyl value, not greater than 56, as determined by the methods entitled "Acid Value," p. 1220 and "Hydroxyl Value," p. 1223, respectively, in the Food Chemicals Codex, 7th ed. (2010), which is incorporated by reference. The Director of the Office of