AUTHENTICATED
U.S. GOVERNMENT
INFORMATION
GPO

immediate container label with a statement directed to pharmacists not to remove the label or (ii) by including in the package a printed warning with instructions to pharmacists to place the warning on the container prior to dispensing.

(d) The marketing of isoproterenol inhalation preparations may be continued if all the following conditions are met:

(1) Within 30 days following the date of publication of this section in the FEDERAL REGISTER:

(i) The label and labeling of such preparations shipped within the jurisdiction of the act are in accordance with paragraphs (b) and (c) of this section.

(ii) The holder of an approved new-drug application for such preparation submits a supplement to his new-drug application to provide for appropriate labeling changes as described in paragraphs (b) and (c) of this section.

(2) Within 90 days following the date of publication of this section in the FEDERAL REGISTER, the manufacturer, packer, or distributor of any drug containing isoproterenol intended for inhalation for which a new-drug approval is not in effect submits a new-drug application containing satisfactory information of the kinds required by § 314.50 of this chapter, including appropriate labeling as described in paragraphs (b) and (c) of this section.

(3) The applicant submits additional information required for the approval of the application as may be specified in a written communication from the Food and Drug Administration.

(e) After 270 days following expiration of said 90 days, regulatory proceedings based on section 505(a) of the Federal Food, Drug, and Cosmetic Act may be initiated with regard to any such drug shipped within the jurisdiction of the act for which an approved new-drug application is not in effect.

[40 FR 13998, Mar. 27, 1975, as amended at 55 FR 11576, Mar. 29, 1990]

§ 201.306 Potassium salt preparations intended for oral ingestion by man.

(a) The Food and Drug Administration will initiate no regulatory action with respect to the continued marketing of coated tablets containing potassium chloride or other potassium salts which supply 100 milligrams or more of potassium per tablet provided all the following conditions are met:

(1) Within 30 days from the date of publication of this statement of policy in the FEDERAL REGISTER:

(i) The labeling of the drug bears the prescription caution statement quoted in section 503(b)(4) of the Federal Food, Drug, and Cosmetic Act;

(ii) The labeling on or within the package from which the drug is to be dispensed bears adequate information for its use by practitioners in accord with the ''full disclosure'' labeling requirements of § 201.100 of this chapter, including the following warning statement:

Warning—There have been several reports, published and unpublished, concerning nonspecific small-bowel lesions consisting of stenosis, with or without ulceration, associated with the administration of enteric-coated thiazides with potassium salts. These lesions may occur with enteric-coated potassium tablets alone or when they are used with nonenteric-coated thiazides, or certain other oral diuretics. These small-bowel lesions have caused obstruction, hemorrhage, and perforation. Surgery was frequently required and deaths have occurred. Based on a large survey of physicians and hospitals, both United States and foreign, the incidence of these lesions is low, and a causal relationship in man has not been definitely established. Available information tends to implicate enteric-coated potassium salts, although lesions of this type also occur spontaneously. Therefore, coated potassium-containing formulations should be administered only when indicated, and should be discontinued immediately if abdominal pain, distention, nausea, vomiting, or gastrointestinal bleeding occur. Coated potassium tablets should be used only when adequate dietary supplementation is not practicable.

(Although the warning statement includes references to enteric-coated potassium salt preparations, it applies to any capsule or coated tablet of a potassium salt intended for oral ingestion without prior dilution with an adequate volume of liquid to preclude gastrointestinal injury.)

(iii) Any other labeling or additional advertising for the drug conforms to the labeling described in paragraph (a)(1)(ii) of this section, in accordance with §§ 202.1 and 201.100 of this chapter.

(2) Within 90 days from the date of publication of this statement of policy in the FEDERAL REGISTER, the manufacturer, packer, or distributor of the drug shall submit a new-drug application containing satisfactory information of the kind required by § 314.50 of this chapter, with appropriate labeling as described in this paragraph.

(b) The Food and Drug Administration may initiate regulatory proceedings after 30 days from the date of publication of this section, with respect to the marketing of uncoated tablets containing potassium chloride or other potassium salts which supply 100 milligrams or more of potassium per tablet or with respect to liquid preparations containing potassium chloride or other potassium salts which supply 20 milligrams or more of potassium per milliliter, labeled or intended for human use, unless all the following conditions are met:

(1) The labeling of the drug bears the prescription statement quoted in section 503(b)(4) of the Federal Food, Drug, and Cosmetic Act; and

(2) The labeling on or within the package from which the drug is to be dispensed bears adequate information for its use by practitioners in accord with the ''full disclosure'' labeling requirements of § 201.100 of this chapter, including a recommendation that patients be directed to dissolve any such tablets in an appropriate amount of liquid and to dilute any such liquid preparations adequately to assure against gastrointestinal injury associated with the oral ingestion of concentrated potassium salt preparations.

[40 FR 13998, Mar. 27, 1975, as amended at 55 FR 11576, Mar. 29, 1990; 67 FR 4906, Feb. 1, 2002]

§ 201.307 Sodium phosphates; package size limitation, warnings, and directions for over-the-counter sale.

(a) Reports in the medical literature and data accumulated by the Food and Drug Administration indicate that multiple container sizes of sodium phosphates oral solution available in the marketplace have caused consumer confusion and appear to have been involved in several consumer deaths. Sodium phosphates oral solution has been marketed in 45-milliliter (mL), 90-mL, and 240-mL container sizes. The 45-mL and 90-mL container sizes of sodium phosphates oral solution are often recommended and prescribed by physicians for bowel cleansing prior to surgery and diagnostic procedures of the colon. Sodium phosphates oral solution (adult dose 20 mL to 45 mL) is also used as an over-the-counter (OTC) laxative for the relief of occasional constipation. Accidental overdosing and deaths have occurred because the 240-mL container was mistakenly used instead of the 45-mL or 90-mL container. The Food and Drug Administration is limiting the amount of sodium phosphates oral solution to not more than 90 mL (3 ounces (oz)) per OTC container because of the serious health risks associated with the ingestion of larger than intended doses of this product. Further, because an overdose of either oral or rectal enema sodium phosphates can cause an electrolyte imbalance, additional warning and direction statements are required for the safe use of any OTC laxative drug product containing sodium phosphates.

(b) Any OTC drug product for laxative or bowel cleansing use containing sodium phosphates as an active ingredient when marketed as described in paragraph (a) of this section is misbranded within the meaning of section 502 of the Federal Food, Drug, and Cosmetic Act unless packaged and labeled as follows:

(1) Package size limitation for sodium phosphates oral solution: Container shall not contain more than 90 mL (3 oz).

(2) Warnings. The following sentences shall appear in boldface type as the first statement under the heading ''Warnings.''

(i) Oral dosage forms. ''Taking more than the recommended dose in 24 hours can be harmful.''

(ii) Rectal enema dosage forms. ''Using more than one enema in 24 hours can be harmful.''

(3) Directions—(i) The labeling of all orally or rectally administered OTC drug products containing sodium phosphates shall contain the following directions in boldface type immediately preceding the dosage information: ''Do not'' (''take'' or ''use'')