

“more unless directed by a doctor. See Warnings.”

(ii) For products containing dibasic sodium phosphate/monobasic sodium phosphate identified in § 334.16(d) marketed as a solution. Adults and children 12 years of age and over: Oral dosage is dibasic sodium phosphate 3.42 to 7.56 grams (g) and monobasic sodium phosphate 9.1 to 20.2 g (20 to 45 mL dibasic sodium phosphate/monobasic sodium phosphate oral solution) as a single daily dose. “Do not take more than 45 mL (9 teaspoonfuls or 3 tablespoonfuls) in a 24-hour period.” Children 10 and 11 years of age: Oral dosage is dibasic sodium phosphate 1.71 to 3.78 g and monobasic sodium phosphate 4.5 to 10.1 g (10 to 20 mL dibasic sodium phosphate/monobasic sodium phosphate oral solution) as a single daily dose. “Do not take more than 20 mL (4 teaspoonfuls) in a 24-hour period.” Children 5 to 9 years of age: Oral dosage is dibasic sodium phosphate 0.86 to 1.89 g and monobasic sodium phosphate 2.2 to 5.05 g (5 to 10 mL dibasic sodium phosphate/monobasic sodium phosphate oral solution) as a single daily dose. “Do not take more than 10 mL (2 teaspoonfuls) in a 24-hour period.” Children under 5 years of age: ask a doctor.

(c) After June 22, 1998, for package size limitation and September 18, 1998, for labeling in accord with paragraph (b) of this section, any such OTC drug product initially introduced or initially delivered for introduction into interstate commerce, or any such drug product that is repackaged or relabeled after these dates regardless of the date the product was manufactured, initially introduced, or initially delivered for introduction into interstate commerce, that is not in compliance with this section is subject to regulatory action.

[63 FR 27843, May 21, 1998]

§ 201.308 Ipecac syrup; warnings and directions for use for over-the-counter sale.

(a) It is estimated that each year about 500,000 accidental poisonings occur in the United States and result in approximately 1,500 deaths, of which over 400 are children. In the emergency treatment of these poisonings, ipecac syrup is considered the emetic of

choice. The immediate availability of this drug for use in such situations is critical, since rapid treatment may be the difference between life and death. The restriction of this drug to prescription sale limits its availability in emergencies. On the other hand, it is the consensus of informed medical opinion that ipecac syrup should be used only under medical supervision in the emergency treatment of poisonings. In view of these facts, the question of whether ipecac syrup labeled as an emergency treatment for use in poisonings should be available over the counter has been controversial.

(b) In connection with its study of this problem, the Food and Drug Administration has obtained the views of medical authorities. It is the unanimous recommendation of the American Academy of Pediatrics, the American Association of Poison Control Centers, the American Medical Association, and the Medical Advisory Board of the Food and Drug Administration that ipecac syrup in 1 fluid ounce containers be permitted to be sold without prescription so that it will be readily available in the household for emergency treatment of poisonings, under medical supervision, and that the drug be appropriately packaged and labeled for this purpose.

(c) In view of the above recommendations, the Commissioner of Food and Drugs has determined that it is in the interest of the public health for ipecac syrup to be available for sale without prescription, provided that it is packaged in a quantity of 1 fluid ounce (30 milliliters), and its label bears, in addition to other required label information, the following, in a prominent and conspicuous manner:

(1) A statement conspicuously boxed and in red letters, to the effect: “For emergency use to cause vomiting in poisoning. Before using, call physician, the Poison Control Center, or hospital emergency room immediately for advice.”

(2) A warning to the effect: “Warning—Keep out of reach of children. Do not use in unconscious persons. Ordinarily, this drug should not be used if strychnine, corrosives such as alkalies (lye) and strong acids, or petroleum

§ 201.309

distillates such as kerosine, gasoline, coal oil, fuel oil, paint thinner, or cleaning fluid have been ingested.”

(3) Usual dosage: 1 tablespoon (15 milliliters) in persons over 1 year of age.

§ 201.309 Acetophenetidin (phenacetin)-containing preparations; necessary warning statement.

(a) In 1961, the Food and Drug Administration, pursuant to its statutory responsibility for the safety and effectiveness of drugs shipped in interstate commerce, began an active investigation of reports of possible toxic effects and renal damage due to misuse of the drug acetophenetidin. This study led to the decision that there was probable cause to conclude that misuse and prolonged use of the drug were in fact responsible for kidney lesions and disease. The Commissioner of Food and Drugs, in December 1963, appointed an ad hoc Advisory Committee of Inquiry on Possible Nephrotoxicity Associated With the Abuse of Acetophenetidin (Phenacetin)-Containing Preparations. This committee, composed of scientists in the fields of pharmacology and medicine, on April 23, 1964, submitted its findings and conclusions in the matter and recommended that all acetophenetidin (phenacetin)-containing preparations bear a warning as provided in section 502(f)(2) of the Federal Food, Drug, and Cosmetic Act.

(b) On the basis of the studies made by the Food and Drug Administration and the report of the Advisory Committee, the Commissioner of Food and Drugs has concluded that it is necessary for the protection of users that the label and labeling of all acetophenetidin (phenacetin)-containing preparations bear a warning statement to the following effect: “Warning—This medication may damage the kidneys when used in large amounts or for a long period of time. Do not take more than the recommended dosage, nor take regularly for longer than 10 days without consulting your physician.”

§ 201.310 Phenindione; labeling of drug preparations intended for use by man.

(a) Reports in the medical literature and data accumulated by the Food and Drug Administration indicate that

21 CFR Ch. I (4–1–24 Edition)

phenindione, a synthetic anticoagulant drug, has caused a number of cases of agranulocytosis (with two fatalities). There are also reports implicating the drug in cases of hepatitis and hypersensitivity reactions. In view of the potentially serious effects found to be associated with preparations of this drug intended for use by man, the Commissioner of Food and Drugs will regard such preparations as misbranded within the meaning of section 502(f) (1) and (2) of the Federal Food, Drug, and Cosmetic Act, unless the label and labeling on or within the package from which the drug is to be dispensed, and any other labeling furnishing or purporting to furnish information for use of the drug, bear a conspicuous warning statement to the following effect: “Warning: Agranulocytosis and hepatitis have been associated with the use of phenindione. Patients should be instructed to report promptly prodromal symptoms such as marked fatigue, chill, fever, and sore throat. Periodic blood studies and liver function tests should be performed. Use of the drug should be discontinued if leukopenia occurs or if evidence of hypersensitivity, such as dermatitis or fever, appears.”

(b) Regulatory action may be initiated with respect to preparations of phenindione intended for use by man found within the jurisdiction of the act on or after November 25, 1961, unless such preparations are labeled in accordance with paragraph (a) of this section.

§ 201.311 [Reserved]

§ 201.312 Magnesium sulfate heptahydrate; label declaration on drug products.

Magnesium sulfate heptahydrate should be listed on the label of a drug product as epsom salt, which is its common or usual name.

§ 201.313 Estradiol labeling.

The article presently recognized in The National Formulary under the heading “Estradiol” and which is said to be “17-cis-beta estradiol” is the same substance formerly recognized in the United States Pharmacopeia under the designation “Alpha Estradiol.” The