

§ 201.315

with such directions for use conspicuous warning statements to the effect: "Caution: If pain persists for more than 10 days, or redness is present, or in conditions affecting children under 12 years of age, consult a physician immediately." The salicylate dosage should not exceed 60 grains in a 24-hour period or 10 grains in a 4-hour period. If the article contains other analgesics, the salicylate dosage should be appropriately reduced.

(g)(1) The label of any drug containing more than 5 percent methyl salicylate (wintergreen oil) should bear a conspicuous warning such as: "Do not use otherwise than as directed." These drug products must also include the "Keep out of reach of children" warning and the accidental ingestion warning as required in § 330.1(g) of this chapter.

(2) If the preparation is a counter-irritant or rubefacient, it should also bear a caution such as, "Caution: Discontinue use if excessive irritation of the skin develops. Avoid getting into the eyes or on mucous membranes." (See also § 201.303.)

(h)(1) The labeling of orally or rectally administered over-the-counter drug products containing aspirin or nonaspirin salicylates as active ingredients subject to this paragraph is required to prominently bear the following warning: "Reye's syndrome [subheading in bold type]: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness."

(2) This warning statement shall appear on the immediate container labeling. In cases where the immediate container is not the retail package, the retail package also must bear the warning statement. In addition, the warning statement shall appear on any labeling that contains warnings and, in such cases, the warning statement shall be the first warning statement under the heading "Warnings."

(3) Over-the-counter drug products subject to this paragraph and labeled solely for use by children (pediatric

21 CFR Ch. I (4-1-24 Edition)

products) shall not recommend the product for use in treating flu or chicken pox.

(4) Any product subject to paragraphs (h)(1), (h)(2), and (h)(3) of this section that is not labeled as required by these paragraphs and that is initially introduced or initially delivered for introduction into interstate commerce after the following dates is misbranded under sections 201(n) and 502(a) and (f) of the Federal Food, Drug, and Cosmetic Act.

(i) Compliance by October 18, 2004, for OTC drug products containing aspirin and nonaspirin salicylates as an active ingredient and marketed under a new drug application or abbreviated new drug application.

(ii) Compliance by April 19, 2004, for OTC antidiarrheal and overindulgence drug products that contain bismuth subsalicylate as an active ingredient and have annual sales greater than \$25,000.

(iii) Compliance by April 18, 2005, for OTC antidiarrheal and overindulgence drug products that contain bismuth subsalicylate as an active ingredient and have annual sales less than \$25,000.

(iv) Compliance dates for all other OTC drug products containing aspirin and nonaspirin salicylates as an active ingredient and marketed under an OTC drug monograph (for internal analgesic, antipyretic, and antirheumatic drug products, or for menstrual drug products) will be established when the final monographs for those products are published in a future issue of the FEDERAL REGISTER. In the interim, these products should continue to be labeled with the previous Reye's syndrome warning that appears in paragraph (h)(1) of this section.

[40 FR 13998, Mar. 27, 1985, as amended at 51 FR 8182, Mar. 7, 1986; 53 FR 21637, June 9, 1988; 53 FR 24830, June 30, 1988; 64 FR 13291, Mar. 17, 1999; 65 FR 8, Jan. 3, 2000; 68 FR 18869, Apr. 17, 2003]

§ 201.315 Over-the-counter drugs for minor sore throats; suggested warning.

The Food and Drug Administration has studied the problem of the labeling of lozenges or troches containing a local anesthetic, chewing gum containing aspirin, various mouth washes

and gargles and other articles sold over the counter for the relief of minor irritations of the mouth or throat. It will not object to the labeling of suitable articles of this type "For the temporary relief of minor sore throats", provided this is immediately followed in the labeling with a warning statement in prominent type essentially as follows: "Warning—Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult physician promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by physician."

§ 201.316 Drugs with thyroid hormone activity for human use; required warning.

(a) Drugs with thyroid hormone activity have been promoted for, and continue to be dispensed and prescribed for, use in the treatment of obesity, although their safety and effectiveness for that use have never been established.

(b) Drugs for human use with thyroid hormone activity are misbranded within the meaning of section 502 of the Federal Food, Drug, and Cosmetic Act unless their labeling bears the following boxed warning at the beginning of the "Warnings" section:

Drugs with thyroid hormone activity, alone or together with other therapeutic agents, have been used for the treatment of obesity. In euthyroid patients, doses within the range of daily hormonal requirements are ineffective for weight reduction. Larger doses may produce serious or even life-threatening manifestations of toxicity, particularly when given in association with sympathomimetic amines such as those used for their anorectic effects.

[43 FR 22009, May 23, 1978]

§ 201.317 Digitalis and related cardiotonic drugs for human use in oral dosage forms; required warning.

(a) Digitalis and related cardiotonic drugs for human use in oral dosage forms have been promoted for, and continue to be dispensed and prescribed

for, use in the treatment of obesity, although their safety and effectiveness for that use have never been established.

(b) Digitalis and related cardiotonic drugs for human use in oral dosage forms are misbranded within the meaning of section 502 of the Federal Food, Drug, and Cosmetic Act unless their labeling bears the following boxed warning at the beginning of the "Warnings" section:

Digitalis alone or with other drugs has been used in the treatment of obesity. This use of digoxin or other digitalis glycosides is unwarranted. Moreover, since they may cause potentially fatal arrhythmias or other adverse effects, the use of these drugs in the treatment of obesity is dangerous.

(c) This section does not apply to digoxin products for oral use.

[43 FR 22009, May 23, 1978, as amended at 85 FR 72907, Nov. 16, 2020]

§ 201.319 Water-soluble gums, hydrophilic gums, and hydrophilic mucilloids (including, but not limited to agar, alginic acid, calcium polycarbophil, carboxymethylcellulose sodium, carrageenan, chondrus, glucomannan ((B-1,4 linked polymannose acetate), guar gum, karaya gum, kelp, methylcellulose, plantago seed (psyllium), polycarbophil tragacanth, and xanthan gum) as active ingredients; required warnings and directions.

(a) Reports in the medical literature and data accumulated by the Food and Drug Administration indicate that esophageal obstruction and asphyxiation have been associated with the ingestion of water-soluble gums, hydrophilic gums, and hydrophilic mucilloids including, but not limited to, agar, alginic acid, calcium polycarbophil, carboxymethylcellulose sodium, carrageenan, chondrus, glucomannan ((B-1,4 linked) polymannose acetate), guar gum, karaya gum, kelp, methylcellulose, plantago seed (psyllium), polycarbophil, tragacanth, and xanthan gum. Esophageal obstruction and asphyxiation due to orally-administered drug products containing water-