

§ 369.21

21 CFR Ch. I (4–1–23 Edition)

SALTS). (See also § 201.314 of this chapter.)

“Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away;” or “Keep out of reach of children.”

If the article is an aspirin preparation, it should bear the first of the above two warning statements. In either case, the above information should appear on the label.

Caution—For children under 3 years of age, consult your physician; or

Caution—For younger children, consult your physician.

One of the two immediately preceding caution statements is required on the label of all aspirin tablets, but such a statement is not required on the labels of other salicylates clearly offered for administration to adults only.

If offered for use in arthritis or rheumatism, in juxtaposition therewith, the statement:

Caution—If pain persists for more than 10 days, or redness is present, or in conditions affecting children under 12 years of age, consult a physician immediately.

SALICYLATES: METHYL SALICYLATE (WINTERGREEN OIL). (See also §§ 201.303 and 201.314 of this chapter.)

“Do not use otherwise than as directed. Keep out of reach of children to avoid accidental poisoning. If swallowed, get medical help or contact a Poison Control Center right away.”

If the preparation is a counter-irritant or rubefacient the statement:

Caution—Discontinue use if excessive irritation of the skin develops. Avoid getting into the eyes or on mucous membranes.

If offered for use in arthritis or rheumatism, in juxtaposition therewith, the statement:

Caution—If pain persists for more than 10 days, or redness is present, or in conditions affecting children under 12 years of age consult a physician immediately.

SILVER.

Caution—Frequent or prolonged use of this preparation may result in permanent discoloration of skin and mucous membranes.

SODIUM PERBORATE MOUTHWASH AND GARGLE AND TOOTHPASTE.

Caution—Discontinue use if irritation or inflammation develops, or increases. Avoid swallowing.

SULFONAMIDE NOSE DROPS.

Caution—Do not use if a known allergy to sulfonamide drugs exists.

SULFUR PREPARATION FOR EXTERNAL USE.

Caution—If undue skin irritation develops or increases, discontinue use and consult physician.

THROAT PREPARATIONS FOR TEMPORARY RELIEF OF MINOR SORE THROAT: LOZENGES, TROCHES, WASHES, GARGLES, ETC. (See also § 201.315 of this chapter.)

Warning—Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult physician promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by physician.

TOOTHACHE PREPARATIONS.

For temporary use only until a dentist can be consulted.

ZINC STEARATE DUSTING POWDERS.

“Keep out of reach of children; avoid inhaling. If swallowed, get medical help or contact a Poison Control Center right away.”

[39 FR 11745, Mar. 29, 1974, as amended at 40 FR 8917, Mar. 3, 1975; 40 FR 13496, Mar. 27, 1975; 41 FR 10885, Mar. 15, 1976; 51 FR 27760, Aug. 1, 1986; 51 FR 35340, Oct. 2, 1986; 52 FR 15893, Apr. 30, 1987; 52 FR 30057, Aug. 12, 1987; 52 FR 47324, Dec. 11, 1987; 53 FR 7093, Mar. 4, 1988; 55 FR 31783, Aug. 3, 1990; 57 FR 58376, Dec. 9, 1992; 59 FR 43412, Aug. 23, 1994; 64 FR 13296, Mar. 17, 1999; 68 FR 18882, Apr. 17, 2003; 68 FR 34293, June 9, 2003]

§ 369.21 Drugs; warning and caution statements required by regulations.

ACETAMINOPHEN (N-ACETYL-*p*-AMINOPHENOL) (See § 310.201(a)(1) of this chapter.)

Warning—Do not give to children under 3 years of age or use for more than 10 days unless directed by a physician.

If offered for use in arthritis, or rheumatism, in juxtaposition therewith, the statement:

Caution—If pain persists for more than 10 days, or redness is present, or in conditions affecting children under 12 years of age consult a physician immediately.

ALCOHOL RUBBING COMPOUND. (See 26 CFR 182.855(a)(5); The National Formulary, Tenth Edition 1955, pp. 27–28; and section 502(g) of the act).

Warning—For external use only. If taken internally serious gastric disturbances will result.

ANTI-HISTAMINICS, ORAL (PHENYL-TOLOXAMINE DIHYDROGEN CITRATE AND CHLOROTHEN CITRATE PREPARATIONS). (See § 310.201(a)(4) and (a)(24) of this chapter.)

Caution—This preparation may cause drowsiness. Do not drive or operate machinery while taking this medication. Do not give to children under 6 years of age or exceed the recommended dosage unless directed by physician.

If offered for symptoms of colds, the statement:

Caution—If relief does not occur within 3 days, discontinue use and consult physician.

DICYCLOMINE HYDROCHLORIDE WITH AN ANTACID. (See § 310.201(a)(8) of this chapter.)

Warning—Do not exceed the recommended dosage. Do not administer to children under 12 years of age or use for a prolonged period unless directed by physician, since persistent or recurring symptoms may indicate a serious disease requiring medical attention.

DIPHEMANIL METHYLSULFATE FOR EXTERNAL USE. (See § 310.201(a)(22) of this chapter.)

Caution—If redness, irritation, swelling, or pain persists or increases, discontinue use and consult physician.

DRUGS IN DISPENSERS PRESSURIZED BY GASEOUS PROPELLANTS. (See also § 310.201(a) (11) and (18) of this chapter.)

The warnings herein shall appear prominently and conspicuously, but in no case may the letters be less than $\frac{1}{16}$ inch in height.

If the label of any package is too small to accommodate the warnings, the Commissioner may establish by regulation an acceptable alternative method, e.g., a type size smaller than $\frac{1}{16}$ inch in height. A petition requesting such a regulation, as an amendment to this paragraph, shall be submitted to the Division of Dockets Management in the form established in part 10 of this chapter.

Warning—Avoid spraying in eyes. Contents under pressure. Do not puncture or incinerate. Do not store at temperature above 120 °F. Keep out of reach of children.

In the case of products packaged in glass containers, the word “break” may be substituted for the word “puncture.”

The words “Avoid spraying in eyes” may be deleted from the warning in the case of a product not expelled as a spray, or that is intended to be used in the eyes.

In addition to the above warning, the label of a drug packaged in a self-purized container in which the propellant consists in whole or in part of a halocarbon or hydrocarbon shall bear the following warning:

Warning—Use only as directed. Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.

The warning is not required for the following products:

(a) Products expelled in the form of a foam or cream, which contain less than ten percent propellant in the container;

(b) Products in a container with a physical barrier that prevents escape of the propellant at the time of use;

(c) Products of a net quantity of contents of less than 2 ozs. that are designed to release a measured amount of product with each valve actuation;

(d) Products of a net quantity of contents of less than $\frac{1}{2}$ oz.

DYCLONINE HYDROCHLORIDE. (See § 310.201(a)(23) of this chapter.)

Caution—Do not use in the eyes. Not for prolonged use. Do not apply to large areas of the body. If redness, irritation, swelling, or pain persists or increases, discontinue use unless directed by physician. Do not use, but consult physician for deep or puncture wounds

§ 369.21

21 CFR Ch. I (4–1–23 Edition)

or serious burns. Do not use in case of rectal bleeding, as this may indicate serious disease.

HEXADENOL. (See § 310.201(a)(11) of this chapter.)

Caution—Do not use for treatment of serious burns or skin conditions or for conditions which persist for prolonged periods. In such cases, consult your physician. Do not spray in vicinity of eyes, mouth, nose, or ears. Do not store above 120 °F.

IPECAC SYRUP IN ONE-FLUID OUNCE CONTAINERS FOR EMERGENCY TREATMENT OF POISONING, TO INDUCE VOMITING. (See § 201.308 of this chapter.)

Ipecac syrup packaged for over-the-counter sale must bear statements to the following effect, in a prominent and conspicuous manner:

The following statement (boxed and in red letters):

“For emergency use to cause vomiting in poisoning. Before using, call physician, the Poison Control Center, or hospital emergency room immediately for advice.”

The following warning: *Warning*—Keep out of reach of children. Do not use in unconscious persons. Ordinarily, this drug should not be used if strychnine, corrosives such as alkalis (lye) and strong acids, or petroleum distillates such as kerosene, gasoline, coal oil, fuel oil, paint thinner, or cleaning fluid have been ingested.

ISOAMYLHYDROCUPREINE AND ZOLAMINE HYDROCHLORIDE RECTAL PREPARATIONS FOR EXTERNAL USE (See § 310.201(a)(3) of this chapter.)

Warning—Do not use this preparation in case of rectal bleeding, as this may indicate serious disease.

NEOMYCIN SULFATE WITH A VASOCONSTRICTOR, IN NASAL PREPARATIONS (SPRAY OR DROPS).

Caution—Do not exceed recommended dosage. Do not administer to children under 3 years of age unless directed by physician.

PRAMOXINE HYDROCHLORIDE FOR EXTERNAL USE. (See § 310.201(a)(19) of this chapter.)

Caution—Do not use in the eyes or nose. Not for prolonged use. Do not

apply to large areas of the body. If redness, irritation, swelling, or pain persists or increases, discontinue use unless directed by a physician.

SODIUM GENTISATE. (See §§ 201.314 and 310.301(a)(2) of this chapter.)

Warning—Do not use in children under 6 years of age or use for prolonged period unless directed by physician.

“Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.”

If offered for use in arthritis or rheumatism, in juxtaposition therewith, the statement:

Caution—If pain persists for more than 10 days, or redness is present, or in conditions affecting children under 12 years of age, consult a physician immediately.

TUAMINOHEPTANE SULFATE NASAL PREPARATIONS. (See § 310.201(a)(16) of this chapter.)

Caution—Do not exceed recommended dosage. Overdosage may cause nervousness, restlessness, or sleeplessness. Individuals with high blood pressure, heart disease, diabetes, or thyroid disease should use only as directed by physician. Do not use for more than 3 or 4 consecutive days unless directed by physician.

VIBESATE PREPARATIONS. (See § 310.201(a)(18) of this chapter.)

Caution—Do not use but consult physician for deep or puncture wounds or serious burns. If redness, irritation, swelling, or pain persists or increases, discontinue use and consult physician.

Warning—Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130 °Fahrenheit may cause bursting. Never throw container into fire or incinerator.

[39 FR 11745, Mar. 29, 1974]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 369.21, see the List of CFR Sections Affected, which appears in the Finding Aids section of the printed volume and at www.govinfo.gov.

PARTS 370–499 [RESERVED]