

§ 520.1638

horses. Not for use in horses intended for food. Do not administer to mares during the last month of pregnancy. Trichlorfon is a cholinesterase inhibitor. Do not use this product in animals simultaneously with, or within a few days before or after treatment with or exposure to, cholinesterase-inhibiting drugs, pesticides, or chemicals. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[50 FR 50291, Dec. 10, 1985, as amended at 61 FR 5506, Feb. 13, 1996; 78 FR 28825, May 20, 2014]

§ 520.1638 Oxibendazole.

(a) *Specifications.* (1) Each gram of paste contains 227 milligrams (mg) (22.7 percent) oxibendazole.

(2) Each milliliter of suspension contains 100 mg (10 percent) oxibendazole.

(b) *Sponsor.* See No. 054771 in § 510.600(c) of this chapter.

(c) *Special considerations.* (1) See § 500.25 of this chapter.

(2) Suspension product described in paragraph (a)(2) of this section shall be labeled: "Federal law restricts this drug to use by or on the order of a licensed veterinarian."

(d) *Conditions of use in horses—*(1) *Amount.* For uses other than for threadworms (*Strongyloides westeri*), 10 mg oxibendazole per kilogram (kg) body weight; for threadworms (*Strongyloides westeri*), 15 mg/kg. Horses maintained on premises where reinfection is likely to occur should be re-treated in 6 to 8 weeks. Administer suspension product by stomach tube in 3 to 4 pints of warm water, or by top dressing or mixing into a portion of the normal grain ration.

(2) *Indications for use.* For removal and control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*); small strongyles (genera *Cylicostephanus*, *Cylicocyclus*, *Cyathostomum*, *Triodontophorus*, *Cylicodontophorus*, and *Gyalocephalus*); large roundworms (*Parascaris equorum*); pinworms (*Oxyuris equi*) including various larval stages; and threadworms (*Strongyloides westeri*).

(3) *Limitations.* Do not use in horses intended for human consumption.

[78 FR 28825, May 20, 2014]

21 CFR Ch. I (4–1–23 Edition)

§ 520.1660 Oxytetracycline.

§ 520.1660a Oxytetracycline and carbomycin.

(a) *Specifications.* (1) Oxytetracycline: The antibiotic substance produced by growth of *Streptomyces rimosus* or the same antibiotic substance produced by any other means.

(2) Carbomycin: The antibiotic substance produced by growth of *Streptomyces halstedii* or the same antibiotic substance produced by any other means.

(b) *Sponsor.* See No. 054771 in § 510.600(c) of this chapter.

(c) *Special considerations.* The quantities of oxytetracycline in paragraph (e) of this section refer to the activity of oxytetracycline hydrochloride and the quantities of carbomycin listed refer to the activity of an appropriate standard.

(d) *Related tolerances.* See §§ 556.110 and 556.500 of this chapter.

(e) *Conditions of use.* It is used as oxytetracycline hydrochloride plus carbomycin base in drinking water of chickens as follows:

(1) *Amount.* Administer 1.0 gram of oxytetracycline and 1.0 gram carbomycin per gallon for not more than 5 days.

(2) *Indications for use.* As an aid in the prevention and treatment of complicated chronic respiratory disease (air-sac infection) caused by *Mycoplasma gallisepticum* and secondary bacterial organisms associated with chronic respiratory disease such as *E. coli*.

(3) *Limitations.* Not for use in chickens producing eggs for human consumption. Withdraw 24 hours before slaughter. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 78 FR 28825, May 20, 2014; 81 FR 94989, Dec. 27, 2016]

§ 520.1660b Oxytetracycline hydrochloride capsules.

(a) *Specifications.* The drug is in capsule form with each capsule containing 125 or 250 milligrams of oxytetracycline hydrochloride. Oxytetracycline is the antibiotic substance produced by growth of *Streptomyces rimosus* or the