

(4) *Honey bees*—(i) *Amount*. Mix 200 milligrams tylosin in 20 grams confectioners'/powdered sugar. Use immediately. Apply (dust) this mixture over the top bars of the brood chamber once weekly for 3 weeks.

(ii) *Indications for use*. For the control of American foulbrood (*Paenibacillus larvae*).

(iii) *Limitations*. The drug should be fed early in the spring or fall and consumed by the bees before the main honey flow begins, to avoid contamination of production honey. Complete treatments at least 4 weeks before main honey flow.

[40 FR 13838, Mar. 27, 1975, as amended at 50 FR 49841, Dec. 5, 1985; 59 FR 14365, Mar. 28, 1994; 62 FR 39443, July 23, 1997; 68 FR 24879, May 9, 2003; 70 FR 69439, Nov. 16, 2005; 73 FR 76946, Dec. 18, 2008; 75 FR 76259, Dec. 8, 2010; 76 FR 59024, Sept. 23, 2011; 77 FR 29217, May 17, 2012; 79 FR 37620, July 2, 2014; 79 FR 53136, Sept. 8, 2014; 79 FR 64116, Oct. 28, 2014; 80 FR 34278, June 16, 2015; 83 FR 14587, Apr. 5, 2018; 84 FR 8973, Mar. 13, 2019; 84 FR 32992, July 11, 2019; 87 FR 76421, Dec. 14, 2022; 88 FR 16547, Mar. 20, 2023]

#### § 520.2645 Tylvalosin.

(a) *Specifications*. Granules containing 62.5 percent tylvalosin (w/w) as tylvalosin tartrate.

(b) *Sponsor*. See No. 066916 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.748 of this chapter.

(d) *Conditions of use in swine*—(1) *Amount*. Administer 50 parts per million (ppm) tylvalosin continuously in drinking water for 5 consecutive days.

(2) *Indications for use*. For control of porcine proliferative enteropathy (PPE) associated with *Lawsonia intracellularis* infection in groups of swine intended for slaughter and female swine intended for breeding in buildings experiencing an outbreak of PPE; and for control of swine respiratory disease (SRD) associated with *Bordetella bronchiseptica*, *Glaesserella (Haemophilus) parasuis*, *Pasteurella multocida*, *Streptococcus suis*, and *Mycoplasma hyopneumoniae* in groups of swine intended for slaughter and female swine intended for breeding in buildings experiencing an outbreak of SRD.

(3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[77 FR 55415, Sept. 10, 2012, as amended at 83 FR 13635, Mar. 30, 2018; 86 FR 57997, Oct. 20, 2021; 89 FR 14410, Feb. 27, 2024]

#### § 520.2654 Velagliflozin.

(a) *Specifications*. Each milliliter of solution contains 15 milligrams (mg) velagliflozin.

(b) *Sponsor*. See No. 000010 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. Administer orally 0.45 mg per pound of body weight (1 mg per kilogram) velagliflozin once daily.

(2) *Indications for use*. To improve glycemic control in otherwise healthy cats with diabetes mellitus not previously treated with insulin.

(3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[88 FR 84700, Dec. 6, 2023]

### PART 522—IMPLANTATION OR INJECTABLE DOSAGE FORM NEW ANIMAL DRUGS

Sec.	
522.23	Acepromazine.
522.52	Alfaxalone.
522.56	Amikacin.
522.62	Aminopentamide.
522.82	Aminopropazine.
522.88	Amoxicillin.
522.90	Ampicillin injectable dosage forms.
522.90a	Ampicillin trihydrate suspension.
522.90b	Ampicillin trihydrate powder for injection.
522.90c	Ampicillin sodium.
522.144	Arsenamides.
522.147	Atipamezole.
522.150	Azaperone.
522.158	Bedinvetmab.
522.161	Betamethasone.
522.163	Betamethasone dipropionate and betamethasone sodium phosphate aqueous suspension.
522.167	Betamethasone sodium phosphate and betamethasone acetate.
522.204	Boldenone.
522.224	Bupivacaine.
522.230	Buprenorphine.
522.234	Butamisolol.
522.246	Butorphanol.
522.275	N-Butylscopolammonium.
522.304	Carprofen.
522.311	Cefovecin.
522.313	Ceftiofur injectable dosage forms.
522.313a	Ceftiofur crystalline free acid.