

§ 520.903d

(2) *Indications for use.* (i) Dogs and puppies: For removal of hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*), whipworms (*Trichuris vulpis*), ascarids (*Toxocara canis* and *Toxascaris leonina*), and tapeworms (*Dipylidium caninum* and *Taenia pisiformis*).

(ii) Cats and kittens: For removal of hookworms (*Ancylostoma tubaeforme*), ascarids (*Toxocara cati*) and tapeworms (*Dipylidium caninum* and *Taenia taeniaeformis*).

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 19167, May 7, 1985, as amended at 53 FR 48533, Dec. 1, 1988; 56 FR 50813, Oct. 9, 1991; 79 FR 28821, May 20, 2014. Redesignated at 85 FR 18119, Apr. 1, 2020; 86 FR 14818, Mar. 19, 2021]

§ 520.903d Febantel tablets.

(a) *Specifications.* Each scored tablet contains 27.2 milligrams of febantel for use in dogs, puppies, cats, and kittens or 163.3 milligrams of febantel for use in dogs, puppies, and cats.

(b) *Sponsor.* See No. 058198 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount—(i) Dogs and cats.* Ten milligrams per kilogram body weight. Administer once daily for 3 consecutive days.

(ii) *Puppies and kittens fewer than 6 months of age.* Fifteen milligrams per kilogram body weight. Administer once daily for 3 consecutive days.

(2) *Indications for use.* (i) For removal of hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*), ascarids (*Toxocara canis* and *Toxascaris leonina*) and whipworms (*Trichuris vulpis*) in dogs and puppies.

(ii) For removal of hookworms (*Ancylostoma tubaeforme*) and ascarids (*Toxocara cati*) in cats and kittens.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[56 FR 50655, Oct. 8, 1991, as amended at 79 FR 28821, May 20, 2014. Redesignated at 85 FR 18119, Apr. 1, 2020, as amended at 86 FR 14818, Mar. 19, 2021]

21 CFR Ch. I (4–1–25 Edition)

§ 520.905 Fenbendazole oral dosage forms.

§ 520.905a Fenbendazole suspension.

(a) *Specifications.* Each milliliter of suspension contains 100 milligrams (mg) fenbendazole for use as in paragraphs (e)(1), (2), (3), and (4) of this section; or 200 mg fenbendazole for use as in paragraphs (e)(5) and (6) of this section.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.275 of this chapter.

(d) *Special considerations.* (1) See § 500.25 of this chapter.

(2) Fenbendazole suspension 10 percent and approved forms of trichlorfon, when used concomitantly for treating the indications provided in paragraph (e) of this section and for treating infections of stomach bot as provided in § 520.2520, have been shown to be compatible and not to interfere with one another.

(e) *Conditions of use—(1) Horses—(i) Amount.* Administer orally 5 mg per kilogram (/kg) (2.3 mg per pound (/lb)) for the control of large strongyles, small strongyles, and pinworms; 10 mg/kg for the control of ascarids.

(ii) *Indications for use.* For the treatment and control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*, *Triodontophorus* species), small strongyles (*Cyathostomum* species, *Cylicocycclus* species, *Cylicostephanus* species, *Cylicodontophorus* species), pinworms (*Oxyuris equi*) and ascarids (*Parascaris equorum*).

(iii) *Limitations.* Do not use in horses intended for human consumption.

(2) *Beef and dairy cattle—(i) Amount.* Administer orally 2.3 mg/lb of body weight (5 mg/kg).

(ii) *Indications for use.* For the treatment and control of: Lungworms: Adult (*Dictyocaulus viviparus*); Stomach worms: Adult brown stomach worms (*Ostertagia ostertagi*); adult and fourth-stage larvae barberpole worms (*Haemonchus contortus* and *H. placei*); adult and fourth-stage larvae small stomach worms (*Trichostrongylus axei*); Intestinal worms (adult and fourth-stage larvae): Hookworms (*Bunostomum phlebotomum*), thread-necked intestinal worms (*Nematodirus helvetianus*), small