

§ 522.480

(ii) *Indications for use.* Control of early mortality associated with *Escherichia coli* organisms susceptible to colistin.

(iii) *Limitations.* For subcutaneous injection in the neck of 1- to 3-day-old chickens. Not for use in laying hens producing eggs for human consumption. Do not use in turkeys. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[63 FR 13123, Mar. 18, 1998, as amended at 79 FR 16185, Mar. 25, 2014; 84 FR 32992, July 11, 2019]

§ 522.480 Corticotropin.

(a) *Specifications.* Each milliliter of aqueous solution contains 40 or 80 U.S.P. (I.U.) units of repository corticotropin.

(b) *Sponsor.* See sponsors in § 510.600(c) of this chapter.

(1) No. 061133 for use as in paragraphs (c)(1) and (2) of this section.

(2) No. 043264 for use as in paragraph (c)(2) and (3) of this section.

(c) *Conditions of use*—(1) *Dogs*—(i) *Amount.* Administer one unit per pound of body weight by intramuscular injection.

(ii) *Indications for use.* As a diagnostic aid to test for adrenal dysfunction.

(iii) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Dogs and cats*—(i) *Amount.* Administer one unit per pound of body weight by intramuscular or subcutaneous injection, to be repeated as indicated.

(ii) *Indications for use.* For stimulation of the adrenal cortex where there is a general deficiency of corticotropin (ACTH).

(iii) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(3) *Cattle*—(i) *Amount.* Administer 200 to 600 units by intramuscular or subcutaneous injection as an initial dose, followed by a dose daily or every other day of 200 to 300 units.

(ii) *Indications for use.* As a therapeutic agent for primary bovine ketosis; and for stimulation of the adrenal cortex where there is a general deficiency of ACTH.

21 CFR Ch. I (4–1–25 Edition)

(iii) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[79 FR 16185, Mar. 25, 2014, as amended at 84 FR 8973, Mar. 13, 2019; 85 FR 45308, July 28, 2020]

§ 522.522 Danofloxacin.

(a) *Specifications.* Each milliliter of solution contains 180 milligrams (mg) danofloxacin as the mesylate salt.

(b) *Sponsor.* See No. 054771 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.169 of this chapter.

(d) *Conditions of use in cattle*—(1) *Amount and indications for use.* Administer by subcutaneous injection either:

(i) 6 mg per kilogram (/kg) of body weight, repeated in 48 hours, for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica* and *Pasteurella multocida*; or

(ii) 8 mg/kg of body weight as a single dose for the treatment of BRD associated with *M. haemolytica* and *P. multocida* and for the control of BRD in beef cattle at high risk of developing BRD associated with *M. haemolytica* and *P. multocida*.

(2) *Limitations.* Animals intended for human consumption should not be slaughtered within 4 days from the last treatment. Do not use in cattle intended for dairy production. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Federal law restricts this drug to use by or on the order of a licensed veterinarian. Federal law prohibits the extra-label use of this drug in food-producing animals.

[67 FR 78972, Dec. 27, 2002, as amended at 77 FR 4227, Jan. 27, 2012; 79 FR 16185, Mar. 25, 2014; 79 FR 53136, Sept. 8, 2014]

§ 522.533 Deslorelin.

(a) *Specifications.* (1) Each implant contains 2.1 milligrams (mg) deslorelin acetate.

(2) Each milliliter (mL) of suspension contains 1.8 mg deslorelin acetate.

(b) *Sponsors.* See sponsor numbers in § 510.600(c) of this chapter as follows:

(1) No. 051311 for use of product described in paragraph (a)(1) as in paragraph (c)(1) of this section.