

§ 556.375

§ 556.375 Maduramicin.

- (a) [Reserved]
- (b) *Tolerances.* The tolerance for maduramicin (marker residue) is:
 - (1) *Chickens.* Fat (target tissue): 0.38 ppm.
 - (2) [Reserved]
- (c) *Related conditions of use.* See § 558.340 of this chapter.

§ 556.380 Melengestrol.

- (a) [Reserved]
- (b) *Tolerances.* The tolerance for melengestrol is:
 - (1) *Cattle.* Fat: 25 ppb.
 - (2) [Reserved]
- (c) *Related conditions of use.* See § 558.342 of this chapter.

§ 556.410 Metoserpate.

- (a) [Reserved]
- (b) *Tolerances.* The tolerance for metoserpate is:
 - (1) *Chickens.* Edible tissues (excluding eggs): 0.02 ppm.
 - (2) [Reserved]
- (c) *Related conditions of use.* See § 520.1422 of this chapter.

§ 556.420 Monensin.

- (a) *Acceptable daily intake (ADI).* The ADI for total residue of monensin is 12.5 µg/kg of body weight per day.
- (b) *Tolerances.* The tolerances for monensin are:
 - (1) *Cattle.* (i) Liver: 0.10 ppm.
(ii) Muscle, kidney, and fat: 0.05 ppm.
(iii) Milk: Not required.
 - (2) *Chickens and turkeys.* Edible tissues (excluding eggs): Not required.
 - (3) *Goats.* Edible tissues (excluding milk): 0.05 ppm.
 - (4) *Quail.* Edible tissues (excluding eggs): Not required.
- (c) *Related conditions of use.* See § 558.355 of this chapter.

§ 556.425 Morantel.

- (a) *Acceptable daily intake (ADI).* The ADI for total residue of morantel tartrate is 10 µg/kg of body weight per day.
- (b) *Tolerances.* The tolerances for N-methyl-1,3-propanediamine (marker residue) are:
 - (1) *Cattle.* (i) Liver (target tissue): 0.7 ppm.
(ii) Milk: Not required.

21 CFR Ch. I (4–1–25 Edition)

- (2) *Goats.* (i) Liver (target tissue): 0.7 ppm.
(ii) Milk: Not required.
- (c) *Related conditions of use.* See §§ 520.1450a, 520.1450b, 520.1450c, and 558.360 of this chapter.

§ 556.426 Moxidectin.

- (a) *Acceptable daily intake (ADI).* The ADI for total residue of moxidectin is 4 µg/kg of body weight per day.
- (b) *Tolerances.* The tolerances for moxidectin (marker residue) are:
 - (1) *Cattle.* (i) Fat (target tissue): 900 ppb.
(ii) Liver: 200 ppb.
(iii) Muscle: 50 ppb.
(iv) Milk: 40 ppb.
 - (2) *Sheep.* (i) Fat (target tissue): 900 ppb.
(ii) Liver: 200 ppb.
(iii) Muscle: 50 ppb.
- (c) *Related conditions of use.* See §§ 520.1454, 522.1450, and 524.1450 of this chapter.

§ 556.428 Narasin.

- (a) *Acceptable daily intake (ADI).* The ADI for total residue of narasin is 5 µg/kg of body weight per day.
- (b) *Tolerances.* The tolerance for narasin (marker residue) is:
 - (1) *Chickens.* Abdominal fat (target tissue): 480 ppb.
 - (2) [Reserved]
- (c) *Related conditions of use.* See §§ 558.363 and 558.364 of this chapter.

§ 556.430 Neomycin.

- (a) *Acceptable daily intake (ADI).* The ADI for total residue of neomycin is 6 µg/kg of body weight per day.
- (b) *Tolerances.* The tolerances for neomycin are:
 - (1) *Cattle.* (i) Kidney (target tissue): 7.2 ppm.
(ii) Liver: 3.6 ppm.
(iii) Muscle: 1.2 ppm.
(iv) Fat: 7.2 ppm.
(v) Milk: 0.15 ppm.
 - (2) *Sheep and goats.* (i) Kidney (target tissue): 7.2 ppm.
(ii) Liver: 3.6 ppm.
(iii) Muscle: 1.2 ppm.
(iv) Fat: 7.2 ppm.
(v) Milk: 0.15 ppm.
 - (3) *Swine.* (i) Kidney (target tissue): 7.2 ppm.
(ii) Liver: 3.6 ppm.