

§ 556.36 Altrenogest.

(a) *Acceptable daily intake (ADI)*. The ADI for total residue of altrenogest is 0.04 µg/kg of body weight per day.

(b) *Tolerances*. The tolerances for altrenogest (marker residue) are:

(1) *Swine*. (i) Liver (target tissue): 4 ppb.

(ii) Muscle: 1 ppb.

(2) [Reserved]

(c) *Related conditions of use*. See § 520.48 of this chapter.

§ 556.38 Amoxicillin.

(a) [Reserved]

(b) *Tolerances*. The tolerance for amoxicillin is:

(1) *Cattle*. Edible tissues: 0.01 ppm.

(2) [Reserved]

(c) *Related conditions of use*. See §§ 520.88d, 520.88e, 522.88, and 526.88 of this chapter.

[84 FR 32993, July 11, 2019, as amended at 86 FR 13188, Mar. 8, 2021]

§ 556.40 Ampicillin.

(a) [Reserved]

(b) *Tolerances*. The tolerances for ampicillin are:

(1) *Cattle*. Edible tissues: 0.01 ppm.

(2) *Swine*. Edible tissues: 0.01 ppm.

(c) *Related conditions of use*. See §§ 520.90c, 522.90a, and 522.90b of this chapter.

[84 FR 32993, July 11, 2019, as amended at 85 FR 18120, Apr. 1, 2020]

§ 556.50 Amprolium.

(a) [Reserved]

(b) *Tolerances*. The tolerances for amprolium are:

(1) *Cattle*. (i) Liver, kidney, and muscle: 0.5 ppm.

(ii) Fat: 2.0 ppm.

(2) *Chickens and turkeys*. (i) Liver and kidney: 1 ppm.

(ii) Muscle: 0.5 ppm.

(iii) Eggs:

(A) Egg yolks: 8 ppm.

(B) Whole eggs: 4 ppm.

(3) *Pheasants*. (i) Liver: 1 ppm.

(ii) Muscle: 0.5 ppm.

(c) *Related conditions of use*. See §§ 520.100, 558.55, and 558.58 of this chapter.

§ 556.52 Apramycin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residue of apramycin is 25 µg/kg of body weight per day.

(b) *Tolerances*. The tolerance for apramycin (marker residue) is:

(1) *Swine*. Kidney (target tissue): 0.1 ppm.

(2) [Reserved]

(c) *Related conditions of use*. See §§ 520.110 and 558.59 of this chapter.

§ 556.60 Avilamycin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residue of avilamycin is 1.1 mg/kg of body weight per day.

(b) *Tolerances*. The tolerances for avilamycin are:

(1) *Chickens*. Edible tissues (excluding eggs): Not required.

(2) *Swine*. Edible tissues: Not required.

(c) *Related conditions of use*. See § 558.68 of this chapter.

§ 556.68 Azaperone.

(a) *Acceptable daily intake (ADI)*. The ADI for total residue of azaperone is 0.63 µg/kg of body weight per day.

(b) *Tolerances*. The tolerance for azaperone is:

(1) *Swine*. Edible tissues: Not required.

(2) [Reserved]

(c) *Related conditions of use*. See § 522.150 of this chapter.

§ 556.70 Bacitracin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residue of bacitracin is 0.05 mg/kg of body weight per day.

(b) *Tolerances*. The tolerances for bacitracin are:

(1) *Cattle*. Edible tissues: 0.5 ppm.

(2) *Chickens, turkeys, pheasants, quail*. Edible tissues: 0.5 ppm.

(3) *Swine*. Edible tissues: 0.5 ppm.

(c) *Related conditions of use*. See §§ 520.154a, 520.154c, 558.76, and 558.78 of this chapter.

§ 556.75 Bambermycins.

(a) [Reserved]

(b) *Tolerances*. The tolerances for bambermycins are:

(1) *Cattle*. Edible tissues (excluding milk): Not required.

(2) *Chickens and turkeys*. Edible tissues (excluding eggs): Not required.