

Diclazuril grams/ton	Combination grams/ton	Indications for use	Limitations	Sponsor
(iii) 0.91 .....	Bambermycins 1 to 2.	Growing turkeys: For the prevention of coccidiosis caused by <i>Eimeria adenoides</i> , <i>E. gallopavonis</i> , and <i>E. meleagrimitis</i> , and for improved feed efficiency.	Feed continuously as the sole ration. Do not feed to breeding turkeys. Do not feed to birds producing eggs for human consumption. Bambermycins as provided by No. 016592 in § 510.600(c) of this chapter.	058198
(iv) 0.91 .....	Bambermycins 2 ..	Growing turkeys: For the prevention of coccidiosis caused by <i>Eimeria adenoides</i> , <i>E. gallopavonis</i> , and <i>E. meleagrimitis</i> , and for increased rate of weight gain and improved feed efficiency.	Feed continuously as the sole ration. Do not feed to breeding turkeys. Do not feed to birds producing eggs for human consumption. Bambermycins as provided by No. 016592 in § 510.600(c) of this chapter.	058198

(3) Diclazuril may also be used in combination with virginiamycin as in § 558.635.

[64 FR 35923, July 2, 1999, as amended at 65 FR 50134, Aug. 17, 2000; 66 FR 47962, 47963, Sept. 17, 2001; 66 FR 62917, Dec. 4, 2001; 67 FR 34830, May 16, 2002; 67 FR 47257, July 18, 2002; 67 FR 48549, July 25, 2002; 69 FR 9947, Mar. 3, 2004; 72 FR 60552, Oct. 25, 2007; 79 FR 10982, Feb. 27, 2014; 79 FR 13545, Mar. 11, 2014; 81 FR 17609, Mar. 30, 2016; 81 FR 95004, Dec. 27, 2016. Redesignated and amended at 84 FR 12497, 12498, Apr. 2, 2019; 87 FR 10970, Feb. 28, 2022]

**§ 558.235 Efrotomycin.**

(a) *Specifications.* Type A medicated articles containing 14.5 grams efrotomycin per pound.

(b) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.224 of this chapter.

(d) *Conditions of use in swine—*

Efrotomycin in grams/ton	Indications for use	Limitations	Sponsor
(1) 3.6 .....	Swine: For improved feed efficiency .....	Feed continuously as sole ration. Not to be used in swine weighing more than 250 pounds.	000010
(2) 3.6 to 14.5 .....	Swine: For increased rate of weight gain	Feed continuously as sole ration. Not to be used in swine weighing more than 250 pounds.	000010

[57 FR 38442, Aug. 25, 1992, as amended at 62 FR 63271, Nov. 28, 1997; 84 FR 33001, July 11, 2019; 84 FR 39185, Aug. 9, 2019; 85 FR 45309, July 28, 2020]

**§ 558.248 Erythromycin.**

(a) *Specifications.* Type A medicated articles containing 92.5 grams per pound erythromycin (as the thiocyanate salt).

(b) *Sponsor.* See No. 061133 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.230 of this chapter.

(d) *Special considerations.* (1) Federal law restricts medicated feed containing

this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian. See § 558.6 for additional requirements.

(2) The expiration date of VFDs for erythromycin medicated feeds must not exceed 6 months from the date of issuance. VFDs for erythromycin shall not be refilled.

(e) *Conditions of use—(1) Chickens—*

§ 558.254

21 CFR Ch. I (4–1–25 Edition)

Erythromycin in grams/ton	Combination in grams/ton	Indications for use	Limitations	Sponsor
(i) 92.5 .....	.....	Chickens: As an aid in the prevention of chronic respiratory disease during periods of stress.	Feed for 2 days before stress and 3 to 6 days after stress. Withdraw 24 hours before slaughter.	061623
(ii) 92.5 .....	.....	Chickens: As an aid in the prevention of infectious coryza.	Feed for 7 to 14 days. Withdraw 24 hours before slaughter.	061623
(iii) 185 .....	.....	Chickens: As an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease (CRD).	Feed for 5 to 8 days. Withdraw 48 hours before slaughter. Do not use in birds producing eggs for food.	061623

(2) *Turkeys*—

Erythromycin thiocyanate in grams/ton	Combination in grams/ton	Indications for use	Limitations	Sponsor
(i) 92.5 .....	.....	Turkeys: As an aid in the prevention of chronic respiratory disease during periods of stress.	Feed for 2 days before stress and 3 to 6 days after stress.	061623
(ii) 185 .....	.....	Turkeys: As an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease (CRD).	Feed for 5 to 8 days. Do not use in birds producing eggs for food.	061623

[41 FR 10999, Mar. 15, 1976, as amended at 45 FR 56799, Aug. 26, 1980; 49 FR 31281, Aug. 6, 1984; 51 FR 7397, Mar. 3, 1986; 52 FR 2684, Jan. 26, 1987; 54 FR 12189, Mar. 24, 1989; 66 FR 14074, Mar. 9, 2001; 68 FR 4915, Jan. 31, 2003; 79 FR 10982, Feb. 27, 2014; 81 FR 36790, June 8, 2016; 81 FR 95004, Dec. 27, 2016; 84 FR 8975, Mar. 13, 2019]

§ 558.254 **Famphur.**

(a) *Specifications.* Type A medicated articles containing 13.2 or 33.3 percent famphur.

Famphur amount	Indications for use	Limitations	Sponsor
(1) To provide 1.1 milligrams per pound (mg/lb) body weight per day.	Beef cattle and nonlactating dairy cattle: For control of grubs and as an aid in control of sucking lice.	Feed for 30 days. Withdraw from dry dairy cows and heifers 21 days prior to freshening. Withdraw 4 days prior to slaughter.	000061
(2) To provide 2.3 mg/lb body weight per day.	Beef cattle and nonlactating dairy cattle: For control of grubs.	Feed for 10 days. Withdraw from dry dairy cows and heifers 21 days prior to freshening. Withdraw 4 days prior to slaughter.	000061

[84 FR 39185, Aug. 9, 2019, as amended at 87 FR 17947, Mar. 29, 2022]

§ 558.258 **Fenbendazole.**

(a) *Specifications.* Type A medicated articles: 4 percent (18.1 grams per pound (g/lb)), 8 percent (36.2 g/lb), and 20 percent (90.7 g/lb) fenbendazole.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.273 of this chapter.

(d) *Special considerations.* Famphur is a cholinesterase inhibitor. Do not use this product in animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(e) *Conditions of use.* It is used in cattle feed as follows:

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.275 of this chapter.

(d) *Special considerations.* See § 500.25 of this chapter.

(e) *Conditions of use*—(1) *Turkeys.*