

§ 558.550

21 CFR Ch. I (4–1–25 Edition)

Robenidine hydrochloride in grams/ton	Combination in grams/ton	Indications for use	Limitations	Sponsor
.....	Bacitracin (as bacitracin zinc) 27 to 50.	For broiler and fryer chickens: As an aid in the prevention of coccidiosis caused by <i>E. mivati</i> , <i>E. brunetti</i> , <i>E. tenella</i> , <i>E. acervulina</i> , <i>E. maxima</i> , and <i>E. necatrix</i> . For improved feed efficiency..	Feed continuously as sole ration. Do not feed to laying chickens. Withdraw 5 days prior to slaughter..	054771 054771

(f) Robenidine may also be used in combination with:

- (1) Chlortetracycline as in § 558.128.
- (2) Lincomycin as in § 558.325.
- (3) Oxytetracycline as in § 558.450.

[40 FR 13959, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.515, see the List of CFR Sections Affected, which appears in the Finding Aids section of the printed volume and at [www.govinfo.gov](http://www.govinfo.gov).

§ 558.550 **Salinomycin.**

(a) *Specifications.* Type A medicated articles containing 30 or 60 grams of

salinomycin sodium activity per pound.

(b) *Sponsor.* See No. 016592 in § 510.600(c) of this chapter for use as in paragraph (e) of this section.

(c) *Related tolerances.* See § 556.592 of this chapter.

(d) *Special considerations.* Not approved for use with pellet binders.

(e) *Conditions of use.* It is used as follows:

- (1) *Chickens*—

Salinomycin in grams/ton	Combination in grams/ton	Indications for use	Limitations	Sponsor
(i) 40 to 60 .....	.....	Broiler, roaster, and replacement (breeder and layer) chickens: For the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> , and <i>E. mivati</i> .	Feed continuously as sole ration. Do not feed to birds producing eggs for human consumption. May be fatal if accidentally fed to adult turkeys or horses.	016592
(ii) 40 to 60 .....	Bacitracin methylenedisalicylate, 4 to 50.	Broiler, roaster, and replacement (breeder and layer) chickens: For the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> , and <i>E. mivati</i> , and for increased rate of weight gain and improved feed efficiency.	Feed continuously as sole ration. Do not feed to laying chickens. May be fatal if fed to adult turkeys or horses. Salinomycin as provided by No. 016592; bacitracin methylenedisalicylate as provided by No. 054771 in § 510.600(c) of this chapter.	016592 054771
(iii) 40 to 60 .....	Bacitracin methylenedisalicylate, 50.	Broiler chickens: For the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> , and <i>E. mivati</i> , and as an aid in the prevention of necrotic enteritis caused or complicated by <i>Clostridium</i> spp. or other organisms susceptible to bacitracin.	Feed continuously as sole ration. Do not feed to laying chickens. May be fatal if fed to adult turkeys or to horses. Salinomycin as provided by No. 016592; bacitracin methylenedisalicylate as provided by No. 054771 in § 510.600(c) in this chapter.	054771

Salinomycin in grams/ton	Combination in grams/ton	Indications for use	Limitations	Sponsor
(iv) 40 to 60 .....	Bacitracin methylenedisalicylate, 100 to 200.	Broiler chickens: For the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> , and <i>E. mivati</i> , and as an aid in the control of necrotic enteritis caused or complicated by <i>Clostridium</i> spp. or other organisms susceptible to bacitracin.	Feed continuously as sole ration. To control necrotic enteritis, start medication at first clinical signs of disease; vary dosage based on the severity of infection; administer continuously for 5 to 7 days or as long as clinical signs persist, then reduce bacitracin to prevention level (50 grams per ton). Do not feed to laying chickens. May be fatal if fed to adult turkeys or to horses. Salinomycin as provided by No. 016592; bacitracin methylenedisalicylate as provided by No. 054771 in §510.600(c) in this chapter.	054771
(v) 40 to 60 .....	Bacitracin zinc, 10 to 50.	Broiler chickens: For the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> , and <i>E. mivati</i> , and for increased rate of weight gain.	Feed continuously as sole ration. Not approved for use with pellet binders. Do not feed to layers. May be fatal if accidentally fed to adult turkeys or horses. Bacitracin zinc as provided by No. 054771 in §510.600(c) of this chapter.	016592 054771
(vi) 40 to 60 .....	Bambermycins, 1 to 3.	Broiler chickens: For the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> , and <i>E. mivati</i> , and for improved feed efficiency.	Feed continuously as sole ration. Do not feed to laying chickens. Not approved for use with pellet binders. May be fatal if accidentally fed to adult turkeys or horses. Salinomycin and bambermycins as provided by No. 016592 in §510.600(c) in this chapter.	016592

(2) Game birds—

Salinomycin in grams/ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 50 .....	.....	Quail: For the prevention of coccidiosis caused by <i>Eimeria dispersa</i> and <i>E. lettyae</i> .	Feed continuously as sole ration. Do not feed to birds producing eggs for human consumption. May be fatal if accidentally fed to adult turkeys or horses.	016592
(ii) [Reserved] .....	.....	.....	.....	.....

(3) *Combinations.* Salinomycin may also be used in combination with:

- (i) Avilamycin as in § 558.68.
- (ii) Chlortetracycline as in § 558.128.
- (iii) Lincomycin as in § 558.325.
- (iv) Oxytetracycline as in § 558.450.
- (v) Virginiamycin as in § 558.635.

[48 FR 30616, July 5, 1983]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.550, see the List of CFR Sections Affected, which appears in the Finding Aids section of the printed volume and at [www.govinfo.gov](http://www.govinfo.gov).

§ 558.555 Semduramicin.

(a) *Specifications.* Type A medicated article containing:

(1) 22.7 grams (g) per pound (lb) (50 g/kilogram (kg)) semduramicin (as semduramicin sodium).

(2) 22.7 g/lb (50 g/kg) semduramicin (as semduramicin sodium biomass).

(b) *Sponsor.* See No. 066104 in § 510.600(c) of this chapter for use of product described in paragraph (a)(1) of this section as in paragraph (d) of this section; for use of product described in