

§ 573.696

21 CFR Ch. I (4–1–25 Edition)

broiler chicken feeds in accordance with the following specifications:

(a) The additive is a stereoisomer of 22-[(3-amino-3,6-dideoxy-β-D-mannopyranosyl)oxy]-1,3,26-trihydroxy-12-methyl-10-oxo-6,11,28-trioxatricyclo[22.3.1.0^{5,7}] octacos-8,14,16,18,20-pentaene-25-carboxylic acid with the empirical formula C₃₃H₄₇NO₁₃.

(b) The additive shall conform to U.S.P. specifications.

(c) The additive (as part of a premix composed of calcium carbonate, natamycin, and lactose) is used for retarding the growth of *Aspergillus parasiticus* in broiler chicken feeds for up to 14 days after the addition of natamycin.

(d) Each pound (454 grams (g)) of the premix shall contain 434 (g) of calcium carbonate, 10 g of natamycin activity, and 10 g of lactose. The premix shall be mixed into broiler chicken feed at the rate of 1 pound (0.454 kilograms (kg)) per ton (908 kg) of feed to provide natamycin at a level of 11 parts per million (ppm). The premix shall be thoroughly mixed into the dry components of the broiler chicken feed before adding the liquid components. Broiler feeds to which the natamycin premix is added shall be used within 4 weeks of addition of the premix.

(e) To assure the safe use of the additive, the label or labeling of the additive shall bear, in addition to other information required by the Federal Food, Drug, and Cosmetic Act, the following:

(1) The name and CAS number of the additive, and its purpose.

(2) A listing of ingredients consisting of calcium carbonate, the additive, and lactose and their proportions in the premix as prescribed under paragraph (d) of this section.

(3) Adequate directions for use to ensure a broiler chicken feed that is in compliance with the limitations prescribed in paragraph (d) of this section.

(4) An appropriate cautionary statement: "Caution: Store in a tightly-closed, light-resistant container in a cool, dry place."

(5) An expiration date of 1 year from the date of manufacture.

(6) A contact address and telephone number for reporting adverse reactions experienced by users, or to request a

copy of the Material Safety Data Sheet for natamycin.

[69 FR 19321, Apr. 13, 2004]

§ 573.696 Feed grade sodium formate.

The food additive, feed grade sodium formate, may be safely used in the manufacture of complete swine and poultry feeds in accordance with the following prescribed conditions:

(a) The additive is manufactured by the reaction of 99 percent formic acid and 50 percent sodium hydroxide in water to produce a solution made up of at least 20.5 percent sodium salt of formic acid and not more than 61 percent formic acid.

(b) The additive is used or intended for use as a feed acidifying agent, to lower the pH, in complete swine and poultry feeds at levels not to exceed 1.2 percent of the complete feed.

(c) To assure safe use of the additive, formic acid and formate salts from all added sources cannot exceed 1.2 percent of complete feed when multiple sources of formic acid and its salts are used in combination.

(d) To assure safe use of the additive, in addition to the other information required by the Federal Food, Drug, and Cosmetic Act, the label and labeling shall contain:

(1) The name of the additive.

(2) Adequate directions for use, including a statement that feed grade sodium formate must be uniformly applied and thoroughly mixed into complete feeds and that the complete feeds so treated shall be labeled as containing feed grade sodium formate.

(3) Cautions for use including this statement: Caution: Follow label directions. Formic acid and formate salts from all added sources cannot exceed 1.2 percent of complete feed when multiple sources of formic acid and its salts are used in combination.

(e) To assure safe use of the additive, in addition to the other information required by the act and paragraph (d) of this section, the label and labeling shall contain:

(1) Appropriate warnings and safety precautions concerning feed grade sodium formate.

(2) Statements identifying feed grade sodium formate as a corrosive and possible severe irritant.

Food and Drug Administration, HHS

§ 573.760

(3) Information about emergency aid in case of accidental exposure as follows:

(i) Statements reflecting requirements of applicable sections of the Superfund Amendments and Reauthorization Act (SARA), and the Occupational Safety and Health Administration (OSHA) human safety guidance regulations.

(ii) Contact address and telephone number for reporting adverse reactions or to request a copy of the Material Safety Data Sheet (MSDS).

[81 FR 67154, Sept. 30, 2016, as amended at 81 FR 95027, Dec. 27, 2016]

§ 573.700 Sodium nitrite.

Sodium nitrite may be safely used in canned pet food containing meat and fish in accordance with the following prescribed conditions:

(a) It is used or intended for use alone as a preservative and color fixative in canned pet food containing fish, meat, and fish and meat byproducts so that the level of sodium nitrite does not exceed 20 parts per million.

(b) To assure safe use of the additive, in addition to the other information required by the act:

(1) The label of the additive shall bear:

(i) The name of the additive.

(ii) A statement of the concentration of the additive in any mixture.

(2) The label or labeling shall bear adequate directions to provide a final product that complies with the limitations prescribed in paragraph (a) of this section.

§ 573.720 Petrolatum.

Petrolatum may be safely used in or on animal feed, subject to the following prescribed conditions:

(a) Petrolatum complies with the specifications set forth in the U.S. Pharmacopeia XVI for white petrolatum or in The National Formulary XII for yellow petrolatum.

(b) Petrolatum meets the following ultraviolet absorbance limits when subjected to the analytical procedure described in §172.886(b) of this chapter.

Ultraviolet absorbance per centimeter path length:

Millimicrons	Maximum
280 to 289	0.25
290 to 29920
300 to 35914
360 to 40004

(c) It is used in animal feed for the following purposes:

(1) To reduce dustiness of feeds or mineral supplements.

(2) To serve as a lubricant in the preparation of pellets, cubes, or blocks, and to improve resistance to moisture of such pellets, cubes, or blocks.

(d) The quantity of petrolatum present in animal feeds from the uses specified in paragraph (c) of this section shall not exceed 3 percent in mineral supplements nor shall it exceed 0.06 percent of the total ration when present in feed or feed concentrates.

(e) When used in combination with technical white mineral oil for the uses described in paragraph (c) of this section, the total quantity of combined petrolatum and technical white mineral oil shall not exceed the limits prescribed in paragraph (d) of this section.

(f) Petrolatum may contain any antioxidant permitted in food by regulations issued in accordance with section 409 of the act, in an amount not greater than that required to produce its intended effect.

§ 573.740 Odorless light petroleum hydrocarbons.

Odorless light petroleum hydrocarbons complying with §172.884(a) and (b) of this chapter may be safely used in an amount not in excess of that required as a component of insecticide formulations used in compliance with regulations issued in this part.

§ 573.760 Poloxalene.

The food additive poloxalene may be safely used in accordance with the following prescribed conditions:

(a) The additive consists of polyoxypropylene-polyoxyethylene glycol non-ionic block polymer meeting the following specifications:

(1) Molecular weight range: 2,850–3,150.

(2) Hydroxyl number: 35.7–39.4.

(3) Cloud point (10 percent solution): 42 °C–46 °C.

(4) Structural formula: