§ 573.940

- (ii) Minimum and maximum guarantees for a total selenium content of not less than 2.08 percent (weight/weight) and not more than 2.24 percent;
- (iii) Minimum guarantee for selenomethionine hydroxy analogue content of not less than 5.2 percent;
- (iv) The following statement, "Storage Conditions: Selenomethionine hydroxy analogue must be stored in a closed package at temperatures not higher than 20 °C (68 °F)."; and
- (v) An expiration date not to exceed 1 year from the date of manufacture.
- (4) The additive, as selenomethionine hydroxy analogue, shall be incorporated into each ton of complete feed by adding no less than 1 pound of a premix containing no more than 272.4 milligrams of added selenium per pound.
- (5) Usage of this additive must conform to the requirements of paragraphs (d) and (e) of this section.

[52 FR 10888, Apr. 6, 1987; 52 FR 21001, June 4, 1987, as amended at 54 FR 14215, Apr. 10, 1989; 54 FR 15874, Apr. 19, 1989; 60 FR 53703, Oct. 17, 1995; 65 FR 35824, June 6, 2000; 65 FR 53167, Sept. 1, 2000; 67 FR 46851, July 17, 2002; 68 FR 52340, Sept. 3, 2003; 72 FR 39561, July 19, 2007; 84 FR 7993, Mar. 6, 2019; 86 FR 37036, July 14, 2021

EFFECTIVE DATE NOTE: At 58 FR 47973, Sept. 13, 1993, the amendments to §573.920 that were published at 52 FR 10887, Apr. 6, 1987; 52 FR 21001, June 4, 1987; and 54 FR 14214, Apr. 10, 1989 were stayed until further notice. At 59 FR 45973, Sept. 6, 1994, the stay was confirmed.

§ 573.940 Silicon dioxide.

The food additive silicon dioxide may be safely used in animal feed in accordance with the following conditions:

- (a) The food additive is manufactured by vapor phase hydrolysis or by other means whereby the particle size is such as to accomplish the intended effect.
- (b) It is used or intended for use as an anticaking agent, antifoaming agent, carrier, and/or grinding aid in animal feed, including ingredients, intermediate premixes, premixes, supplements, concentrates, and complete feed
- (c) To ensure safe use of the additive, silicon dioxide is to be used in an amount not to exceed that reasonably required to accomplish its intended effect, and silicon dioxide from all

sources cannot exceed 2 percent by weight of the complete feed.

- (d) To ensure safe use of the additive, the label and labeling of the additive and ingredients, intermediate premixes, premixes, supplements, concentrates, and complete feed containing the additive shall meet the requirements of the Federal Food, Drug, and Cosmetic Act, including part 501 of this chapter.
- (e) To ensure safe use of the additive, in addition to the other information required by the Federal Food, Drug, and Cosmetic Act, the label and labeling of the additive and ingredients, intermediate premixes, premixes, supplements, and concentrates containing the additive shall have:
- (1) A statement of the concentration of the additive.
- (2) A statement that silicon dioxide from all sources cannot exceed 2 percent by weight of the complete feed.
- [41 FR 38652, Sept. 10, 1976, as amended at 83 FR 8930, Mar. 2, 2018; 84 FR 7993, Mar. 6, 2019; 85 FR 33539, June 2, 2020]

§573.960 Sorbitan monostearate.

The food additive sorbitan monostearate may be safely used alone or in combination with polysorbate 60 as an emulsifier in mineral premixes and dietary supplements for animal feeds.

§ 573.980 Taurine.

The food additive taurine (2-aminoethanesulfonic acid) may be safely used in feed in accordance with the following prescribed conditions:

- (a) It is used as a nutritional supplement in the feed of growing chickens.
- (b) It is added to complete feeds so that the total taurine content does not exceed 0.054 percent of the feed.
- (c) To assure safe use of the additive, the label and labeling shall bear in addition to the other information required by the Act:
 - (1) The name of the additive.
- (2) The quantity of the additive contained therein.
 - (3) Adequate directions for use.

§ 573.1000 Verxite.

The food additive verxite may be safely used in animal feed in accordance with the following prescribed conditions: