

Food and Drug Administration, HHS

§ 640.31

by number so as to relate it to the donor.

[40 FR 4304, Jan. 29, 1975, as amended at 42 FR 10983, Feb. 25, 1977; 47 FR 49021, Oct. 29, 1982; 50 FR 4139, Jan. 29, 1985; 63 FR 16685, Apr. 6, 1998; 64 FR 45372, Aug. 19, 1999; 66 FR 1836, Jan. 10, 2001; 66 FR 40890, Aug. 6, 2001; 72 FR 45887, Aug. 16, 2007; 73 FR 7464, Feb. 8, 2008]

§ 640.25 General requirements.

(a) *Storage.* Immediately after resuspension, Platelets shall be placed in storage at the selected temperature range. If stored at 20 to 24 °C, a continuous gentle agitation of the platelet concentrate shall be maintained throughout the storage period. Agitation is optional if stored at a temperature between 1 and 6 °C.

(b) *Quality control testing.* Each month four units prepared from different donors shall be tested at the end of the storage period as follows:

(1) Platelet count.

(2) pH of not less than 6.2 measured at the storage temperature of the unit.

(3) Measurement of actual plasma volume.

(4) If the results of the quality control testing indicate that the product does not meet the prescribed requirements, immediate corrective action shall be taken and a record maintained of such action.

(c) *Manufacturing responsibility.* All manufacturing of Platelets shall be performed at the same licensed establishment, except that the quality control testing under paragraph (b) of this section may be performed by a clinical laboratory which meets the standards of the Clinical Laboratories Improvement Amendments of 1988 (CLIA) (42 U.S.C. 263a) and is qualified to perform platelet counts. Such arrangements must be approved by the Director, Center for Biologics Evaluation and Research, Food and Drug Administration. Such testing shall not be considered as divided manufacturing, as described in §610.63 of this chapter, provided the following conditions are met:

(1) The results of each test are received within 10 days of the preparation of the platelet concentrate, and are maintained by the establishment licensed for Platelets so that they may be reviewed by an authorized rep-

resentative of the Food and Drug Administration.

(2) The licensed Platelets manufacturer has obtained a written agreement that the testing laboratory will permit an authorized representative of the Food and Drug Administration to inspect its testing procedures and facilities during reasonable business hours.

(3) The testing laboratory will participate in any proficiency testing programs undertaken by the Center for Biologics Evaluation and Research, Food and Drug Administration.

[40 FR 4304, Jan. 29, 1975, as amended at 47 FR 49021, Oct. 29, 1982; 49 FR 23834, June 8, 1984; 50 FR 4139, Jan. 29, 1985; 55 FR 11013, Mar. 26, 1990; 66 FR 1836, Jan. 10, 2001; 72 FR 45888, Aug. 16, 2007]

Subpart D—Plasma

§ 640.30 Plasma.

(a) *Proper name and definition.* The proper name of this component is Plasma. The component is defined as:

(1) The fluid portion of one unit of human blood intended for intravenous use which is collected in a closed system, stabilized against clotting, and separated from the red cells; or

(2) The fluid portion of human blood intended for intravenous use which is prepared by apheresis methods as specified in the directions for use for the blood collecting, processing, and storage system including closed and open systems.

(b) *Source.* (1) Plasma shall be obtained by separating plasma from blood collected from blood donors or by plasmapheresis.

(2) Plasma may be obtained from a unit of Whole Blood collected by another licensed establishment.

[42 FR 59878, Nov. 22, 1977; 48 FR 13026, Mar. 29, 1983, as amended at 50 FR 4139, Jan. 29, 1985; 72 FR 45888, Aug. 16, 2007]

§ 640.31 Eligibility of donors.

(a) Whole Blood donors must meet the criteria for donor eligibility prescribed in §§630.10 and 630.15 of this chapter.

§ 640.32

(b) Collection establishments must determine the eligibility of plasmapheresis donors in accordance with §§ 630.10 and 630.15 of this chapter.

[80 FR 29904, May 22, 2015]

§ 640.32 Collection of source material.

(a) Whole Blood must be collected, transported, and stored as prescribed in § 640.4. When whole blood is intended for Plasma, Fresh Frozen Plasma, and Liquid Plasma, until the plasma is removed, the whole blood must be maintained at a temperature between 1 and 6 °C or as specified in the directions for use for the blood collecting, processing, and storage system approved for such use by the Director, Center for Biologics Evaluations and Research. Whole blood intended for Platelet Rich Plasma must be maintained as prescribed in § 640.24 until the plasma is removed. The red blood cells must be placed in storage at a temperature between 1 and 6 °C immediately after the plasma is separated.

(b) Plasma obtained by plasmapheresis shall be collected as prescribed in § 640.64 (except that paragraph (c)(3) of § 640.64 shall not apply), and § 640.65.

[42 FR 59878, Nov. 22, 1977, as amended at 45 FR 27927, Apr. 25, 1980; 50 FR 4139, Jan. 29, 1985; 64 FR 45372, Aug. 19, 1999; 72 FR 45888, Aug. 16, 2007; 80 FR 29905, May 22, 2015]

§ 640.33 Testing the blood.

(a) Blood from which plasma is separated shall be tested as prescribed in § 610.40 of this chapter and § 640.5 (b) and (c).

(b) Manufacturers of Plasma collected by plasmapheresis shall have testing and recordkeeping responsibilities equivalent to those prescribed in §§ 640.71 and 640.72.

[42 FR 59878, Nov. 22, 1977, as amended at 44 FR 17658, Mar. 23, 1979; 50 FR 4139, Jan. 29, 1985; 53 FR 117, Jan. 5, 1988; 66 FR 31165, June 11, 2001; 80 FR 29905, May 22, 2015]

§ 640.34 Processing.

(a) *Plasma.* Plasma shall be separated from the red blood cells and shall be stored at -18 °C or colder within 6 hours after transfer to the final container or within the timeframe specified in the directions for use for the

21 CFR Ch. I (4-1-24 Edition)

blood collecting, processing, and storage system unless the product is to be stored as Liquid Plasma.

(b) *Fresh Frozen Plasma.* Fresh frozen plasma shall be prepared from blood collected by a single uninterrupted venipuncture with minimal damage to and minimal manipulation of the donor's tissue. The plasma must be separated from the red blood cells or collected by an apheresis procedure, and placed in a freezer within 8 hours or within the timeframe specified in the directions for use for the blood collecting, processing, and storage system, and stored at -18 °C or colder.

(c) *Liquid Plasma.* Liquid Plasma shall be separated from the red blood cells and shall be stored at a temperature of 1 to 6 °C within 4 hours after filling the final container or within the timeframe specified in the directions for use for the blood collecting, processing, and storage system.

(d) *Platelet Rich Plasma.* Platelet rich plasma shall be prepared from blood collected by a single uninterrupted venipuncture with minimal damage to and manipulation of the donor's tissue. The plasma shall be separated from the red blood cells by centrifugation within 4 hours after completion of the phlebotomy or within the timeframe specified in the directions for use for the blood collecting, processing, and storage system. The time and speed of the centrifugation shall have been shown to produce a product with at least 250,000 platelets per microliter. The plasma shall be stored at a temperature between 20 and 24 °C immediately after filling the final container. A gentle and continuous agitation of the product shall be maintained throughout the storage period, if stored at a temperature of 20 to 24 °C.

(e) *Modifications of Plasma.* It is possible to separate Platelets and/or Cryoprecipitated AHF from Plasma. When these components are to be separated, the plasma shall be collected as described in § 640.32 for Plasma.

(1) Platelets shall be separated as prescribed in subpart C of part 640, prior to freezing the plasma. The remaining plasma may be labeled as